

Determinant

source: <http://physics.stackexchange.com/questions/885/why-does-calculus-of-variations-work>

1 Independence

Why is the Lagrangian a function of two variables? Put differently, you can choose position and velocity independently as initial conditions, that's why the Lagrangian function treats them as independent; but the calculus of variation does not vary them independently, a variation in position induces a fitting variation in velocity.

$$S = \int L(q, \dot{q}, t) dt \quad (1)$$

$$\delta S = \int \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \delta \dot{q} dt = 0 \quad (2)$$

the second step seems suspect; why are q and \dot{q} varied independently?
now

$$\delta \dot{q} = \frac{d}{dt} \delta q \quad (3)$$

the independence of q and \dot{q} is removed by this identity.

$$S = \int L(q, \dot{q}, t) dt \quad (4)$$

$$\delta S = \int \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \delta \dot{q} dt = 0 \quad (5)$$

$$\delta S = \left[\frac{\partial L}{\partial \dot{q}} \delta q \right]_{t_1}^{t_2} + \int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right) \delta q dt = 0 \quad (6)$$

and the bracketed expression is zero because the endpoints are held fixed. And then we can pull out the Euler-Lagrange equation:

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = 0. \quad (7)$$

Now it makes more sense to me. You start by treating the variables as independent but then remove the independence by imposing a condition *during* the derivation.