Prof. Kavita Dorai, Department of Physics IISER Mohali. kavita@iisermohali.ac.in

- 1. Suppose a delta-function bump is put in the center of the infinite square well: $H' = \alpha \delta(x a/2)$ where α is a constant. Find the first-order correction to the allowed energies.
- 2. Consider a harmonic oscillator with a slightly increased spring constant $k' = (1 + \epsilon)k$. Calculate the first order perturbation in the energy.
- 3. Consider a charged particle in a one-dimensional harmonic oscillator potential. Suppose a weak electric field is turned on so that the energy is shifted by an amount H' = -qEx. Show that there is no first order change in the energy levels and calculate the second order correction. The Schrodinger equation can be solved exactly for this case by a change of variables $x' = x (qE/m\omega^2)$. Find the exact energies and show that they are consistent with the perturbation theory expressions.
- 4. Consider an isotropic harmonic oscillator in two dimensions with the Hamiltonian

$$H_0 = \frac{p_x^2}{2m} + \frac{p_y^2}{2m} + \frac{m\omega^2}{2}(x^2 + y^2)$$

- (a) Now apply a perturbation $H_p = \delta m \omega^2 xy$, where δ is a dimensionless real number <<1. Find the energy upto first order for the ground state. (b) Solve the $H_0 + H_p$ problem exactly and compare with the perturbation results.
- 5. Calculate the energy of the ground state using first-order perturbation theory for a particle moving in a 1D box potential of length L with walls at x=0 and x=L, when a weak potential $H_p=\lambda x^2$ is added,where $\lambda << 1$.
- 6. Consider two identical spin-1/2 particles confined in an isotropic 3D harmonic oscillator potential of frequency ω . (a) Find the ground state energy and the corresponding wave function of this system when the two particles do not interact. (b) Consider now that there exists a weakly attractive spin-dependent potential between the two particles, $V(r_1, r_2) = -kr_1r_2 \lambda S_{1z}S_{2z}$, where k and λ are two small positive real numbers. Find the ground state to first-order time-independent perturbation theory.