

Assignment 10

PHY310: Mathematical Methods for Physicists I

Instructor: **Dr. Prasenjit Das**

No Need to Submit

- Find the real and imaginary parts $u(x, y)$ and $v(x, y)$ of the following functions:
(a) e^{iz} and (b) $\sqrt{1+z^2}$.
- Show whether or not the functions (a) $f(z) = x$ and (b) $f(z) = z^*$ is analytic.
- If $f(z) = u(x, y) + iv(x, y)$ is analytic, show that $u(x, y)$ and $v(x, y)$ cannot both have either a maximum or a minimum in the interior of any region in which $f(z)$ is analytic.
- Find the analytic function $f(z) = u(x, y) + iv(x, y)$ if (a) $u(x, y) = x^3 - 3xy^2$ and (b) $v(x, y) = e^{-y} \sin x$.
- If there is some common region in which $w_1 = u(x, y) + iv(x, y)$ and $w_2 = u_1^*$ are both analytic, prove that $u(x, y)$ and $v(x, y)$ are constants.
- The function $f(z) = u(x, y) + iv(x, y)$ is analytic. Show that $f^*(z^*)$ is also analytic.
- Using $f(z) = f(re^{i\theta}) = R(r, \theta)e^{i\phi(r, \theta)}$ in which $R(r, \theta)$ and $\phi(r, \theta)$ are differentiable real functions of r and θ , show that the *Cauchy-Riemann* conditions in polar coordinates becomes

$$\frac{\partial R}{\partial r} = \frac{R}{r} \frac{\partial \phi}{\partial \theta} \quad \text{and} \quad \frac{1}{r} \frac{\partial R}{\partial \theta} = -R \frac{\partial \phi}{\partial r}.$$

Applying these conditions, study the analyticity of $|z|$.

- The function $f(z)$ is analytic. Show that the derivative of $f(z)$ with respect to z^* does not exist unless $f(z)$ is a constant.
- For each of the following functions $f(z)$, find $f'(z)$ and identify the maximal region within which $f(z)$ is analytic (a) $f(z) = \sin z/z$, (b) $f(z) = z^2 - 3z + 2$, (c) $f(z) = 1/z(z+1)$, and (d) $f(z) = \tanh(z)$.
- The functions $f(z)$ have a derivative for what complex values (a) $f(z) = z^{3/2}$, (b) $f(z) = \tan^{-1} z$, (c) $f(z) = z^{-3/2}$, and (d) $f(z) = \tanh^{-1}(z)$.
- Show that the integral

$$\int_{3+i4}^{4-i3} (4z^2 - i3z) dz$$

has the same value on the two paths: (a) the straight line connecting the integration limits and (b) an arc on the circle $|z| = 5$.

- Verify that $\int_0^{1+i} z^* dz$ depends on the path by evaluating the integral for two paths (a) a straight line connecting (0,0) and (1,1) (b) two straight lines (0,0) to (1,0) and (1,0) to (1,1). Explain the outcome.

13. Show that

$$\oint \frac{dz}{z^2 + z} = 0$$

in which the contour C is a circle defined by $|z| = R > 1$.

14. For a square shaped contour with vertices located at $\pm 1 \pm i$, evaluate

$$\oint \frac{dz}{z}.$$

15. For a circular contour C defined by $|z - 1| = 1$, evaluate

$$\oint \frac{dz}{z^2 - 1}.$$

16. Show that the following integral is a representation of the Kronecker $\delta_{m,n}$:

$$\frac{1}{2\pi i} \oint z^{m-n-1} dz,$$

where m, n are integers.