

1. In an effort to clear impurities from a fabricated nano-wire, a laser beam is swept repeatedly along the wire, in the presence of a parallel electric field. After one sweep, an impurity initially at $x = 0$ has the following probability density of being found at a new position x

$$p(x) = \begin{cases} p(x) = \frac{1}{3}\delta(x) + \frac{2}{3a}e^{-x/a} & x \geq 0 \\ 0 & \text{otherwise,} \end{cases}$$

where a is some characteristic length. Calculate the probability density for the total distance d , the impurity has moved along the wire after 36 sweeps.

2. Consider one cubic centimeter of a dilute gas of atoms of mass M in thermal equilibrium at temperature $T = 0^\circ\text{C}$ and atmospheric pressure. The number of atoms (or molecules) in a cubic meter of an ideal gas at $T = 0^\circ\text{C}$ and atmospheric pressure has the value $2.69 \times 10^{25} \text{m}^{-3}$.

- a) For the kinetic energy of a single atom, find a numerical value for the ratio of standard deviation (the square root of the variance) to the mean.
- b) Find the same ratio for total energy of the gas, assumed to be all kinetic.

3. Consider a magnetic system which is at thermodynamic equilibrium at a temperature T . Let μ be the magnetic moment of each spin; and let M be the magnetization per spin, so that $-\mu < M < \mu$. The free energy per spin is given by $F(M)$.

- a) suppose $F(M) = \lambda \left[\left(\frac{M}{\mu} \right)^4 - \left(\frac{M}{\mu} \right)^2 \right]$. Is this a valid expression for the free energy?
- b) what would go wrong if this is indeed a valid free energy?


4. Consider the velocity of a gas in one dimension ($-\infty < v < \infty$)

- a) Find the **unbiased probability density** $p_1(v)$, subject only to the constraint that the average speed is c , that $\langle |v| \rangle = c$
- b) Now find the probability density $p_2(v)$, given only the constraint of average kinetic energy, $\langle mv^2/2 \rangle = mc^2/2$.
- c) Which of the above statements provide more information on the velocity? Quantify the difference in information in terms of $l_2 - l_1 = (\langle \ln p_2 \rangle - \langle \ln p_1 \rangle)/2$

5. You are given several unusual three-sided dice which, when rolled, show either one, two, or three spots. There are three games played with these dice: *Distinguishable*, *Bosons*, and *Fermions*. In each turn in these games, the player rolls one die at a time, starting over if required by the rules, until a legal combination occurs. In *Distinguishable*, all rolls are legal. In *Bosons*, a roll is legal only if the new number is larger or equal to the preceding number. In *Fermions*, a roll is legal only if the new number is strictly larger than the preceding number. Let the sum of the spots as result of throwing n dices be S_n .

- a) construct a table showing the different values for S_2 . Which of these correspond to *Fermions* and which of these are *Bosons*?
- b) Presume the dice are fair: each of the three numbers of dots shows up $1/3$ of the time. For a legal turn rolling a die twice in *Bosons*, what is the probability of $S_2 = 4$ of rolling a 4? Similarly, among the legal *Fermion* turns rolling two dice, what is the probability of $S_2 = 4$?

- c) For a legal turn rolling three three-sided dice in Fermions, what is the probability of $S_3 = 6$ of rolling a 6?
- d) Consider rolling M dice with N sides each. How many legal turns for Bosons are there?
- e) In a turn of three rolls, what is the factor by which the probability of getting triples in Bosons is enhanced over that in Distinguishable? In a turn of M rolls, what is the enhancement factor for generating an M -tuple (all rolls having the same number of dots showing) for a three sided dice? What happens to the same expression in the general case when you have a N faced dice?
6. The Carnot process is an idealized, quasistatic process with maximum efficiency. Taking a finite heat conduction between the working substance and the reservoirs into account, the duration of a cycle, however, is infinitely long and the power output is zero. To speed up the heat transfer, temperature differences between the working substance and the reservoirs are required. In the Curzon-Ahlborn heat engine, the working substance performs a Carnot cycle as depicted to the right. It alternates between temperatures $T_{1w} < T_{2w}$, where it is in contact with either of two reservoirs at T_1 and T_2 . Assume that the heat flux of the isothermal stages lasts for periods $\tau_i (i = 1, 2)$ and is described by $\dot{Q}_i = K\Delta T_i$ with heat conductivity K . Neglect the duration of the fast adiabatic stages and possible associated heat fluxes.



- a) Determine the entropy changes of the working substance, S_{AB} and S_{CD} , and of the reservoirs, \tilde{S}_{AB} and \tilde{S}_{CD} , during the isothermal stages. Calculate the rate of entropy production due to heat conduction.
- b) Find the optimal temperatures T_{1w}, T_{2w} for given T_1, T_2 by maximizing the power output, $P = (Q_2 Q_1) / (\tau_1 + \tau_2)$, with respect to the variables $\tau_1, \tau_2, T_{1w}, T_{2w}$ under the constraint $S_{AB} = S_{CD}$.
- c) Calculate the efficiency of the optimized engine in terms of T_1 and T_2 . Compare with the efficiency of the Carnot process and with existing engines, e.g., a Diesel engine or a coal-fired power plant.