

1. Calculate the number of accessible microstates of a system of two localized independent quantum oscillators with fundamental frequencies  $\omega_0$  and  $3\omega_0$ , and total energy of  $10\hbar\omega_0$ .
2. The spin Hamiltonian of a system of  $N$  localized magnetic ions is given by

$$\mathcal{H} = D \sum_{j=1}^N S_j^2,$$

where  $D > 0$  and spin variables  $S_j$  may assume values  $\pm 1$  or  $0$ , for  $j = 1, 2, 3, \dots$ . This spin Hamiltonian describes the effects of the electrostatic environment on spin-1 ions. An ion in state  $\pm 1$  has energy  $D > 0$  and in state  $0$  has zero energy.

- a) Show that the number of accessible microstates of the system is given by

$$\Omega(U, N) = \frac{N!}{(N - U/D)!} \sum_{N_-} \frac{1}{(U/D - N_-)! N_-!}$$

- b) Calculate the binomial sum and obtain an exact result for  $\Omega(U, N)$
  - c) Now use Sterling's approximation and obtain the entropy of the system. Is it extensive?
  - d) Calculate the specific heat of the system as a function of temperature.
3. Consider  $N$  particles distributed in a volume  $V$ . Now divide the volume into cell of size  $b$ , with  $N \leq V/b$ . Suppose that each cell may be either empty or can be occupied by a single particle.
    - a) calculate the number of microstates accessible to the system.
    - b) from the above result, calculate the entropy of the system and hence the quantity  $P/T$ , where the symbols have their usual meaning.
    - c) do you see any difference with an ideal gas? If yes what do think is the reason behind this difference?
  4. In the class we have worked out the problem of  $N$  quantum harmonic oscillators in the microcanonical ensemble. Now assume that the fundamental frequency has volume dependence given by:

$$\omega = \omega(v) = \omega_0 - A \ln \left( \frac{v}{v_0} \right),$$

where  $v = V/N$ , and  $\omega_0$ ,  $A$ , and  $v_0$  are positive constants. Calculate the expansion coefficient and the compressibility of the system.

5. Consider system of ideal gas of  $N$  particles which can be in discrete energy states  $\epsilon_j$ ;  $j = 0, 1, 2, 3, \dots$ 
  - a) how would you specify the microscopic state of the system?
  - b) calculate the number of accessible microstates for the system.
  - c) assuming that the total particle number and total energy of the system is fixed, write down the constraint equations.
  - d) can you now extremize the system and find out probability of finding  $N_i$  particle in an energy state  $\epsilon_i$ ? Do you recognize the distribution.

6. Consider a magnetic system with a total energy  $E$  and having  $N$  spins. The Hamiltonian for the system is  $\mathcal{H} = -\mu H \sum_i \sigma_i$ , with  $\sigma_i = \pm 1$ . In a microcanonical ensemble, we want to calculate the total number of accessible microstates.
- a) Assume that there are  $N_+$  up spins and  $N_-$  down spins. Express the  $N_+$  and  $N_-$  in terms of  $E$  and  $N$ .
  - b) From this calculate the total number of microstates accessible to the system for  $E, N \rightarrow \infty$  and  $E/N = u$  fixed.
  - c) Hence calculate the entropy per spin and derive an expression for the energy per spin of the system.
  - d) Using all the above informations (not all are required though) derive an expression for the magnetization of the system.
7. We want to look at the above problem using a different approach.
- a) The total number of accessible microstates is given by  $\Omega = \sum_{\{\sigma_i = \pm 1\}} \delta(E + h \sum_i \sigma_i)$ . Using the integral representation of the delta function  $\delta(x - a) = \int_{-\infty}^{\infty} e^{ik(x-a)} \frac{dk}{2\pi}$ , rewrite the expression for  $\Omega$ .
  - b) Now do the sum over the states of the individual spins  $\sigma_i$  to express  $\Omega$  as  $\Omega = 2^N \int_{-\infty}^{\infty} e^{f(k)} \frac{dk}{2\pi}$ . Determine  $f(k)$ .
  - c) Verify that the argument of the exponential goes as  $\mathcal{O}(N)$ . We can therefore use a saddle point approximation. To do this find the minimum  $k_0$  of the function  $f(k)$  and verify that it is indeed a minimum (i.e.  $f''(k_0) < 0$ ). You need the following identity :  $\arctan(ix) = \frac{1}{2} \log \left( \frac{1+x}{1-x} \right)$
  - d) Once you have the value of  $k_0$ , you want to re-express the integral in  $\Omega$ . Recall how we did it while establishing the connection between the canonical partition function and the free energy  $F$ . You have to approximate the function  $f(k)$  by a Taylor series around  $k_0$  and retain the second order term. Once you have done that carry out the integral over and determine  $\Omega$  as function of  $e = E/Nh$ .