

## PHY 635: Gravitation & Cosmology : Problem Sheet 3

Submit before 05th March 2025

1. In the case if the spacetime has torsion, i.e.  $T_{\mu\nu}^{\lambda} \equiv \Gamma_{\mu\nu}^{\lambda} - \Gamma_{\nu\mu}^{\lambda} \neq 0$ , a vector (or a co-vector) fails to comeback to itself when parallely translated along a closed loop, even in the globally flat spacetime (no curvature).
2. Using the relation  $\log \text{Det} M = \text{Tr}(\log M)$  for a matrix  $M$ , find out the expression for  $\delta(-g)^{-n/2} / \delta g^{\mu\nu}$ , in terms of  $\delta g_{\mu\nu}$  and  $\delta g^{\mu\nu}$  with  $g$  being the determinant of matrix form of  $g_{\mu\nu}$ .
3. Show that the Einstein's field equation  $G_{\mu\nu} = 8\pi G T_{\mu\nu}$  suggests that all vacuum spacetimes are *Ricci flat*, i.e.  $R_{\mu\nu} = 0$ . What is the condition if the Einstein's equation gets corrected by a presence of cosmological constant to  $G_{\mu\nu} = 8\pi G T_{\mu\nu} + \Lambda g_{\mu\nu}$  ?
4. A scalar field has the Lagrangian density as

$$\mathcal{L} = \frac{1}{2\alpha} \sin [\alpha(\partial_{\mu}\phi)(\partial^{\mu}\phi)] - V(\phi).$$

Obtain the expression of the stress energy tensor prescribed as

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S[\phi]}{\delta g^{\mu\nu}},$$

where  $S[\phi]$  is the action for the scalar field.

5. A simple spacetime has a certain quantity called geodesic deviation expansion parameter  $\theta$  defined as

$$\frac{d\theta}{d\lambda} = -\alpha\theta^2 - R_{\mu\nu}u^{\mu}u^{\nu},$$

where  $R_{\mu\nu}$  is the Ricci tensor,  $u^{\mu}$  is tangent along the geodesic which is parameterized by  $\lambda$  and  $\alpha$  is a constant. Find out for the geodesic expansion rate to become positive (i.e. for gravity to become repulsive), what is the necessary condition for the matter stress energy tensor  $T_{\mu\nu}$ , in Einstein's theory of relativity ?