

# COVID-19 Medical Inventory Prediction

Deepak Padmanabhan\*

P Sunthar<sup>†</sup>

April 3, 2020

## Abstract

This document describes the assumptions behind the estimated medical inventory requirement for attending to COVID-19 patients in hospitals, as projected in <https://covid19medinventory.in>. The estimated patient counts have been obtained from <https://mesoscalelab.github.io/covid19/>. The projections are themselves dynamic, and can change every week depending on newly available evidence. The medical inventory list and the formulae for its dependence on the type of patients is also dynamic, and will change as we find better relationships with actual field data.

## 1 Basic Assumptions

The number of COVID-19 positive patients obtained from the district-wise projection<sup>1</sup> is denoted as  $N$ . This could be the current number or the projected number in future weeks. According to WHO statistics<sup>2</sup>, the number of patients who actually need various levels of care<sup>3</sup> is given in the table below:

Type of patient	Type of care	approx %	Symbol used here
Total positive	–	100%	$N$
Mild	Symptomatic, Home Quarantine/isolation (out-patients)	40%	$N_q$
Moderate	In-patient ward	40%	$N_i$
Severe	Supportive care, oxygen therapy	15%	$N_s$
Critical	ICU, mechanical ventilation	5%	$N_c$
Deaths	–	2.5%	$N_d$

## 2 Inventory Estimates

The tables below give the inventory list, the estimation logic, and formula, and an example calculation for  $N = 100$  positive cases.

In the example case the patient estimates are given below. In the website, some of the numbers are rounded higher to the nearest 10 or 50, depending on their magnitude.

Category	Symbol	Estimate
Total positives	$N$	100
Mild	$N_q$	40
Moderate	$N_i$	40
Severe	$N_s$	15
Critical	$N_c$	5
Deaths	$N_d$	2.5

\*MBBS, MD DM, Cardiologist, Sri Jayadeva Institute of Cardiology, Bengaluru

<sup>†</sup>Professor, Department of Chemical Engineering, IIT Bombay, Mumbai [p.sunthar@iitb.ac.in](mailto:p.sunthar@iitb.ac.in)

Table 1: Persons and Protective Equipment. The last column is rounded number assuming a total positive cases of  $N = 100$ .

Item [Unit]	Reason	Formula	$N = 100$
Doctors [per day]	2 per 12-hr shift for 10 critical and severe patients and 2 per 12-hr shift for 15 in-patients	$\frac{2}{5} \left( N_c + N_s + \frac{N_i}{1.5} \right)$	19
Nurses [per day]	1 per 8-hr shift for 3 critical and severe patients and 1 per 8-hr shift for 5 in-patients	$\left( N_c + N_s + \frac{N_i}{1.5} \right)$	47
Total staff [per day]	Doctors, Nurses	$S \equiv \frac{7}{5} \left( N_c + N_s + \frac{N_i}{1.5} \right)$	66
Staff PPE (Gowns, Masks, Goggles etc.) [per day]	Two per staff	$2 S$	132
Patient PPE: Masks [per day]	4 per severe and moderate patients	$4(N_s + N_i)$	220
Sterile gloves [per day]	3 per patient per 6-hr shift for critical and half that for severe and moderate	$12N_c + 6(N_s + N_i)$	390
Non-sterile gloves	6 per patient per 6-hr shift for critical and half that for severe and moderate	$24N_c + 12(N_s + N_i)$	780
Dead body bags	1 per death	$N_d$	3

Table 2: Medical Equipment. The last column is rounded number assuming a total positive cases of  $N = 100$ .

Item [Unit]	Reason	Formula	$N = 100$
Ventilators, Ambu bags, Glass case	1 per critical patient	$N_c$	5
Laryngoscopes, Defibrillator	3 per 10 critical patients	$3N_c/10$	2
ECG	1 per 20 critical patients	$N_c/20$	1
Arterial BP monitors	1 per critical patient	$N_c$	5
Arterial blood gas machine	1 per 30 critical patients	$N_c/30$	1
Bedside X-ray	1 per 20 critical patients	$N_c/20$	1
Infusion pumps	3 per critical patient	$3N_c$	15
Oxymeter	1 per 20 severe patients	$N_s/20$	1
High flow nasal canula	1 per severe patient	$N_s$	15
Nebuliser	1 per severe patient	$N_s$	15
Non-contact Thermometer	1 for 20 out-patients	$N_q$	2
Patient cot	1 per severe and moderate	$N_s + N_i$	55
Wheel chair	1 per 20 severe patients	$N_s/20$	1
Stretcher	1 per critical and 3 per 20 severe patients	$N_c + 3 * N_s/20$	7
Ambulance	3 times total positive cases, each making 20 trips per day	$3N/20$	15

Table 3: Medical Consumables. The last column is rounded number assuming a total positive cases of  $N = 100$ .

Item [Unit]	Reason	Formula	$N = 100$
Needles [per day]	10 per critical, 5 per severe, 2 per moderate	$10N_c + 5N_s + 2N_i$	205
Disposable Bags [per day]	3 per critical 2 per severe and 1 per 4 moderate	$3N_c + 2N_s + 0.25N_i$	55
Sanitizer [lt/day]	250 ml per all in-patients	$0.25(N_c + N_s + N_i)$	15
Endotracheal Tube [per day]	1 per critical patients for 3 days	$N_c/3$	2
Oxygen (medium) [cylinders per day]	4 cylinders per critical patient	$4N_c$	20
Central and Peripheral lines [per day]	1 per critical patient for 3 days	$N_c/3$	2
IV Fluids [lt/day]	5 bottles of 500 ml per critical patient	$2.5N_c$	13
Suction catheter [per day]	1 per critical patient	$N_c$	5
Test kits [per day]	3 times the number of positive cases	$3N$	300

### 3 Acknowledgements

The authors thank Dr Santosh Ansumali (JNCASR) and Dr. Alope Kumar (IISc) for providing the projection maps, and Dr. Soumyadeep Bhattacharjee (Sankhyasutra Labs Pvt. Ltd) for data API. We also thank the office of the Principal Scientific Advisor of Govt. of India for providing an elaborate list of inventory from which some critical ones have been used here.

### 4 Errors and Omissions

Though the authors have taken sufficient care in estimating the quantity of the inventory, the proportions used here may change depending on the actual field data in India for the patients and the inventory usage pattern. Please email your comments and suggestions with justifications to [p.sunthar@iitb.ac.in](mailto:p.sunthar@iitb.ac.in) for inclusion in the future updates of the software.

### References

- [1] Santosh Ansumali and Alope Kumar. A district-wise projection map for covid-19 in india. Technical report, JNCASR and IISc, 2020.
- [2] *Operational considerations for case management of COVID-19 in health facility and community*. World Health Organisation. 19 March 2020.
- [3] *COVID-19 Preparedness document*. AIIMS, New Delhi. 27 March 2020.