

## 《线性代数》 期末试题答案 (A/B)

### 1. Fill in the blank (10\*3=30 Pts)

(1)  $x = \begin{bmatrix} 11 \\ 2 \end{bmatrix}$

(2)  $\det A = 6$

(3)  $a = 8, b = -7$

(4)  $\hat{y} = \frac{2}{3} \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}$

(5)  $A = \begin{bmatrix} -2 & 1 \\ 4 & -1 \end{bmatrix}$

(6) A basis for Row  $A$  is  $\{ (2, -3, 6, 2, 5), (0, 0, 3, -1, 1), (0, 0, 0, 1, 3) \}$ ,

$\dim \text{Nul } A$  is 2,  $\text{rank } A^T$  is 3

(7)  $v_1 = \begin{bmatrix} 1+3i \\ 2 \end{bmatrix}$

### 2. Mark each statement True or False, and describe your reasons. (5\*4=20 Pts)

(1) T

(2) T

(3) F

(4) T

(5) F

### 3. Calculation issues (5\*6=30 Pts)

(1)

**Solution**

**a. Determine if the equation  $c_1v_1+c_2v_2+c_3v_3=x$  is consistent.**

$$\begin{bmatrix} 5 & 3 & 1 & -2 \\ 0 & 2 & -1 & 3 \\ 0 & -2 & 1 & 0 \\ 0 & -2 & 1 & -3 \end{bmatrix} \sim \begin{bmatrix} 5 & 3 & 1 & -2 \\ 0 & 2 & -1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}. \quad \text{Thus } x \text{ isn't in } W.$$

**b. Since  $y \cdot v_1 = 0, y \cdot v_2 = 0, y \cdot v_3 = 0$ , so  $y$  is in  $W^\perp$ .**

(2)

**Solution**

$$AB + I = A^2 + B \Rightarrow AB - B = A^2 - I \Rightarrow (A - I) B = A^2 - I \Rightarrow B = (A - I)^{-1}(A^2 - I)$$

$$\text{thus } B = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ -1 & 0 & 2 \end{bmatrix}$$

(3)

**Solution**

**a. Computer and find**  $Av_1 = -5v_1, Av_2 = 4v_2, Av_3 = 4v_3$ , and  $\{v_2, v_3\}$  must be replace by an orthogonal basis.

$$\hat{v}_3 = \left( \frac{v_3 \cdot v_2}{v_2 \cdot v_2} \right) v_2 = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad v_3 - \hat{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/2 \\ -1/2 \\ 2 \end{bmatrix},$$

**And normalize the eigenvectors.**

$$\text{Place eigenvectors into } P = \begin{bmatrix} -2/3 & 1/\sqrt{2} & 1/\sqrt{18} \\ 2/3 & 1/\sqrt{2} & -1/\sqrt{18} \\ 1/3 & 0 & 4/\sqrt{18} \end{bmatrix}, \text{ and set } D = \begin{bmatrix} -5 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}_1$$

$$\text{Equation } A = PDP^{-1}$$

**b.**  $3x_3^2 + 8x_1x_2 + 4x_1x_3 - 4x_2x_3$ . **The quadratic form is indefinite.**

(4)

**Solution**

$$A^T A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 5 & 4 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 5 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 14 & 21 \\ 21 & 42 \end{bmatrix}, \quad A^T b = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 5 & 4 \end{bmatrix} \begin{bmatrix} -1 \\ 10 \\ 3 \end{bmatrix} = \begin{bmatrix} 28 \\ 63 \end{bmatrix}.$$

$$\text{Solving } A^T Ax = A^T b, \quad \begin{bmatrix} 14 & 21 & 28 \\ 21 & 42 & 63 \end{bmatrix} \sim \dots \sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}, \text{ so the solution is } \hat{x} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

$$\text{Or } \hat{x} = (A^T A)^{-1} A^T b = \frac{1}{147} \begin{bmatrix} 42 & -21 \\ -21 & 14 \end{bmatrix} \begin{bmatrix} 28 \\ 63 \end{bmatrix} = \frac{1}{147} \begin{bmatrix} -147 \\ 294 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$\text{And } b - A\hat{x} = \begin{bmatrix} -1 \\ 10 \\ 3 \end{bmatrix} - \begin{bmatrix} 1 & -1 \\ 2 & 5 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ -2 \end{bmatrix}, \text{ thus } \|b - A\hat{x}\| = \sqrt{2^2 + 2^2 + (-2)^2} = \sqrt{12}$$

(5)

**Solution**

A typical element of  $H$  can be written as

$$\begin{bmatrix} 3a+7b-c \\ -5b+8c-2d \\ 3d-4e \\ 5b-8c-d+4e \end{bmatrix} = a \begin{bmatrix} 3 \\ 0 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} 7 \\ -5 \\ 0 \\ 5 \end{bmatrix} + c \begin{bmatrix} -1 \\ 8 \\ 0 \\ -8 \end{bmatrix} + d \begin{bmatrix} 0 \\ -2 \\ 3 \\ -1 \end{bmatrix} + e \begin{bmatrix} 0 \\ 0 \\ -4 \\ 4 \end{bmatrix}$$

$\mathbf{u}_1 \quad \mathbf{u}_2 \quad \mathbf{u}_3 \quad \mathbf{u}_4 \quad \mathbf{u}_5$

a.  $H$  is a vector space because it is the set of all linear combinations of a set of vectors.

b. Row reduce:

$$\begin{bmatrix} 3 & 7 & -1 & 0 & 0 \\ 0 & -5 & 8 & -2 & 0 \\ 0 & 0 & 0 & 3 & -4 \\ 0 & 5 & -8 & -1 & 4 \end{bmatrix} \sim \begin{bmatrix} 3 & 7 & -1 & 0 & 0 \\ 0 & -5 & 8 & -2 & 0 \\ 0 & 0 & 0 & 3 & -4 \\ 0 & 0 & 0 & -3 & 4 \end{bmatrix} \sim \begin{bmatrix} 3 & 7 & -1 & 0 & 0 \\ 0 & -5 & 8 & -2 & 0 \\ 0 & 0 & 0 & 3 & -4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The pivot columns are 1,2, and 4. So  $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_4\}$  is a basis for  $H$ .

#### 4. Prove issues (2\*10=20 Pts)

(1)

**Proof** Since  $A^T = (I_n - \frac{2}{\|x\|^2} xx^T)^T = I_n - \frac{2}{\|x\|^2} xx^T = A$ ,  $A$  is symmetric.

$$\text{And because } A^T A = A^2 = (I_n - \frac{2}{\|x\|^2} xx^T)(I_n - \frac{2}{\|x\|^2} xx^T)$$

$$= I_n - \frac{4}{\|x\|^2} xx^T + \frac{4}{\|x\|^4} x(x^T x)x^T$$

$$= I_n - \frac{4}{\|x\|^2} xx^T + \frac{4}{\|x\|^4} x(\|x\|^2)x^T$$

$$= I_n - \frac{4}{\|x\|^2} xx^T + \frac{4}{\|x\|^2} xx^T$$

$$= I_n,$$

thus  $A$  is a orthogonal matrix.

Hence  $A$  is both orthogonal and symmetric.

(2)

**Proof** If there is an orthogonal matrix  $T$  such that  $T^{-1}AT = B$ , then  $A$  is similar to  $B$ , thus they have the same characteristic polynomial.

Conversely, if they have the same characteristic polynomial, then they have the same eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ , thus there is orthogonal matrix  $T_1$  and  $T_2$  such that

$$T_1^{-1}AT_1=[\lambda_1,\lambda_2,...,\lambda_n]$$

$$T_2^{-1}BT_2=[\lambda_1,\lambda_2,...,\lambda_n]$$

**Then**

$$T_1^{-1}AT_1=T_2^{-1}BT_2$$

$$(T_1T_2^{-1})^{-1}AT_1T_2^{-1}=B$$

**Let  $T=T_1T_2^{-1}$  , then  $T$  is an orthogonal matrix such that  $T^{-1}AT=B$**