

《线性代数》 期末试题 (A/B 卷)

(考试形式：闭 卷 考试时间：2 小时)



《中山大学授予学士学位工作细则》第六条

考试作弊不授予学士学位

方向：_____ 姓名：_____ 学号：_____

出卷：_____ 复核：_____

1. Fill in the blank (10*3=30 Pts)

(1) If the vector x determined by the coordinate vector $[x]_B = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$ and the basis

$$B = \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}, \text{ then } x = \underline{\hspace{2cm}}.$$

(2) Let A be a 4×4 matrix, and suppose the eigenvalues of A are 3, 1, 1, and 2, then $\det A = \underline{\hspace{2cm}}$

(3) The matrix $A = \frac{1}{9} \begin{bmatrix} -1 & 4 & a \\ a & 4 & -1 \\ 4 & b & 4 \end{bmatrix}$ is an orthogonal matrix, then $a = \underline{\hspace{2cm}}$,

$b = \underline{\hspace{2cm}}.$

(4) If $v_1 = \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}, v_2 = \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}$, and $W = \text{Span}\{v_1, v_2\}$, then the closest point in

W to y is $\underline{\hspace{2cm}}.$

(5) If the transformation $x \mapsto Ax$ maps $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 7 \end{bmatrix}$ into $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$, respectively, then the matrix $A = \underline{\hspace{2cm}}.$

(6) Assume the matrix A is row equivalent to B .

$$A = \begin{bmatrix} 2 & -3 & 6 & 2 & 5 \\ -2 & 3 & -3 & -3 & -4 \\ 4 & -6 & 9 & 5 & 9 \\ -2 & 3 & 3 & -4 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & -3 & 6 & 2 & 5 \\ 0 & 0 & 3 & -1 & 1 \\ 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

A basis for Row A is _____, and $\dim \text{Nul } A$ is _____, and $\text{rank } A^T$ is _____.

(7) $A = \begin{bmatrix} 1 & 5 \\ -2 & 3 \end{bmatrix}$ act on C^n , the eigenvalues of A is $\lambda = 2 \pm 3i$, and a basis for the eigenspace corresponding to $\lambda_1 = 2 - 3i$ is _____.

2. Mark each statement True or False, and describe your reasons. (5*4=20 Pts)

(1) Let $T: R^n \mapsto R^n$ be a linear transformation defined by $T(x) = Ax$. Then T is onto if and only if $\det(A) \neq 0$.

(2) If A is an $n \times n$ matrix, then $\text{rank } A < n$ if and only if some eigenvalue of A is zero.

(3) The set of all solution to the linear system $Ax = b$, where A is $m \times n$ and $b \neq 0$, is a subspace of R^n .

(4) Every set of five orthonormal vectors is a basis for R^5 .

(5) The linear transformation $T: P_2 \mapsto P_2$ defined by $T(at^2 + bt + c) = 2at + b$ is one-to-one.

3. Calculation issues (5*6=30 Pts)

(1) Let $v_1 = \begin{bmatrix} 5 \\ 0 \\ 0 \\ 0 \end{bmatrix}$, $v_2 = \begin{bmatrix} 3 \\ 2 \\ -2 \\ -2 \end{bmatrix}$, $v_3 = \begin{bmatrix} 2 \\ -1 \\ 1 \\ 1 \end{bmatrix}$, $x = \begin{bmatrix} -2 \\ 3 \\ 0 \\ -3 \end{bmatrix}$, $y = \begin{bmatrix} 0 \\ 3 \\ 6 \\ -3 \end{bmatrix}$, and $W = \text{Span}\{v_1, v_2, v_3\}$

a. Determine whether x is in W

b. Determine whether y is in the orthogonal of W .

(2) If A and B are 3×3 matrices, I is the identity matrix, and $AB + I = A^2 + B$, where

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}. \quad \text{Find } B.$$

(3) Let $A = \begin{bmatrix} 0 & 4 & 2 \\ 4 & 0 & -2 \\ 2 & -2 & 3 \end{bmatrix}$, $v_1 = \begin{bmatrix} -2 \\ 2 \\ 1 \end{bmatrix}$, $v_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $v_3 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$. The v_1, v_2 , and v_3 are

eigenvectors for A .

- Find matrices P and D that orthogonally diagonalize A , and write the equation that relates A to P and D .
- Write the quadratic form associated with A using variables x_1, x_2 , and x_3 . And classify the quadratic form as positive (or negative) definite, or indefinite.

(4) Let $A = \begin{bmatrix} 1 & -1 \\ 2 & 5 \\ 3 & 4 \end{bmatrix}$, $b = \begin{bmatrix} -1 \\ 10 \\ 3 \end{bmatrix}$. Find the least-squares solution of $Ax = b$, and

determine the least-squares error in the least-squares solution of $Ax = b$.

(5) Let $H = \left\{ \begin{bmatrix} 3a+7b-c \\ -5b+8c-2d \\ 3d-4e \\ 5b-8c-d+4e \end{bmatrix} : a, b, c, d, e \text{ any real numbers} \right\},$

- Show H is a subspace of R^4 .
- Find a basis for H .

4. Prove issues (2*10=20 Pts)

- (1) Show that if x is a nonzero vector in R^n , then the $n \times n$ matrix

$$A = I_n - \frac{2}{\|x\|^2} xx^T$$

is both orthogonal and symmetric.

- (2) Suppose A and B are both real symmetric matrix, then they have the same characteristic polynomial if and only if there is an orthogonal matrix T such that $T^{-1}AT = B$.