

《线性代数》期中试题答案

1. Fill in the blank (5×4=20 Pts)

(1) $\det(2A) = 320$

(2) a, b

(3) $X = B^{-1}, Y = -B^{-1}AC$

(4) $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

(5) $3, \left\{ \begin{bmatrix} 2 \\ -1 \\ 3 \\ 5 \end{bmatrix}, \begin{bmatrix} -1 \\ -4 \\ 6 \\ 5 \end{bmatrix}, \begin{bmatrix} 2 \\ -7 \\ 13 \\ 10 \end{bmatrix} \right\}$

2. Mark each statement True or False, and descript your reasons. (6×4=24 Pts)

(1) F (2) F (3) T (4) F (5) T (6) F

3. Calculation issues (5×8=40 Pts)

(1) Solution

$$\begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 2 & 4 & 1 & 1 & 5 \\ -1 & -2 & -2 & 1 & -4 \end{bmatrix} \sim \dots \sim \begin{bmatrix} 1 & 2 & 0 & 1 & 2 \\ 0 & 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -1 \\ 0 \\ 1 \\ 1 \end{bmatrix} \quad (\text{with } x_2, x_4 \text{ free})$$

(2) Solution

$$AB + I = A^2 + B \Rightarrow AB - B = A^2 - I \Rightarrow (A - I)B = A^2 - I \Rightarrow B = (A - I)^{-1}(A^2 - I)$$

$$\text{thus } B = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ -1 & 0 & 2 \end{bmatrix}$$

(3) Solution

$$A_1 = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}, A_2 = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 2 & 4 \\ 0 & 0 & 3 \end{bmatrix},$$

$$A = \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} A_1^{-1} & 0 \\ 0 & A_2^{-1} \end{bmatrix} = \begin{bmatrix} 3 & -5 & 0 & 0 & 0 \\ -1 & 2 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 4/3 \\ 0 & 0 & 0 & 1/2 & -2/3 \\ 0 & 0 & 0 & 0 & 1/3 \end{bmatrix}$$

(4) Solution

Take $U = \begin{bmatrix} 2 & -6 & 14 & 4 & 18 \\ 0 & 3 & -12 & 6 & 3 \\ 0 & 0 & 0 & 3 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$, further row reduction of U is not needed. For L , copy the

pivot columns that were found during the row reduction to U : $\begin{bmatrix} 2 \\ -1 \\ -2 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -5 \end{bmatrix}$. Divide each

column by the pivot at the top and place into the lower triangular part of L , then fill in the remaining column of L with the last column of the 4×4 identity matrix:

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1/2 & 1 & 0 & 0 \\ -1 & 1/3 & 1 & 0 \\ 3/2 & 1/3 & -5/3 & 1 \end{bmatrix}$$

(5) Solution

a. Standard matrix of T is $A = [T(e_1) \quad T(e_2) \quad T(e_3)] = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & -7 \\ 4 & 5 & 5 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & -7 \\ 0 & 0 & -2 \end{bmatrix}$. Since

A has three pivot positions, the IMT implies that A is invertible and the mapping is one-to-one.

b. $\begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}$

4. Prove issues (2×8=16 Pts)

(1) Proof

Since $\{v_1, v_2, v_3\}$ is a linearly independent set, $v_1 \neq 0$. Also, Theorem 7 shows that v_2 cannot be a multiple of v_1 , and v_3 cannot be a linear combination of v_1 and v_2 . By hypothesis, v_4 is not a linear combination of v_1, v_2 and v_3 . Thus, by Theorem 7, $\{v_1, v_2, v_3, v_4\}$ cannot be a linearly dependent set and so must be linearly independent.

(2) Proof

Let A be a 5×3 matrix and B a 3×5 matrix. Since B has more columns than rows, its five columns are linearly dependent and there is a nonzero x such that $Bx = 0$. Thus $ABx = A0 = 0$. This shows that the matrix AB is not invertible, by the IMT.