

Communication

The Asymmetric Piers Hydrosilylation

Lars Süsse, Julia Hermeke, and Martin Oestreich

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.6b03443 • Publication Date (Web): 23 May 2016

Downloaded from <http://pubs.acs.org> on May 24, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

The Asymmetric Piers Hydrosilylation

Lars Süsse, Julia Hermeke, and Martin Oestreich*

Institut für Chemie, Technische Universität Berlin, Strasse des 17. Juni 115, 10623 Berlin, Germany

Supporting Information Placeholder

ABSTRACT: An axially chiral, cyclic borane decorated with just one C_6F_5 group at the boron atom promotes the highly enantioselective hydrosilylation of acetophenone derivatives without assistance of an additional Lewis base (up to 99% ee). The reaction is an unprecedented asymmetric variant of Piers' $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed carbonyl hydrosilylation. The steric congestion imparted by the 3,3'-disubstituted binaphthyl backbone of the borane catalyst as well as the use of reactive trihydrosilanes as reducing agents are key to success.

Piers' discovery that $\text{B}(\text{C}_6\text{F}_5)_3$ catalyzes carbonyl hydrosilylation¹ opened a new chapter in reduction methodology that still continues to grow.² Part of the fascination with this reaction came from its at the time peculiar mechanism. Early insight had already suggested that it proceeds through activation of the hydrosilane reagent by $\text{B}(\text{C}_6\text{F}_5)_3$ rather than conventional Lewis pair formation with the carbonyl substrate.³ Over recent years, the full mechanistic picture evolved,⁴ and η^1 coordination of the Si-H bond to $\text{B}(\text{C}_6\text{F}_5)_3$ followed by $\text{S}_{\text{N}}2$ -Si displacement of hydride at the silicon atom with the carbonyl group as the nucleophile is now well accepted. The borohydride emerging from that step is the actual reducing agent.

An asymmetric variant of the Piers hydrosilylation would therefore require a chiral $\text{B}(\text{C}_6\text{F}_5)_3$ congener that is sufficiently electron-deficient to promote the Si-H bond activation and meets the challenge of inducing enantioselectivity as its borohydride. For this, we introduced axially chiral (S)-**1**-THF with one C_6F_5 group at the boron atom a few years ago (Figure 1, left)⁵ but enantioinduction was low ($\leq 15\%$ ee).⁶ Substantially better levels of enantioselection were obtained with (S)-**1**-THF in the related catalytic hydrosilylation of imines⁷ ($\leq 62\%$ ee).⁸ Even higher enantiomeric excesses ($\leq 87\%$ ee) were achieved by borrowing from the concept of frustrated Lewis pairs (FLPs).^{9,10} Klankermayer and co-workers employed an FLP- H_2 adduct composed of a terpene-derived borane and a bulky phosphine in the imine hydrosilylation; the hydrosilylation of acetophenone was again inferior (37% ee).¹¹ The FLP strategy was also successful in the hydrosilylation of α -keto carbonyl and carboxyl compounds ($>99\%$ ee), using one of Du's in-situ-generated catalysts (S)-**2** (Figure 1, right).^{12,13} The same catalytic setup afforded 42% ee in the reduction of acetophenone. To date, the asymmetric Piers hydrosilylation catalyzed by an electron-deficient borane alone¹⁴ is elusive, and we disclose here a solution to this long-standing problem.





Figure 1. Axially chiral congeners of $\text{B}(\text{C}_6\text{F}_5)_3$ for Si-H bond activation (LB = Lewis base).

To refine catalyst (S)-**1**-THF, we created steric congestion in the proximity of the boron atom by installation of phenyl groups in the 3 and 3' positions. These substituents had a dramatic effect on both catalyst preparation and purification. Introduction of the $\text{B}(\text{C}_6\text{F}_5)_3$ unit by conventional tin-boron exchange failed,⁵ and we had to develop a dummy-ligand strategy to overcome chemoselectivity issues [(S)-**4** \rightarrow (S)-**3**, Scheme 1].¹⁵ Moreover, Lewis-pair formation between (S)-**3** and various Lewis bases to precipitate or crystallize adducts of type (S)-**3**-LB was hampered by the steric situation around the boron atom in (S)-**3**. We eventually succeeded using 3,3-dimethyloxetane (3,3-DMO) and were able to crystallographically characterize (S)-**3**-3,3-DMO.¹⁵ However, considerable experimentation was required to reliably remove stoichiometrically formed **5** by precipitation and several washing cycles [(S)-**3** \rightarrow (S)-**3**-3,3-DMO, Scheme 1]. Alternatively, dimethyl sulfide (DMS) worked equally well, and (S)-**3**-DMS was isolated with less than 1% tin contamination [(S)-**3** \rightarrow (S)-**3**-DMS, Scheme 1; see the Supporting Information for the molecular structure of (S)-**3**-DMS]. Although both complexes of (S)-**3** as well as the free borane (S)-**3** (burdened with **5**) induced similar levels of enantioselection (*vide infra*), we decided to continue with (S)-**3**-DMS.

Scheme 1. Catalyst Preparation



With catalyst (S)-3-DMS, we tested representative hydrosilanes as reductants in the hydrosilylation of acetophenone [6 \rightarrow (S)-7, Table 1, entries 1–8]. Not surprisingly, monohydrosilanes including EtMe_2SiH were too sterically hindered. This was also true for Ph_2SiH_2 but with MePhSiH_2 carbonyl compound 6 was fully converted within two days, affording (S)-7 with 28% ee. Trihydrosilanes such as PhSiH_3 and MeSiH_3 performed even better, reaching enantioselectivities of 81 and 87% ee, respectively; *t*-BuSiH₃ did not react. The use of an equimolar amount of the trihydrosilane is likely to be detrimental to enantioinduction as intermediate chiral alkoxy-substituted hydrosilanes will potentially act as reductants (Table 1, entry 6 vs entries 9–12). With 2.0 instead of 1.0 equiv of PhSiH_3 , the level of enantiocontrol was improved to 93% ee. No further increase of the enantiomeric excess was seen with more hydrosilane. It made no difference whether the catalysis was run in 1,2- $\text{F}_2\text{C}_6\text{H}_4$ as solvent or neat. The catalyst loading also had an effect on conversion and enantioinduction (for a discussion of this observation, see below):^{7c} 1.0 mol % resulted in 80% ee at 42% conversion and 5.0 mol % yielded 94% ee at full conversion (both within one day; see the Supporting Information for details).

Table 1. Optimization of the Carbonyl Hydrosilylation

entry	hydrosilane	equiv	solvent	conv. (%) ^a	ee (%) ^b
1	Ph_3SiH	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	—	—
2	Me_2PhSiH	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	—	—
3	EtMe_2SiH	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	—	—
4	Ph_2SiH_2	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	—	—
5	MePhSiH_2	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	quant.	28
6	PhSiH_3	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	quant.	81
7	MeSiH_3	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	quant.	87
8	<i>t</i> -BuSiH ₃	1.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	traces	—
9	PhSiH_3	2.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	quant.	93
10	PhSiH_3	3.0	1,2- $\text{F}_2\text{C}_6\text{H}_4$	quant.	93
11	PhSiH_3	3.0	neat	quant.	93
12	PhSiH_3	5.0	neat	quant.	93


^aDetermined by GLC analysis using tetracosane as internal standard. ^bDetermined by HPLC analysis using a chiral stationary phase.

Application of the optimized procedure (Table 1, entry 11) to ether-coordinated (S)-3-3,3-DMO required double the reaction time; the enantiomeric excess (90% ee) was in the same range. Importantly, the free borane (S)-3 in an almost equimolar mixture with tin byproduct 5 catalyzed the hydrosilylation as efficiently as (S)-3-DMS with 92% ee. Hence, we believe that neither Lewis base interferes in the reaction as is the case with phosphine additives.^{11,12}

We then examined various electronically modified acetophenone derivatives 8–14, and the effects were substantial (Scheme 2). Compared to parent 6 (full conversion in one day), CF_3 - and NO_2 -substituted 8 and 9 were less reactive (four and two days, respectively). Conversely, MeO -substituted 14 showed full conversion in one day. The difference in enantioselectivity between these extremes was even more pronounced with 80% ee for 9 but 28% ee for 14. Remarkably, the carboxyl group in 10 was tolerated, and impressive 98% ee in 87% isolated yield were reached. The results for Cl- and Me-substituted 11 and 12 were also satisfactory but the enantiomeric excess collapsed for the corresponding Ph-substituted acetophenone 13. Steric hindrance was also detrimental to both reactivity and enantioinduction; mesityl-substituted 15 afforded 24% conversion after four days and no better than 17% ee. A similar result was obtained from the systematic investigation of the three regioisomeric monobrominated acetophenones 16–18. The *para*- and *meta*-substituted compounds 16 and 17 reacted as good as acetophenone itself, and the level of enantioselection was excellent (95% ee and 99%, respectively). However, *ortho* substitution as in 18 again slowed down the hydrosilylation, and enantioinduction dropped to 73% ee. Replacing the methyl by a benzyl or cyclohexyl group at the carbonyl carbon atom

led to far less reactive **19** and **20**, and asymmetric induction gradually decreased with increasing steric bulk. Good enantiomeric excess was seen with β -naphthyl derivative **21**. Likewise, the enantiomeric excess measured for benzophenone **22** as substrate was even lower. These are the current limitations of the method. — The discrepancy between conversion and isolated yield is mainly due to competing deoxygenation, known to occur with $B(C_6F_5)_3$ /hydrosilane combinations.¹⁶

Scheme 2. Scope and Limitations of the Enantioselective Carbonyl Hydrosilylation^{a,b}




^aDetermined by GLC analysis using mesitylene as internal standard. ^bDetermined by HPLC analysis using chiral stationary phases. ^cFor better solubility, double the amount of PhSiH₃ used. ^dAddition of 1,2-F₂C₆H₄ as solvent to secure homogeneous solution.

Both α -diketones and α -keto esters were the privileged substrates for Du's catalytic system (S)-**2**/HB(C₆F₅)₂ in the presence of Cy₃P (91% yield and >99% ee).¹² Remarkably, benzil was completely inert against our catalyst (S)-**3**-DMS, and ethyl phenylglyoxylate was reduced within four days yet without any enantioselectivity (not shown). These results emphasize that Du's and our catalyst system are complementary.

We had observed before that conversion and enantiomeric excess are dependent on the amount of catalyst employed (see above). Moreover, several of the hydrosilylations displayed slightly deviating values for the enantiomeric excess with 2.4 mol % fixed catalyst loading when conversions were lower or higher than those reported in Scheme 2. We interpret these findings on the basis of our recent mechanistic investigation of the cognate ketimine hydrosilylation.^{7c} In this work, we demonstrated that deprotonation of the intermediate α -C-H-acidic silyliminium ion by the unreacted imine substrate forms the corresponding iminium ion. Both iminium ions engage in the enantioselectivity-determining borohydride reduction, resulting in two competing reaction pathways with potentially different stereochemical outcomes. The same scenario applies to the present ketone hydrosilylation where the catalytic cycle proceeds through either a silylcarboxonium ion I or a "hidden" protonated carboxonium ion II (Scheme 3).

Scheme 3. Generation of Another Activated Carbonyl Group by α -Deprotonation (Borohydride Counteranion Omitted for Clarity)



To summarize, we developed an enantioselective variant of Piers' $B(C_6F_5)_3$ -catalyzed carbonyl hydrosilylation.¹ A combination of a binaphthyl-based boron catalyst with just one C₆F₅ group at the boron atom and reactive trihydrosilanes as the stoichiometric reductant were crucial for achieving acceptable conversion (one day) and high enantioselection (up to 99% ee). The new method distinguishes itself from previous FLP-type approaches^{11–13} as no additional Lewis base is needed. As in the original protocol by Piers,¹ the borane catalyst alone promotes the hydrosilylation.

ASSOCIATED CONTENT

Supporting Information

Experimental details, characterization data, as well as ¹H, ¹³C, and ¹⁹F NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

martin.oestreich@tu-berlin.de

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

M.O. is indebted to the Einstein Foundation (Berlin) for an endowed professorship. We thank Dr. Elisabeth Irran (TU Berlin) for the X-ray analysis.

REFERENCES

- (1) Parks, D. J.; Piers, W. E. *J. Am. Chem. Soc.* **1996**, *118*, 9440–9441.
- (2) For reviews, see: (a) Oestreich, M.; Hermeke, J.; Mohr, J. *Chem. Soc. Rev.* **2015**, *44*, 2202–2220. (b) Piers, W. E.; Marwitz, A. J. V.; Mercier, L. G. *Inorg. Chem.* **2011**, *50*, 12252–12262.
- (3) Parks, D. J.; Blackwell, J. M.; Piers, W. E. *J. Org. Chem.* **2000**, *65*, 3090–3098.
- (4) (a) Rendler, S.; Oestreich, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 5997–6000. (b) Sakata, K.; Fujimoto, H. *J. Org. Chem.* **2013**, *78*, 12505–12512. (c) Houghton, A. Y.; Hurmalainen, J.; Mansikkamäki, A.; Piers, W. E.; Tuononen, H. M. *Nat. Chem.* **2014**, *6*, 983–988.
- (5) Mewald, M.; Fröhlich, R.; Oestreich, M. *Chem.–Eur. J.* **2011**, *17*, 9406–9414.
- (6) Mewald, M. Ph.D. Thesis, Westfälische Wilhelms-Universität Münster, Germany, 2012.
- (7) (a) Blackwell, J. M.; Sonmor, E. R.; Scoccitti, T.; Piers, W. E. *Org. Lett.* **2000**, *2*, 3921–3923. (b) Hog, D. T.; Oestreich, M. *Eur. J. Org. Chem.* **2009**, 5047–5056. (c) Hermeke, J.; Mewald, M.; Oestreich, M. *J. Am. Chem. Soc.* **2013**, *135*, 17537–17546.
- (8) (a) Mewald, M.; Oestreich, M. *Chem.–Eur. J.* **2012**, *18*, 14079–14084. For the enantioselective hydrosilylation catalyzed by (S)-2 (R = 3,5-(3,5-*t*Bu₂C₆H₃)₂; ≤82% ee), see: (b) Zhu, X.; Du, H. *Org. Biomol. Chem.* **2015**, *13*, 1013–1016.
- (9) For authoritative reviews, see: (a) Stephan, D. W. *J. Am. Chem. Soc.* **2015**, *137*, 10018–10032. (b) Stephan, D. W.; Erker, G. *Angew. Chem., Int. Ed.* **2015**, *54*, 6400–6441.
- (10) Asymmetric FLP-type hydrogenation is closely connected to the present hydrosilylation, and Klankermayer's and Du's chiral boranes have been applied in both transformations. For reviews, see: (a) Feng, X.; Du, H. *Tetrahedron Lett.* **2014**, *55*, 6959–6964. (b) Chen, D.; Klankermayer, J. In *Frustrated Lewis Pairs II, Expanding the Scope*; Erker, G., Stephan, D. W., Eds.; Topics in Current Chemistry 334; Springer: Heidelberg, Germany, 2013; pp 1–26. For asymmetric imine hydrogenation, see: (c) Chen, D.; Wang, Y.; Klankermayer, J. *Angew. Chem., Int. Ed.* **2010**, *49*, 9475–9478. (d) Sumerin, V.; Chernichenko, K.; Nieger, M.; Leskelä, M.; Rieger, B.; Repo, T. *Adv. Synth. Catal.* **2011**, *353*, 2093–2110. (e) Stephan, D. W.; Greenberg, S.; Graham, T. W.; Chase, P.; Hastie, J. J.; Geier, S. J.; Farrell, J. M.; Brown, C. C.; Heiden, Z. M.; Welch, G. C.; Ullrich, M. *Inorg. Chem.* **2011**, *50*, 12338–12348. (f) Ghattas, G.; Chen, D.; Pan, F.; Klankermayer, J. *Dalton Trans.* **2012**, 9026–9028.
- (11) Chen, D.; Leich, V.; Pan, F.; Klankermayer, J. *Chem.–Eur. J.* **2012**, *18*, 5184–5187.
- (12) Ren, X.; Du, H. *J. Am. Chem. Soc.* **2016**, *138*, 810–813.
- (13) The net outcome of the FLP-type hydrogenation of silyl enol ethers is that of the carbonyl hydrosilylation: (a) Wei, S.; Du, H. *J. Am. Chem. Soc.* **2014**, *136*, 12261–12264. (b) Ren, X.; Li, G.; Wei, S.; Du, H. *Org. Lett.* **2015**, *17*, 990–993.
- (14) For enantioselective imine hydrogenation catalyzed by electron-deficient boranes without the need for a bulky phosphine as an additive, see: (a) Chen, D.; Klankermayer, J. *Chem. Commun.* **2008**, 2130–2131. (b) Liu, Y.; Du, H. *J. Am. Chem. Soc.* **2013**, *135*, 6810–6813.
- (15) Hermeke, J.; Mewald, M.; Irran, E.; Oestreich, M. *Organometallics* **2014**, *33*, 5097–5100.
- (16) (a) Gevorgyan, V.; Rubin, M.; Benson, S.; Liu, J.-X.; Yamamoto, Y. *J. Org. Chem.* **2000**, *65*, 6179–6186. (b) Nimmagadda, R. D.; McRae, C. *Tetrahedron Lett.* **2006**, *47*, 5755–5758.

