NEPHLUX

Who should get the kidney test first?

Machine learning–based albuminuria risk stratification from PhilHealth YAKAP labs, for early kidney damage.

PJDSC 2026 · Public Health Analytics · Set in Cagayan de Oro City

Stock photo: Josephus Bajo, Malabuyoc, Cebu (Unsplash). Not a patient.

Dialysis is PhilHealth's top-paid procedure.

Hemodialysis led PhilHealth's medical procedure payments in fiscal year 2024 (PhilHealth, 2025).

₱990,600

a year for one patient: **₱6,350** a session × **156** sessions (PhilHealth, 2024b)

<10%

of Filipino adults with CKD have been diagnosed, so most find out late (Orejas, 2024)

Stock photo: Bas Peperzak, Cebu City (Unsplash). Not a patient.

In Cagayan de Oro, kidney disease arrives at stage 5.

One year of adult admissions at Northern Mindanao Medical Center, May 2024 to May 2025.

1,207
CKD admissions, of 32,378 adults

580
from Cagayan de Oro City, the largest share of any city or town

82.4%
arrived at **stage 5**

98.4%
were **already on dialysis**

Leading causes: high blood pressure **35.7%**, diabetes **34.8%**.

Hospital admissions, not a city rate (Mambering & Espiga-Comonsad, 2026). No published source gives CKD by Philippine city or barangay.

YAKAP pays for the labs, but not the test that finds early damage.

YAKAP primary care lab list (PhilHealth, 2024a)

- HbA1c
- Creatinine
- Lipid profile
- CBC with platelets
- Urinalysis (dipstick)
- FBS
- OGTT
- ECG
- Chest X-ray
- Sputum
- Ultrasound
- Mammogram
- Pap smear
- Fecalysis
- FOBT
- uACR · not covered

Highlighted: the YAKAP results NEPHLUX reads. The urine albumin-creatinine ratio (uACR) is the KDIGO test for early kidney damage.

85%

of adults with uACR ≥30 mg/g had a **normal-looking creatinine** (eGFR ≥60) in U.S. survey data

24%

had **neither diabetes nor high blood pressure**, so the usual rule never sends them

A YAKAP clinic cannot send everyone for a uACR test, and has no evidence-based way to choose who goes first.

Rank who gets the uACR test, from results the clinic already has.

The nurse or doctor enters the labs. The doctor decides.

1
Resident check
10 questions on a phone, no labs. Awareness, plus a clinic code.

2
YAKAP check-up
BP, HbA1c, creatinine, lipids, CBC and the urine dipstick YAKAP already covers.

3
Step 2U model
Chance of uACR ≥30 mg/g and its top reasons. No dipstick? It uses Step 2, labs only.

4
Doctor decides
Approve, override with a reason, or no referral. Dipstick protein 1+ always means refer.

5
uACR test
The one test YAKAP does not cover. The clinic follows up its own referrals.

Per 100 adults in the Korean test set, with the same referrals as today's rule



We wrote the pass mark before we trained anything.

Data

Korea's national health survey (KNHANES)

It measures the dipstick **and** uACR on the same adults. The U.S. survey has no dipstick.

Learned on **2021–2024**: **20,465** adults.

Tested once on **2019–2020**: **11,047** adults, 912 cases.

Model

Penalised logistic regression

Inputs: age, sex, history, BMI, measured BP, HbA1c, eGFR (CKD-EPI 2021), cholesterol, blood count, and dipstick protein, blood, glucose and specific gravity.

Simple on purpose: every reason can be shown to the doctor.

Pass mark, written first

Fixed before the test years were opened

1. AUROC 95% CI above 0.70.

2. More cases than "diabetes, high BP or dipstick 1+" at the same number of referrals.

3. Better than the same model without the dipstick.

4. Calibration slope 0.8–1.2.

Script and model hashed (SHA-256) before the one test run. 1,000 bootstrap resamples. Reported with TRIPOD+AI. Adults 20+, not pregnant, no known kidney disease.

STEP 2U · KOREAN TEST SET · SCORED ONCE

+94

more cases found than referring everyone with diabetes, high BP or dipstick protein 1+, with the same number of uACR tests.

Of 912 cases among 11,047 Korean adults (KNHANES 2019–2020) · 95% CI +72 to +118 · 39.4% referred by both · sensitivity 92.8% vs 82.5%

The biggest gain is in people today's rule never sends.

No diabetes and no high blood pressure · 244 cases

80% caught by **NEPHLUX**

34% by today's rule

More than 1 in 4 cases in the test set had neither condition. The dipstick and the labs find them; a yes/no history question cannot.



Share of cases caught, same number of referrals (39.4%)

KNHANES 2019–2020 test set, 11,047 Korean adults, 912 cases.

When tests are scarce, NEPHLUX finds four times the cases.

With 50 uACR tests per 1,000 adults

NEPHLUX order

39.7

cases found · about **₱755** per case

Today's rule or dipstick first

9.1

cases found · about ₱3,281 per case



KNHANES 2019–2020 test set. uACR price assumed at ₱600 (range ₱300–1,200); we found no PhilHealth rate. Cases found, not dialysis avoided.

It ranks better than every simple rule, and its percentages hold.



AUROC, 95% bootstrap CI. Higher means better ranking of who has albuminuria.

0.917

AUROC (95% CI 0.909–0.926). Survey-weighted: 0.927.

1.02

calibration slope (0.96–1.07): a **10%** estimate means about 10%. Mean predicted 8.5%, observed 8.3%.

+0.131

AUROC added by the YAKAP dipstick (0.116–0.147).

KNHANES 2019–2020, 11,047 Korean adults, frozen model, scored once. All four pass-mark criteria met.

The model suggests. The doctor decides.

- A nurse or doctor enters **labs, BP and the YAKAP dipstick** in one form.
- A chance and its **top reasons**, nothing hidden.
- A missing lab **stops the estimate**; nothing is guessed.
- Approve, **override with a reason**, or decline.
- Prints the uACR referral slip.

Mang Ramon, 58, fictional. Dipstick negative, creatinine normal-looking (eGFR 78). Step 2U: **about 10%**, well above the 2.55% cut. Refer.



One web app, three people, one job each.

Resident

10 questions in English, Bisaya or Tagalog. A plain result, a 9-character clinic code and the nearest of 42 YAKAP clinics. Nothing is saved or sent.

YAKAP clinic, nurse or doctor

Loads the code and enters the labs and dipstick. The doctor decides and prints the slip. The clinic follows up referrals with no result after 14 days.

City Health Office

Anonymous counts for 80 barangays, a test budget slider and outreach priorities.



Stock photo: Rainier Ridao, Cagayan de Oro City (Unsplash). Not a patient.

From a phone at home to a referral slip.

DARK MODE



Resident. Two minutes, no account. A "higher" or "lower" chance, never a diagnosis.

LIGHT MODE

PROTOTYPE · SAMPLE RECORD · NOT FOR CLINICAL USE

Referral slip: urine albumin-creatinine ratio (uACR) test

Date 2026-10-02 · Seen at City Health Office (CHO) (public YAKAP clinic), Cagayan de Oro City

Patient name	_____ (write by hand; the app keeps initials only)
Patient	R.S., 58 y, M, Bonbon, Cagayan de Oro City
Resident code	MSJ-5WP-N1Y (answers confirmed with the patient)
BP	148/88 mmHg
HbA1c	7.4%
Creatinine / eGFR	1.10 mg/dL / 78 mL/min/1.73m² (CKD-EPI 2021)
Urine dipstick	protein negative · blood negative · glucose trace · SG 1.020
Screening estimate	10% estimated chance of uACR ≥30 mg/g (higher risk; Step 2U, labs + dipstick)

Main reasons: Systolic blood pressure ↑ · Diabetes diagnosis ↑ · Urine dipstick protein ↓ · HbA1c (blood sugar control) ↑
Request: spot uACR, first-morning sample preferred. A single result ≥30 mg/g is **suspected albuminuria**, not a diagnosis:

Clinic. The slip carries the estimate, the reasons and the repeat-test rule. The doctor signs it.

The clinic code **is** the resident's answers packed into 9 characters with a check value, so nothing is typed twice and nothing is stored.

Every referral becomes local evidence the city does not have yet.



Today, in the app

Each clinic sends the City Health Office **counts only**: screened, higher risk, referred, uACR done, uACR raised, for all 80 barangays. Cells of 1 to 4 are hidden.

Next, with consent

Each returned uACR is a Filipino record with **labs, dipstick and uACR together**: the data needed to test NEPHLUX on Filipinos, and a start toward a Philippine kidney dataset held by the city.

Consent under the Data Privacy Act of 2012 · the CHO as data controller · research extracts only with ethics approval. Clinic visitors, not a survey sample: it shows screening reach, not prevalence.

Dashboard records are SAMPLE data. Clinic pins are approximate. Populations: 2020 census (728,402 residents).

Safe by design, with Nephra to help.

Privacy

- The resident check makes **no storage or network calls**; an automated test enforces it. No account needed.
- The city sees **counts, not names**. The clinic that ordered the test follows it up.

Clinical safety

- The doctor decides **every** referral; an override needs a written reason.
- Dipstick protein 1+ means refer, **whatever the score**. eGFR under 60 raises a warning.
- **39** automated app checks pass. The browser model equals the Python model on 300 test patients.

HAPPY

EXPLAINING

CONCERNED

DARK MODE



LIGHT MODE



Nephra, the Orb, picks one of **78 approved answers**, with no language model. **0** unsafe answers on new test questions; danger signs get an emergency answer.

Next: test it on Filipino patients in Cagayan de Oro.

Tested once on Korean adults. The next test set is ours.

1. **Clinical review** by a YAKAP physician and a nephrologist.
2. **Walkthrough** with one YAKAP nurse or doctor on sample data; fix what they find. A usability test across several clinics comes later.
3. **Cagayan de Oro validation:** YAKAP records with uACR from about 2,800–4,400 adults (Riley et al., 2021), with ethics approval and a data-sharing agreement. Recalibrate, then score the frozen model once.
4. **Later:** a City Health Office pilot with a fixed uACR budget and a few barangays (needs their agreement).



Stock photo: JC Gellidon, Manila (Unsplash).

The labs are already paid for. NEPHLUX decides who gets the urine test first.

+94 cases at the same number of tests. AUROC 0.917, pre-registered and scored once on 11,047 Korean adults. Its next test is a Filipino patient.

NEPHLUX ranks who needs the urine test; the doctor decides and the uACR confirms. A working web app (React, Tailwind CSS, Zustand; NestJS and PostgreSQL API). Every figure regenerates from one script. Independent project, not affiliated with PhilHealth.

Stock photo: Jeremy Brady (Unsplash). Not a patient.

References

1. Collins, G. S., et al. (2024). TRIPOD+AI statement. *BMJ, 385*, e078378. https://doi.org/10.1136/bmj-2023-078378
2. Data Privacy Act of 2012, Republic Act No. 10173 (2012).
3. Inker, L. A., et al. (2021). New creatinine- and cystatin C–based equations to estimate GFR without race. *New England Journal of Medicine, 385*(19), 1737–1749. https://doi.org/10.1056/NEJMoa2102953
4. Kidney Disease: Improving Global Outcomes CKD Work Group. (2024). KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney International, 105*(4S), S117–S314. https://doi.org/10.1016/j.kint.2023.10.018
5. Korea Disease Control and Prevention Agency. (n.d.). *Korea National Health and Nutrition Examination Survey (KNHANES), 2019–2024* [Data sets]. https://knhanes.kdca.go.kr
6. Mambering, N., & Espiga-Comonsad, R. G. (2026). WCN26-4150 Geographic distribution and hospital-based incidence and prevalence of chronic kidney disease in patients admitted at a tertiary government hospital [Abstract]. *Kidney International Reports, 11*, 59–62. https://doi.org/10.1016/j.ekir.2026.104771
7. Orejas, T. T. (2024, February 8). PhilHealth, LGUs urged to cover tests for kidney disease. *Inquirer.net*.
8. Philippine Health Insurance Corporation. (2024a). *Annex B: PhilHealth Konsulta benefit table* (PhilHealth Circular No. 2024-0013).
9. Philippine Health Insurance Corporation. (2024b). *Institutionalization of 156 hemodialysis sessions and coverage expansion* (PhilHealth Circular No. 2024-0023).
10. Philippine Health Insurance Corporation. (2025, May 28). *PhilHealth speeds up payments, releases over P592M to NKTI in less than 5 months* [News release].
11. Philippine Health Insurance Corporation. (2026). *List of PhilHealth YAKAP accredited providers as of September 30, 2026* [Data set].
12. Philippine Statistics Authority. (2021, July 7). *2020 Census of Population and Housing population counts declared official by the President* [Press release].
13. Riley, R. D., et al. (2021). Minimum sample size for external validation of a clinical prediction model with a binary outcome. *Statistics in Medicine, 40*(19), 4230–4251. https://doi.org/10.1002/sim.9025
14. Van Calster, B., et al. (2019). Calibration: The Achilles heel of predictive analytics. *BMC Medicine, 17*, Article 230. https://doi.org/10.1186/s12916-019-1466-7
15. Centers for Disease Control and Prevention. (n.d.). *National Health and Nutrition Examination Survey data, 2011–2012 to 2017–March 2020* [Data sets]. https://wwwn.cdc.gov/nchs/nhanes/

Photos: Unsplash License (Josephus Bajo, Bas Peperzak, Rainier Ridao, JC Gellidon, Jeremy Brady). Full reference list in the project paper.