


# $WW \rightarrow qq l \nu$ Benchmark

Justin Anguiano

October 6, 2019

# Introduction / Motivation



$WW$  is a standard process with a large cross-section

- 15 pb in semileptonic channel at 500 GeV

Three central physics issues addressable by this channel are

- Dynamics of the charged triple gauge couplings
- Measurement of W boson mass, width, cross-section, and BR
- Beam polarization measurement

# 500 GeV Samples

$\sqrt{s} = 500 \text{ GeV}$

Total luminosity :  $4000 \text{ fb}^{-1}$

Polarizations:

Pol.	(-0.8,+0.3)	(+0.8,-0.3)	(-0.8,-0.3)	(+0.8,+0.3)
Lum. [ $\text{fb}^{-1}$ ]	1600	1600	400	400

Reco/Sim: ILCSOftv02-00-02 ILD\_15\_o1\_v02

## MC Background Samples (DBD)–

### – 2-fermion

- Z-bhabhag/hadronic/leptonic

### – 4-fermion

- singleW-leptonic
- Zee/vv-leptonic/semileptonic
- singleZsingleWMix-leptonic
- WW-hadronic/leptonic
- ZZ-hadronic/leptonic/semileptonic
- ZZWWMix-hadronic/leptonic

### – 6-fermion

- eeWW, llWW, vvWW, xxWW
- ttbar
- xxxxZ, yyyyZ

### – SM Higgs

- eeH, qqH,  $\mu\mu\text{H}$ ,  $\tau\tau\text{H}$ ,  $\nu\nu\text{H}$

Note: singleW semileptonic is split into  $WW \rightarrow qqe\nu$  signal and also 4f background events that contain an off shell W ( $\pm 10 \text{ GeV}$  to nominal mass) are considered background

# Analysis Approach

## Step 1-

Treat all lepton flavors universally

Start by identifying signal tau candidates with TauFinder

Optimize TauFinder to efficiently find taus (based on decay products) and also reject tau fakes from hadronic jets

Separate into 4 categories:

$$\tau \rightarrow \mu \nu \nu$$

$$\tau \rightarrow e \nu \nu$$

$$\tau \rightarrow \text{hadronic 1-prong}$$

$$\tau \rightarrow \text{hadronic 3-prong}$$

This approach simultaneously optimizes lepton selection for prompt  $\mu/e$

## Step 2-

With a selected lepton, treat the remaining system as hadronic components of  $W \rightarrow qq$

Use y-cut and kinematic cuts on mini-jets to mitigate pileup ( $\gamma\gamma$ )

**Step 3-** Perform basic event selection for multiple polarization scenarios

# (1) TauFinder

## TauFinder basic operation

Processor starts by seeding tau candidates with tracks ordered by  $|P|$

- Particles that fall within a search cone around the seed track are added to the candidate

- Particles are sequentially added until there are no more in the search cone or until acceptance conditions are violated

## Operating criteria

- Search Cone Angle - The opening angle of the search cone for the tau jet [rad]
- Isolation Cone Angle - Outer isolation cone around the search cone of the tau jet [rad]
- Isolation Energy - The total energy allowed within the isolation cone region [GeV]
- Invariant Mass - The upper limit on tau candidate mass [GeV]

# (1) TauFinder Optimization

Optimization of 3 parameters:

- searchCone  $\in [0, 0.15]$  rad with 0.01 rad steps
- isolationCone  $\in [0, 0.15]$  rad with 0.01 rad steps
- isolationEnergy  $\in [0, 5.5]$  GeV with 0.5 GeV steps

For simplicity, fix invariant mass cut at 3 GeV

Define optimization metrics:

Efficiency using  $WW \rightarrow qq\nu n$  for true leptons:

$$\varepsilon_s = N_{\text{matched}} / N_{\text{Total}}$$

- a tau candidate is considered matched within 100 mrad of the gen lepton
- if the gen lepton is a tau, the jet is matched to the gen visible components – excluding FSR
- $N_{\text{Total}}$  is the Number of events with 3 visible gen fermions  $|\cos\theta| < 0.99$

The optimal working point is chosen from the two tuning parameters

$$\max[(1 - P_{\text{fake}})\varepsilon_s]$$

fake leptons: Use  $WW \rightarrow qqqq$

$$-\varepsilon_b = N_b / N_{\text{Total}}$$

$N_b$  is any event with at least one reconstructed tau jet

4 quarks give 4 chances to create a tau jet  $\varepsilon_b$

– Use a better tuning parameter  $P_{\text{fake}}$  which is the probability of reconstructing a tau jet from a single quark jet

$$P_{\text{fake}} = 1 - (1 - \varepsilon_b)^{\frac{1}{4}}$$
$$\sigma_{P_{\text{fake}}} = \frac{1}{4} \sqrt{\frac{\varepsilon_b}{N_{\text{Total}} \sqrt{1 - \varepsilon_b}}}$$

# (1) TauFinder Optimization Results

Channel	$\varepsilon_s$	$1 - P_{fake}$	% Matched	searchCone [rad]	isoCone [rad]	isoE [GeV]
Prompt $\mu$	0.905	0.974	0.992	0.03	0.15	3.0
Inclusive $\tau$	0.736	0.943	0.958	0.07	0.15	4.5
$\tau \rightarrow \nu\nu\mu$	0.802	0.974	0.984	0.03	0.15	3.0
$\tau \rightarrow \nu\nu e$	0.781	0.963	0.981	0.05	0.15	3.5
$\tau$ Had-1p	0.707	0.943	0.951	0.07	0.15	4.5
$\tau$ Had-3p	0.709	0.930	0.937	0.07	0.15	5.5
Prompt $e$	0.839	0.961	0.97	0.04	0.15	4.0

## (2) Hadronic System


If a lepton(s) have been found by TauFinder,

- select highest energy candidate as signal lepton
- shuffle remaining fakes back into the hadronic system.

At least one quark tends to be very forward, so pileup tends to mix into the jets  
These beam particles cannot be cleanly removed by standard methods e.g. kT algorithm tuned R values

My approach: “Jet Fragmentation”

- tune  $y\text{-cut}(\propto M_{jet}^2)$  values on the durham algorithm (eekt)
- apply simple cuts to the resulting “mini-jets”





## (2) Optimized W Mass

Find best W jet parameters over the ranges:

Use only signal events with gen. prompt muons

- $y_{\text{Cut}}: [1 \times 10^{-3}, 5 \times 10^{-6}]$
- $p_T: [0, 5]$  bins of 0.5 GeV
- $|\cos\theta|: [0.9, 1]$  0.01 bins

Use 2 optimization parameters from the  $M_{qq}^{meas} - M_{qq}^{gen}$  dist.:

- Full Width Half Maximum (FWHM)
- Number of bin Entries in the Mode


The Mode Entries is the number of entries in the Maximum bin + the number of Entries of the nearest left/right neighbor bins

The Mode is the weighted mean of the center of the 3 Mode bins

The Maximum for the FWHM is the "Mode Average" or the average number of entries from the 3 mode bins


The edges of the Width for FWHM are the weighted average between the 2 bins around the half maximum ( 1 bin above 1 bin below)

## (2) Hadronic System Results



Comparison of 3 YCuts with the same kinematic cuts  $P_t > 2$  GeV AND  $\cos\theta < 1$  (optimized for 5e-05)

Small peak around -80 GeV is where the W has been incorrectly thrown out



Significant Improvement !

Mass Difference Statistics:

ycut: 0.001 ptcut: 2 costcut: 1 FWHM: 11.769 RMS: 24.1855 Mode: -0.24211 mean: 0.782898 modeEnt: 5199

ycut: 5e-05 ptcut: 2 costcut: 1 FWHM: 9.7087 RMS: 25.2774 Mode: -0.25127 mean: -3.09776 modeEnt: 6326

ycut: 5e-06 ptcut: 2 costcut: 1 FWHM: 11.567 RMS: 25.7475 Mode: -1.75521 mean: -9.57673 modeEnt: 5475

Best Performance is reached with:

**ycut= 5e-05 and removal of mini-jets with  $p_T < 2$  GeV**

### (3) Event Selection Overview

Perform event selection with two mutually exclusive groups:

1st group will use  $\mu$  cone (optimized for prompt muons)

- "tight" selection will yield some efficiency  $\epsilon_0$  and purity  $p_0$
- tight cuts will be targeted towards prompt signal leptons  $\mu/e$

2nd group will use the  $\tau$  cone (optimized for inclusive  $\tau$  decays)

- "loose" selection will yield some efficiency  $\epsilon_1$  and purity  $p_1$
- "loose" cuts should address  $\tau$ s not reconstructed by muon cone
- orthogonalize selection require 0 tight leptons in loose selection

Optimize selection for some overall efficiency  $\epsilon = \epsilon_0 + \epsilon_1$  times purity

$$p = (N_0 + N_1)/(B_0 + B_1 + N_0 + N_1)$$

Description of current cuts:(currently tight/loose are mostly the same)

adapted from ref. I.Marchesini DESY-THESIS2011


–Note reconstucted particles are boosted against crossing angle boost– (3.5 GeV in x)

- Lepton - Require at least 1 reconstructed lepton
- Track Multiplicity  $> 10$  - at least 10 tracks in the event (targeting rejection of 2f Bkg.)
- Pt  $> 5$  GeV - reject events with no genuine missing Pt
- $E_{vis} < 500$  GeV Sum of the total visible energy in the event
- $E_{com} > 100$  GeV - target rejection of 2f and leptonic eeZ,  
 $E_{com} = E_{vis} + E_{miss}$   $P_{miss}^{\mu} = (|P_{miss}|, -\sum \vec{p}_{vis})$
- $40 < M_{qq} < 120$  - constrains jet system to be W-like
- $-q\cos\theta_W$  - require the  $W^-$  to scatter forward


### (3) Event Selection

Tight Signal  $\Rightarrow$  muon cone for  $\mu, e, \tau$  signal events

Loose Signal  $\Rightarrow$  tau cone for  $\mu, e, \tau$  signal events with no reconstructed tight lepton



N Leptons > 0




N Tracks > 10


### (3) Event Selection

Tight Signal  $\Rightarrow$  muon cone for  $\mu, e, \tau$  signal events

Loose Signal  $\Rightarrow$  tau cone for  $\mu, e, \tau$  signal events with no reconstructed tight lepton



Visible Pt  $> 5$  GeV




Visible Energy  $< 500$  GeV


### (3) Event Selection

Tight Signal  $\Rightarrow$  muon cone for  $\mu, e, \tau$  signal events

Loose Signal  $\Rightarrow$  tau cone for  $\mu, e, \tau$  signal events with no reconstructed tight lepton



$$E_{com} > 100 \text{ GeV}$$




$$40 < M_{qq} < 120$$

### (3) Event Selection

Tight Signal  $\Rightarrow$  muon cone for  $\mu, e, \tau$  signal events

Loose Signal  $\Rightarrow$  tau cone for  $\mu, e, \tau$  signal events with no reconstructed tight lepton

W Scattering angle



$$-q \cos \theta_W > -0.95$$

# (3) Event Selection

Polarization:  $(-0.8, +0.3)$  Luminosity:  $1600 \text{ fb}^{-1}$

Tight Selection with muon cone

	Prompt $\mu$	Prompt $e$	$\tau$	Tot. Sig.	2f	4f	6f	Higgs
nocut	$3.87 \times 10^6$	$3.89 \times 10^6$	$3.90 \times 10^6$	$1.17 \times 10^7$	$4.22 \times 10^7$	$3.22 \times 10^7$	$2.14 \times 10^5$	$4.12 \times 10^5$
lepton	$3.31 \times 10^6$	$3.20 \times 10^6$	$2.28 \times 10^6$	$8.78 \times 10^6$	$1.15 \times 10^7$	$1.18 \times 10^7$	$1.63 \times 10^5$	$1.15 \times 10^5$
ntracks	$3.22 \times 10^6$	$3.11 \times 10^6$	$2.21 \times 10^6$	$8.54 \times 10^6$	$2.84 \times 10^6$	$2.83 \times 10^6$	$1.50 \times 10^5$	$9.36 \times 10^4$
ptcut	$3.18 \times 10^6$	$3.07 \times 10^6$	$2.18 \times 10^6$	$8.44 \times 10^6$	$1.81 \times 10^6$	$2.08 \times 10^6$	$1.47 \times 10^5$	$8.40 \times 10^4$
esum	$3.16 \times 10^6$	$3.00 \times 10^6$	$2.18 \times 10^6$	$8.33 \times 10^6$	$1.66 \times 10^6$	$1.96 \times 10^6$	$1.46 \times 10^5$	$8.11 \times 10^4$
roots	$3.15 \times 10^6$	$2.99 \times 10^6$	$2.16 \times 10^6$	$8.31 \times 10^6$	$1.37 \times 10^6$	$1.63 \times 10^6$	$1.45 \times 10^5$	$7.96 \times 10^4$
mwhad	$2.70 \times 10^6$	$2.56 \times 10^6$	$1.82 \times 10^6$	$7.08 \times 10^6$	$4.02 \times 10^5$	$2.67 \times 10^5$	$2.08 \times 10^4$	$3.06 \times 10^4$
qcostw	$2.69 \times 10^6$	$2.55 \times 10^6$	$1.81 \times 10^6$	$7.05 \times 10^6$	$3.21 \times 10^5$	$2.37 \times 10^5$	$2.01 \times 10^4$	$2.94 \times 10^4$
$\epsilon$	$0.6951 \pm 0.00023$	$0.6551 \pm 0.00024$	$0.4644 \pm 0.00025$	$0.6047 \pm 0.00014$	$0.007589 \pm 1.3e-05$	$0.007363 \pm 1.5e-05$	$0.09395 \pm 0.00063$	$0.0714 \pm 0.0004$

Loose selection with tau cone

	Prompt $\mu$	Prompt $e$	$\tau$	Tot. Sig.	2f	4f	6f	Higgs
nocut	$3.87 \times 10^6$	$3.89 \times 10^6$	$3.90 \times 10^6$	$1.17 \times 10^7$	$4.22 \times 10^7$	$3.22 \times 10^7$	$2.14 \times 10^5$	$4.12 \times 10^5$
lepton	$3.36 \times 10^6$	$3.30 \times 10^6$	$2.82 \times 10^6$	$9.48 \times 10^6$	$1.30 \times 10^7$	$1.36 \times 10^7$	$1.77 \times 10^5$	$1.38 \times 10^5$
mucone	$7.72 \times 10^4$	$1.28 \times 10^5$	$5.70 \times 10^5$	$7.76 \times 10^5$	$1.93 \times 10^6$	$2.15 \times 10^6$	$1.61 \times 10^4$	$3.12 \times 10^4$
ntracks	$7.45 \times 10^4$	$1.23 \times 10^5$	$5.55 \times 10^5$	$7.52 \times 10^5$	$1.61 \times 10^6$	$1.85 \times 10^6$	$1.58 \times 10^4$	$2.81 \times 10^4$
ptcut	$7.34 \times 10^4$	$1.22 \times 10^5$	$5.48 \times 10^5$	$7.43 \times 10^5$	$9.22 \times 10^5$	$1.12 \times 10^6$	$1.36 \times 10^4$	$2.52 \times 10^4$
esum	$7.27 \times 10^4$	$1.19 \times 10^5$	$5.48 \times 10^5$	$7.40 \times 10^5$	$8.75 \times 10^5$	$1.02 \times 10^6$	$1.32 \times 10^4$	$2.46 \times 10^4$
roots	$7.04 \times 10^4$	$1.18 \times 10^5$	$5.41 \times 10^5$	$7.29 \times 10^5$	$7.33 \times 10^5$	$9.83 \times 10^5$	$1.32 \times 10^4$	$2.43 \times 10^4$
mwhad	$4.54 \times 10^4$	$8.08 \times 10^4$	$4.22 \times 10^5$	$5.48 \times 10^5$	$1.85 \times 10^5$	$1.18 \times 10^5$	$1.15 \times 10^3$	$1.28 \times 10^4$
qcostw	$4.00 \times 10^4$	$7.74 \times 10^4$	$4.12 \times 10^5$	$5.29 \times 10^5$	$1.17 \times 10^5$	$1.01 \times 10^5$	$1.11 \times 10^3$	$1.23 \times 10^4$
$\epsilon$	$0.01032 \pm 5.1e-05$	$0.01991 \pm 7.1e-05$	$0.1057 \pm 0.00016$	$0.0454 \pm 6.1e-05$	$0.002775 \pm 8.1e-06$	$0.003146 \pm 9.9e-06$	$0.005167 \pm 0.00015$	$0.0299 \pm 0.00027$



### (3) Event Selection Summary

tight selection

	Prompt $\mu$ O.S.	Prompt $e$ O.S.	Tau O.S.
nocut	$1.15 \times 10^6$	$3.88 \times 10^6$	$1.15 \times 10^6$
lepton	$8.53 \times 10^5$	$2.27 \times 10^6$	$8.53 \times 10^5$
ntracks	$8.34 \times 10^5$	$2.21 \times 10^6$	$8.34 \times 10^5$
ptcut	$8.24 \times 10^5$	$2.19 \times 10^6$	$8.24 \times 10^5$
esum	$8.20 \times 10^5$	$2.18 \times 10^6$	$8.20 \times 10^5$
roots	$8.18 \times 10^5$	$2.17 \times 10^6$	$8.18 \times 10^5$
mwhad	$5.78 \times 10^5$	$1.88 \times 10^6$	$5.78 \times 10^5$
qcostw	$5.76 \times 10^5$	$1.80 \times 10^6$	$5.76 \times 10^5$
$\epsilon$	$0.1487 \pm 0.00047$	$0.4637 \pm 0.00025$	$0.5018 \pm 0.00047$
	Prompt $\mu$ O.S.	Prompt $e$ O.S.	Tau O.S.
nocut	$1.15 \times 10^6$	$3.88 \times 10^6$	$1.15 \times 10^6$
lepton	$9.41 \times 10^5$	$2.47 \times 10^6$	$9.41 \times 10^5$
mucone	$9.65 \times 10^4$	$2.61 \times 10^5$	$9.65 \times 10^4$
ntracks	$9.42 \times 10^4$	$2.48 \times 10^5$	$9.42 \times 10^4$
ptcut	$9.31 \times 10^4$	$2.46 \times 10^5$	$9.31 \times 10^4$
esum	$9.28 \times 10^4$	$2.46 \times 10^5$	$9.28 \times 10^4$
roots	$9.21 \times 10^4$	$2.35 \times 10^5$	$9.21 \times 10^4$
mwhad	$5.30 \times 10^4$	$1.28 \times 10^5$	$5.30 \times 10^4$
qcostw	$5.13 \times 10^4$	$1.18 \times 10^5$	$5.13 \times 10^4$
$\epsilon$	$0.01326 \pm 0.00019$	$0.03039 \pm 8.7e-05$	$0.04475 \pm 0.00019$

## RUN 2

	Prompt $\mu$	Prompt $e$	$\tau$	Tot. Sig.	2f	4f	6f	Higgs
nocut	$2.48 \times 10^5$	$2.53 \times 10^5$	$2.50 \times 10^5$	$7.51 \times 10^5$	$2.82 \times 10^7$	$1.36 \times 10^7$	$7.80 \times 10^4$	$1.30 \times 10^5$
lepton	$2.14 \times 10^5$	$2.08 \times 10^5$	$1.46 \times 10^5$	$5.68 \times 10^5$	$9.52 \times 10^6$	$5.99 \times 10^6$	$5.81 \times 10^4$	$4.45 \times 10^4$
ntracks	$2.08 \times 10^5$	$2.02 \times 10^5$	$1.42 \times 10^5$	$5.53 \times 10^5$	$2.00 \times 10^6$	$9.00 \times 10^5$	$5.53 \times 10^4$	$4.04 \times 10^4$
ptcut	$2.06 \times 10^5$	$2.00 \times 10^5$	$1.40 \times 10^5$	$5.47 \times 10^5$	$1.28 \times 10^6$	$7.37 \times 10^5$	$5.39 \times 10^4$	$3.43 \times 10^4$
esum	$2.04 \times 10^5$	$1.96 \times 10^5$	$1.40 \times 10^5$	$5.40 \times 10^5$	$1.19 \times 10^6$	$7.18 \times 10^5$	$5.36 \times 10^4$	$3.24 \times 10^4$
roots	$2.04 \times 10^5$	$1.95 \times 10^5$	$1.39 \times 10^5$	$5.39 \times 10^5$	$9.81 \times 10^5$	$4.23 \times 10^5$	$5.36 \times 10^4$	$3.22 \times 10^4$
mwhad	$1.76 \times 10^5$	$1.67 \times 10^5$	$1.18 \times 10^5$	$4.61 \times 10^5$	$2.90 \times 10^5$	$9.09 \times 10^4$	$7.26 \times 10^3$	$5.38 \times 10^3$
qcostw	$1.75 \times 10^5$	$1.66 \times 10^5$	$1.17 \times 10^5$	$4.58 \times 10^5$	$2.22 \times 10^5$	$7.47 \times 10^4$	$6.84 \times 10^3$	$5.21 \times 10^3$
$\epsilon$	$0.7037 \pm 0.00092$	$0.6584 \pm 0.00094$	$0.4674 \pm 0.001$	$0.6099 \pm 0.00056$	$0.007886 \pm 1.7e-05$	$0.005502 \pm 2e-05$	$0.08772 \pm 0.001$	$0.03996 \pm 0.00054$
	Prompt $\mu$	Prompt $e$	$\tau$	Tot. Sig.	2f	4f	6f	Higgs
nocut	$2.48 \times 10^5$	$2.53 \times 10^5$	$2.50 \times 10^5$	$7.51 \times 10^5$	$2.82 \times 10^7$	$1.36 \times 10^7$	$7.80 \times 10^4$	$1.30 \times 10^5$
lepton	$2.17 \times 10^5$	$2.15 \times 10^5$	$1.82 \times 10^5$	$6.14 \times 10^5$	$1.05 \times 10^7$	$6.29 \times 10^6$	$6.34 \times 10^4$	$5.20 \times 10^4$
mucone	$4.82 \times 10^3$	$8.45 \times 10^3$	$3.76 \times 10^4$	$5.09 \times 10^4$	$1.25 \times 10^6$	$3.72 \times 10^5$	$6.32 \times 10^3$	$1.05 \times 10^4$
ntracks	$4.65 \times 10^3$	$8.12 \times 10^3$	$3.66 \times 10^4$	$4.94 \times 10^4$	$1.02 \times 10^6$	$2.03 \times 10^5$	$6.27 \times 10^3$	$1.01 \times 10^4$
ptcut	$4.58 \times 10^3$	$8.04 \times 10^3$	$3.62 \times 10^4$	$4.89 \times 10^4$	$5.77 \times 10^5$	$1.38 \times 10^5$	$5.43 \times 10^3$	$8.17 \times 10^3$
esum	$4.54 \times 10^3$	$7.90 \times 10^3$	$3.62 \times 10^4$	$4.87 \times 10^4$	$5.45 \times 10^5$	$1.29 \times 10^5$	$5.26 \times 10^3$	$7.78 \times 10^3$
roots	$4.40 \times 10^3$	$7.81 \times 10^3$	$3.58 \times 10^4$	$4.80 \times 10^4$	$4.32 \times 10^5$	$1.11 \times 10^5$	$5.25 \times 10^3$	$7.74 \times 10^3$
mwhad	$2.86 \times 10^3$	$5.41 \times 10^3$	$2.81 \times 10^4$	$3.64 \times 10^4$	$1.13 \times 10^5$	$1.74 \times 10^4$	$3.75 \times 10^2$	$1.71 \times 10^3$
qcostw	$2.53 \times 10^3$	$5.18 \times 10^3$	$2.74 \times 10^4$	$3.51 \times 10^4$	$6.39 \times 10^4$	$1.42 \times 10^4$	$3.47 \times 10^2$	$1.66 \times 10^3$
$\epsilon$	$0.01017 \pm 0.0002$	$0.02048 \pm 0.00028$	$0.1099 \pm 0.00063$	$0.04679 \pm 0.00024$	$0.002268 \pm 9e-06$	$0.001048 \pm 8.8e-06$	$0.004447 \pm 0.00024$	$0.01271 \pm 0.00031$

# RUN2 OS

tight

	Prompt $\mu$ O.S.	Prompt $e$ O.S.	Tau O.S.
nocut	$7.80 \times 10^4$	$6.62 \times 10^5$	$7.80 \times 10^4$
lepton	$5.84 \times 10^4$	$2.78 \times 10^5$	$5.84 \times 10^4$
ntracks	$5.71 \times 10^4$	$2.70 \times 10^5$	$5.71 \times 10^4$
ptcut	$5.65 \times 10^4$	$2.68 \times 10^5$	$5.65 \times 10^4$
esum	$5.62 \times 10^4$	$2.67 \times 10^5$	$5.62 \times 10^4$
roots	$5.61 \times 10^4$	$2.65 \times 10^5$	$5.61 \times 10^4$
mwhad	$4.05 \times 10^4$	$2.30 \times 10^5$	$4.05 \times 10^4$
qcostw	$4.03 \times 10^4$	$2.15 \times 10^5$	$4.03 \times 10^4$
$\epsilon$	$0.1623 \pm 0.0018$	$0.3241 \pm 0.00058$	$0.5165 \pm 0.0018$

	Prompt $\mu$ O.S.	Prompt $e$ O.S.	Tau O.S.
nocut	$7.80 \times 10^4$	$6.62 \times 10^5$	$7.80 \times 10^4$
lepton	$6.45 \times 10^4$	$3.25 \times 10^5$	$6.45 \times 10^4$
mucone	$6.64 \times 10^3$	$5.65 \times 10^4$	$6.64 \times 10^3$
ntracks	$6.50 \times 10^3$	$5.33 \times 10^4$	$6.50 \times 10^3$
ptcut	$6.43 \times 10^3$	$5.28 \times 10^4$	$6.43 \times 10^3$
esum	$6.41 \times 10^3$	$5.28 \times 10^4$	$6.41 \times 10^3$
roots	$6.37 \times 10^3$	$5.02 \times 10^4$	$6.37 \times 10^3$
mwhad	$3.83 \times 10^3$	$2.37 \times 10^4$	$3.83 \times 10^3$
qcostw	$3.72 \times 10^3$	$2.16 \times 10^4$	$3.72 \times 10^3$
$\epsilon$	$0.01497 \pm 0.00076$	$0.03269 \pm 0.00022$	$0.04766 \pm 0.00076$