MISMIP+ summary of TKL_HOM_TSAI_1000m

Thomas Kleiner

March 30, 2017

1 Model Details

- 1. model: TIM-FD³ (Kleiner and Humbert, 2014).
- 2. repository: TIM-FD³ source codes are available on request.
- 3. englacial stresses: HOM (including vertical resistive stresses R_{zz}), Glen's law, n=3, $A=2.0\times 10^{-17}\,\mathrm{Pa}^{-3}\,\mathrm{a}^{-1}$
- 4. basal traction: Modified power law (Tsai et al., 2015) as in Asay-Davis et al. (2016, Eqns. 7,9,10) with $\alpha^2 = 0.5$, $\beta^2 = 3.16 \times 10^6 \,\mathrm{Pa}\,\mathrm{m}^{-1/3}\,\mathrm{s}^{1/3} = 10^4 \,\mathrm{Pa}\,\mathrm{m}^{-1/3}\,\mathrm{a}^{1/3}$
- 5. space discretisation: Finite Differences on regular grid, equidistant horizontal grid $(\Delta x = \Delta y = 1000 \,\mathrm{m})$, terrain-following vertical coordinates, 9 vertical layer (refined at the base)
- 6. time stepping: fixed time step, $\Delta t = 0.25 a$
- 7. grounding line: hydrostatic approximation, no sub-grid grounding line treatment in the model, model output is generated along the contour $h h_{\rm f} = 0$ (floating condition) via linear interpolation in the post-processing
- 8. MISMIP3d name: TKL1, different mesh resolution, FS instead of HOM used in MISMIP3d

2 Comments

Spin-up: As initial condition for the spin-up a pre-computed geometry from BISICLES is used (same parameters as above, tsai-A2.0e-17-ssa.nc, provided by S. L. Cornford). To account for different model physics and resolution a 500 years relaxation run was performed.

Mesh resolution: Mesh resolution has been tested on 4 km, 2 km and 1 km resolution with the same set of parameters as in the text above for the model spin-up. Only the 1 km resolution set-up shows a stable grounding line position at the end of the spin-up, while both coarser resolution runs end in a cyclic grounding/ungrounding state with a very stable period.

Ice2r: In all Ice2r-type simulations a minimum ice thickness of $10 \,\mathrm{m}$ is used to preserve a stable calving front at $x = 640 \,\mathrm{km}$.

References

- X. S. Asay-Davis, S. L. Cornford, G. Durand, B. K. Galton-Fenzi, R. M. Gladstone, G. H. Gudmundsson, T. Hattermann, D. M. Holland, D. Holland, P. R. Holland, D. F. Martin, P. Mathiot, F. Pattyn, and H. Seroussi. Experimental design for three interrelated marine ice sheet and ocean model intercomparison projects: MISMIP v. 3 (mismip+), isomip v. 2 (isomip+) and misomip v. 1 (misomip1). Geoscientific Model Development, 9(7):2471–2497, 2016. doi: 10.5194/gmd-9-2471-2016.
- T. Kleiner and A. Humbert. Numerical simulations of major ice streams in western Dronning Maud Land, Antarctica, under wet and dry basal conditions. *Journal of Glaciology*, 60 (220):215–232, 2014. doi: 10.3189/2014JoG13J006.
- V. C. Tsai, A. L. Stewart, and A. F. Thompson. Marine ice-sheet profiles and stability under coulomb basal conditions. *Journal of Glaciology*, 61(226):205–215, 2015. doi: 10.3189/2015jog14j221.