

Higher-dimensional categories: induction on extensivity

Thomas Cottrell¹

Soichiro Fujii²

John Power³

¹Department of Mathematical Sciences, University of Bath

²Department of Computer Science, University of Tokyo

³Department of Computer Science, University of Bath

7 July, 2018

Overview

Leinster's method for defining weak n -categories:

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $\mathcal{T}^{(n)}$, which is cartesian.

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.
- Define a notion of *contraction* on $T^{(n)}$ -operads.

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.
- Define a notion of *contraction* on $T^{(n)}$ -operads.
- A weak n -category is an algebra for the initial $T^{(n)}$ -operad with contraction.

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.
- Define a notion of *contraction* on $T^{(n)}$ -operads.
- A weak n -category is an algebra for the initial $T^{(n)}$ -operad with contraction.

Aims:

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.
- Define a notion of *contraction* on $T^{(n)}$ -operads.
- A weak n -category is an algebra for the initial $T^{(n)}$ -operad with contraction.

Aims:

- Enrich this to define weak n -dimensional \mathcal{V} -categories.

Overview

Leinster's method for defining weak n -categories:

- Start with the free strict n -category monad $T^{(n)}$, which is cartesian.
- For a cartesian monad T on \mathcal{C} , define T -operads as monoids in $\mathcal{C}/T1$.
- Define a notion of *contraction* on $T^{(n)}$ -operads.
- A weak n -category is an algebra for the initial $T^{(n)}$ -operad with contraction.

Aims:

- Enrich this to define weak n -dimensional \mathcal{V} -categories.
- Build dimensions through iterated enrichment.

Extensivity

Definition

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if,

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$,

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod: \prod_{i \in I} (\mathcal{V}/X_i) & \longrightarrow & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod: \prod_{i \in I} (\mathcal{V}/X_i) & \longrightarrow & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod: \prod_{i \in I} (\mathcal{V}/X_i) & \longrightarrow & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Examples:

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod: \prod_{i \in I} (\mathcal{V}/X_i) & \longrightarrow & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Examples: **Set**,

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod_{i \in I} (\mathcal{V}/X_i) & \xrightarrow{\quad} & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Examples: **Set**, ω -**Cpo**,

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod_{i \in I} (\mathcal{V}/X_i) & \xrightarrow{\quad} & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \longmapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Examples: **Set**, ω -**Cpo**, **Cat**,

Extensivity

Definition

A category \mathcal{V} (with small coproducts) is *extensive* if, for any set I and family of objects $(X_i)_{i \in I}$, the functor

$$\begin{array}{ccc} \coprod_{i \in I} (\mathcal{V}/X_i) & \xrightarrow{\quad} & \mathcal{V}/(\coprod_{i \in I} X_i) \\ \left(\begin{array}{c} A_i \\ \downarrow f_i \\ X_i \end{array} \right)_{i \in I} & \mapsto & \begin{array}{c} \coprod_{i \in I} A_i \\ \downarrow \coprod_{i \in I} f_i \\ \coprod_{i \in I} X_i \end{array} \end{array}$$

is an equivalence of categories.

Examples: **Set**, ω -**Cpo**, **Cat**, \mathcal{V} -**Cat** and \mathcal{V} -**Gph** (for extensive \mathcal{V}).

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

For each natural number n , $\mathcal{V}\text{-Cat}^{(n)}$ is defined by:

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

For each natural number n , $\mathcal{V}\text{-Cat}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Cat}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Cat}^{(n+1)} = (\mathcal{V}\text{-Cat}^{(n)})\text{-Cat},$$

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

For each natural number n , $\mathcal{V}\text{-Cat}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Cat}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Cat}^{(n+1)} = (\mathcal{V}\text{-Cat}^{(n)})\text{-Cat},$$

and $\mathcal{V}\text{-Gph}^{(n)}$ is defined by:

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

For each natural number n , $\mathcal{V}\text{-Cat}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Cat}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Cat}^{(n+1)} = (\mathcal{V}\text{-Cat}^{(n)})\text{-Cat},$$

and $\mathcal{V}\text{-Gph}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Gph}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Gph}^{(n+1)} = (\mathcal{V}\text{-Gph}^{(n)})\text{-Gph}.$$

$\mathcal{V}\text{-Cat}^{(n)}$ and $\mathcal{V}\text{-Gph}^{(n)}$

Definition

Let \mathcal{V} be a category with finite products.

For each natural number n , $\mathcal{V}\text{-Cat}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Cat}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Cat}^{(n+1)} = (\mathcal{V}\text{-Cat}^{(n)})\text{-Cat},$$

and $\mathcal{V}\text{-Gph}^{(n)}$ is defined by:

$$\mathcal{V}\text{-Gph}^{(0)} = \mathcal{V}; \quad \mathcal{V}\text{-Gph}^{(n+1)} = (\mathcal{V}\text{-Gph}^{(n)})\text{-Gph}.$$

When $\mathcal{V} = \mathbf{Set}$, $\mathbf{Set}\text{-Cat}^{(n)} = n\text{-Cat}$, the category of strict n -categories.

Extensivity and enrichment

Proposition

If \mathcal{V} is extensive and finitely complete,

Extensivity and enrichment

Proposition

If \mathcal{V} is extensive and finitely complete, then $\mathcal{V}\text{-}\mathbf{Gph}$ and $\mathcal{V}\text{-}\mathbf{Cat}$ are also extensive and finitely complete.

Extensivity and enrichment

Proposition

If \mathcal{V} is extensive and finitely complete, then $\mathcal{V}\text{-}\mathbf{Gph}$ and $\mathcal{V}\text{-}\mathbf{Cat}$ are also extensive and finitely complete.

Corollary

If \mathcal{V} is extensive and finitely complete, then $\mathcal{V}\text{-}\mathbf{Gph}^{(n)}$ and $\mathcal{V}\text{-}\mathbf{Cat}^{(n)}$ are also extensive and finitely complete.

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,
- T preserves pullbacks,

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,
- T preserves pullbacks,
- all the naturality squares for η and μ are pullback squares.

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,
- T preserves pullbacks,
- all the naturality squares for η and μ are pullback squares.

Proposition

Let \mathcal{V} be extensive and finitely complete. For each n , there is an adjunction

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,
- T preserves pullbacks,
- all the naturality squares for η and μ are pullback squares.

Proposition

Let \mathcal{V} be extensive and finitely complete. For each n , there is an adjunction

$$\mathcal{V}\text{-}\mathbf{Gph}^{(n)} \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad \perp \quad} \end{array} \mathcal{V}\text{-}\mathbf{Cat}^{(n)}$$

Cartesian monads

Definition

A monad (T, η, μ) on \mathcal{C} is *cartesian* if

- \mathcal{C} has all pullbacks,
- T preserves pullbacks,
- all the naturality squares for η and μ are pullback squares.

Proposition

Let \mathcal{V} be extensive and finitely complete. For each n , there is an adjunction

$$\mathcal{V}\text{-}\mathbf{Gph}^{(n)} \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad \perp \quad} \end{array} \mathcal{V}\text{-}\mathbf{Cat}^{(n)}$$

and the induced monad $T^{(n)}$ is cartesian.

T -operads

For a cartesian monad T on a finitely complete category \mathcal{C} , $\mathcal{C}/T1$ can be given a monoidal structure.

T -operads

For a cartesian monad T on a finitely complete category \mathcal{C} , $\mathcal{C}/T1$ can be given a monoidal structure.

Definition

A T -operad is a monoid in $\mathcal{C}/T1$.

T -operads

For a cartesian monad T on a finitely complete category \mathcal{C} , $\mathcal{C}/T1$ can be given a monoidal structure.

Definition

A T -operad is a monoid in $\mathcal{C}/T1$.

Given a T -operad $m: M \rightarrow T1$, an *algebra for* (M, m) consists of an object X of \mathcal{C} together with an action of (M, m) on X .

T -operads

For a cartesian monad T on a finitely complete category \mathcal{C} , $\mathcal{C}/T1$ can be given a monoidal structure.

Definition

A T -operad is a monoid in $\mathcal{C}/T1$.

Given a T -operad $m: M \rightarrow T1$, an *algebra for* (M, m) consists of an object X of \mathcal{C} together with an action of (M, m) on X .

- Classical operads: $\mathcal{C} = \mathbf{Set}$, $T = \text{free monoid}$.

T -operads

For a cartesian monad T on a finitely complete category \mathcal{C} , $\mathcal{C}/T1$ can be given a monoidal structure.

Definition

A T -operad is a monoid in $\mathcal{C}/T1$.

Given a T -operad $m: M \rightarrow T1$, an *algebra for (M, m)* consists of an object X of \mathcal{C} together with an action of (M, m) on X .

- Classical operads: $\mathcal{C} = \mathbf{Set}$, $T = \text{free monoid}$.
- For weak n -dimensional \mathcal{V} -categories, use $\mathcal{C} = \mathcal{V}\text{-}\mathbf{Gph}^{(n)}$, $T = T^{(n)}$.

Contractions

Contractions

A *contraction* on $m: M \rightarrow T^{(n)}1$ consists of

Contractions

A *contraction* on $m: M \rightarrow T^{(n)}1$ consists of a lifting

$$\begin{array}{ccc} \partial_j & \xrightarrow{h} & M \\ f_j \downarrow & \nearrow & \downarrow m \\ C_j & \xrightarrow{k} & T^{(n)}1 \end{array}$$

for every such commuting square, where f_j is a “cell boundary inclusion”.

Contractions

A *contraction* on $m: M \rightarrow T^{(n)}1$ consists of a lifting

$$\begin{array}{ccc} \partial_j & \xrightarrow{h} & M \\ f_j \downarrow & \nearrow & \downarrow m \\ C_j & \xrightarrow{k} & T^{(n)}1 \end{array}$$

for every such commuting square, where f_j is a “cell boundary inclusion”.

Definition

Let \mathcal{V} be extensive, finitely complete, and locally presentable.

Contractions

A *contraction* on $m: M \rightarrow T^{(n)}1$ consists of a lifting

$$\begin{array}{ccc} \partial_j & \xrightarrow{h} & M \\ f_j \downarrow & \nearrow & \downarrow m \\ C_j & \xrightarrow{k} & T^{(n)}1 \end{array}$$

for every such commuting square, where f_j is a “cell boundary inclusion”.

Definition

Let \mathcal{V} be extensive, finitely complete, and locally presentable.

A *weak n -dimensional \mathcal{V} -category* is an algebra for the initial $T^{(n)}$ -operad with contraction.

Contractions

A *contraction* on $m: M \rightarrow T^{(n)}1$ consists of a lifting

$$\begin{array}{ccc} \partial_j & \xrightarrow{h} & M \\ f_j \downarrow & \nearrow & \downarrow m \\ C_j & \xrightarrow{k} & T^{(n)}1 \end{array}$$

for every such commuting square, where f_j is a “cell boundary inclusion”.

Definition

Let \mathcal{V} be extensive, finitely complete, and locally presentable.

A *weak n -dimensional \mathcal{V} -category* is an algebra for the initial $T^{(n)}$ -operad with contraction.

For $\mathcal{V} = \mathbf{Set}$, this agrees with Leinster’s definition of weak n -category.