AI for Medicine

From diagnosis and treatment to clinical trials and mental health support

Ruishan Liu

CSCI 544 Guest Lecture · October 20, 2026

USC University of Southern California


Thomas Lord Department of Computer Science
USC Viterbi

USC Dornsife
Department of Quantitative and Computational Biology

Keck Medicine of USC

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning




TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench


ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

Four recent works from our lab, following a patient's path through care — and four different ways people and AI work together

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning



TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench

ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

MedExAgent: Training LLM Agents to Ask, Examine, and Diagnose in Noisy Clinical Environments

Yicheng Gao, Xiaolin Zhou, Yahan Li, Yue Zhao, Ruishan Liu

University of Southern California · Arizona State University

NeurIPS 2026

USC University of Southern California

Thomas Lord Department of Computer Science
USC Viterbi

USC Dornsife
Department of Quantitative and Computational Biology

Keck Medicine of USC

Real Diagnosis Is Sequential, Noisy, and Costly

Sequential

Chief complaint → questions → exams as evidence is needed → commit to a diagnosis

Noisy

Patients misreport location, severity, or duration; lab results can be contaminated, partial, or ambiguous


Costly

From a routine blood draw to an invasive biopsy — different financial cost and patient discomfort

Static benchmarks (MedQA, MedMCQA, case vignettes) hand the model a complete case
— **they capture none of these properties**

Gao et al. *NeurIPS* (2026)


One Exam Changes the Diagnosis



Conversation-only LLMs keep asking about sleep and appetite and settle on depression.
MedExAgent orders a TSH / free T4 panel and reaches the correct diagnosis — hypothyroidism.

Gao et al. *NeurIPS* (2026)


MedExAgent Overview




Environment: a partially observable Markov decision process (POMDP)

Gao et al. *NeurIPS* (2026)

MedExAgent Overview






Data: doctor–patient conversations synthesized along the five stages of the **Calgary–Cambridge** clinical interview, with patient and exam noise injected

Gao et al. *NeurIPS* (2026)

MedExAgent Overview



Training: Meditron3-8B → **SFT** on the conversations → **DAPO RL** against a composite reward (accuracy, tool calls, exam cost)

Gao et al. *NeurIPS* (2026)

Seven Patient Noises, Three Exam Noises

	Noise type	Description	Example
Patient	**Body part swap**	Reports symptom in an adjacent body part	“stomach pain” → “liver pain”
Patient	**Symptom confusion**	Related but incorrect descriptor	“burning” → “tingling sensation”
Patient	**Severity change**	Misreports intensity, typically downplaying	“severe headache” → “moderate headache”
Patient	**Temporal change**	Misreports duration or onset	“3 weeks” → “1 week”
Patient	**Omission**	Leaves out one of several symptoms	omits “fatigue” alongside fever and cough
Patient	**Self-diagnosis**	Volunteers an incorrect lay diagnosis	“I looked it up online, it seems like allergies”
Patient	**Vague answer**	Evasive, non-specific reply	“I'm not really sure about that”
Exam	**Body part swap**	Adjacent body part in findings	“inflammation in the stomach lining” → “…liver lining”
Exam	**Exam omission**	Drops one clause from multi-part findings	“elevated WBC; low Hb; normal platelets” → drops “low Hb”
Exam	**Ambiguity**	Wraps findings in equivocal language	“Findings are equivocal — nodule in right lung …”

Gao et al. *NeurIPS* (2026)

A Composite Reward for Diagnosis

$$R = R_{dx} + w_{tool} \cdot R_{tool} - w_{cost} \cdot R_{cost}$$

Diagnosis reward **Tool-call reward** **Exam-cost penalty**

Gao et al. *NeurIPS* (2026)

Data: Typical, Long-Tail, and Held-Out Cases

DDxPlus

~1.3M synthetic patients, 49 pathologies, 110 symptoms

Realistic U.S. base rates → "typical" patients

PMC-Patients-v2

~250K real case reports from PubMed Central


Rare diseases, atypical presentations, multi-condition cases

AgentClinic-MedQA

201 OSCE-style cases grounded in USMLE questions

Held out entirely → out-of-distribution test

5,000 cases for SFT · **20,000** for RL · **2,000** for test — split evenly between DDxPlus and PMC-Patients-v2



SFT conversations follow the five Calgary–Cambridge stages, one of 36 personas per patient

Gao et al. *NeurIPS* (2026)

MedExAgent-8B vs. Seven Baselines

Sim: embedding cosine similarity · Jac: LLM-judge Jaccard over matched conditions · Acc: strict accuracy, all ground-truth conditions matched (Table 1)

	DDxPlus			PMC-Patients-v2			AgentClinic-MedQA (OOD)		
Model	**Sim**	**Jac**	**Acc**	**Sim**	**Jac**	**Acc**	**Sim**	**Jac**	**Acc**
Llama-3.1-8B-Instruct	0.640	0.383	0.605	0.507	0.122	0.157	0.575	0.225	0.298
Qwen3-8B	0.605	0.328	0.460	0.532	0.092	0.120	0.565	0.153	0.219
Qwen3-14B	0.659	0.399	0.538	0.565	0.113	0.143	0.614	0.226	0.284
Qwen3-32B	0.685	0.466	0.626	0.551	0.127	0.157	0.637	0.310	0.368
MedGemma-27B-text-it	0.498	0.241	0.325	0.469	0.108	0.123	0.544	0.214	0.249
Aloe-Beta-70B	0.716	0.503	0.670	0.590	0.171	0.219	**0.684**	**0.395**	**0.395**
HuatuoGPT-o1-70B	0.664	0.410	0.470	0.534	0.120	0.124	0.625	0.238	0.249
MedExAgent-8B (ours)	**0.953**	**0.937**	**0.939**	**0.626**	**0.345**	**0.314**	0.672	0.378	0.378

In distribution: beats every baseline on DDxPlus and PMC-Patients-v2 on all metrics ($p < 0.001$)

Gao et al. *NeurIPS* (2026)

MedExAgent-8B vs. Seven Baselines

Sim: embedding cosine similarity · Jac: LLM-judge Jaccard over matched conditions · Acc: strict accuracy, all ground-truth conditions matched (Table 1)


	DDxPlus			PMC-Patients-v2			AgentClinic-MedQA (OOD)		
Model	**Sim**	**Jac**	**Acc**	**Sim**	**Jac**	**Acc**	**Sim**	**Jac**	**Acc**
Llama-3.1-8B-Instruct	0.640	0.383	0.605	0.507	0.122	0.157	0.575	0.225	0.298
Qwen3-8B	0.605	0.328	0.460	0.532	0.092	0.120	0.565	0.153	0.219
Qwen3-14B	0.659	0.399	0.538	0.565	0.113	0.143	0.614	0.226	0.284
Qwen3-32B	0.685	0.466	0.626	0.551	0.127	0.157	0.637	0.310	0.368
MedGemma-27B-text-it	0.498	0.241	0.325	0.469	0.108	0.123	0.544	0.214	0.249
Aloe-Beta-70B	0.716	0.503	0.670	0.590	0.171	0.219	**0.684**	**0.395**	**0.395**
HuatuoGPT-o1-70B	0.664	0.410	0.470	0.534	0.120	0.124	0.625	0.238	0.249
MedExAgent-8B (ours)	**0.953**	**0.937**	**0.939**	**0.626**	**0.345**	**0.314**	0.672	0.378	0.378

Out of distribution: statistically tied with Aloe-Beta-70B on Sim, Jac and Acc, and with Qwen3-32B on Acc

An 8B specialist matches the strongest 70B medical model at roughly 9× fewer parameters

Gao et al. *NeurIPS* (2026)


8B Matches 70B Out of Distribution




Gao et al. *NeurIPS* (2026)

Summary of MedExAgent

- **A POMDP environment** for interactive diagnosis that unifies asking, exam ordering and diagnosis, with 7 patient + 3 exam noise types and real-world exam costs (CMS fees, invasiveness)
- **A training recipe** — SFT on Calgary–Cambridge-structured conversations, then DAPO against a composite reward over accuracy, tool calls and cost
- **An open 8B specialist** that beats every baseline in distribution and matches 70B medical models out of distribution, while ordering 3–4 well-chosen exams per case





Gao et al. *NeurIPS* (2026)

Code: https://github.com/EndlessCG/medexagent

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning



TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench

ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

From Correction to Collaboration: Reusing Routine Clinician Annotations to Guide AI-Based Tumor Contouring

Shupeng Cheng, Nguyen Phuong Dang, Adam Garsa, Joshua Schiff, Hirsch Matani, Amir Ahari, Ashwin Ganesh, Justin Park, Jason Ye, Zhuoli Zhang, Benjamin Ziemer, Yahan Li, Yutaka Natsuaki, Zhilei Shen, Diane C. Ling, Angel Hsing-Chi Hwang, Eric Lin Chang, Ruishan Liu

USC Computer Science · USC Keck Radiation Oncology · UC Irvine Radiological Sciences · USC Annenberg

TRACE · Tumor contouring Refinement via Annotation-Conditioned Enhancement

In submission

USC University of Southern California


Thomas Lord Department of Computer Science
USC Viterbi

USC Dornsife
Department of Quantitative and Computational Biology

Keck Medicine of USC

Tumor Contouring

3D Images




Contour slice-by-slice



Time-consuming

AI for Tumor Contouring

3D Images




Contour slice-by-slice




TransUNet, UNet++, Attention U-Net, SwinUNet, H2Former, FATNet, MedFormer, …


AI for Tumor Contouring



- Need clinician's review

AI for Tumor Contouring






- Need clinician's review

Accept → Next slice

Reject → Edit / Redraw




(practical)




AI Tumor Contouring

First slice

Traditional AI







AI Tumor Contouring

First slice

Traditional AI







AI Tumor Contouring

First slice

Traditional AI





AI Tumor Contouring



First slice → **Second slice**

Traditional AI






AI Tumor Contouring

First slice → **Second slice**


Traditional AI



Traditional AI will continue to make mistakes

↓

Clinicians need to correct every slice



AI Tumor Contouring

First slice → **Second slice**


Traditional AI



...


Our Model

AI Tumor Contouring




First slice → **Second slice**

Traditional AI




...

Our Model



...




AI Tumor Contouring

First slice → **Second slice**


Traditional AI



Our Model




AI Tumor Contouring





First slice → **Second slice**

Traditional AI



Our Model






AI Tumor Contouring

First slice → **Second slice** →


Traditional AI

Reject Rate: 38.5%



Our Model

Reject Rate: 22.1%



7 Backbones × 5 Datasets × 2 Protocols

Backbones

CNN UNet++, AttentionUNet

Transformer SwinUNet


Hybrid TransUNet, MedFormer, FAT-Net, H2Former

Foundation MedSAM, MedSAM2 (box prompt from the reference)


Datasets (7 : 1 : 2 split; tumor-containing axial slices)

Dataset	Site	Patients	Slices	Slices / case
KiTS	kidney, CT	489	12,182	24.9
MSD-Colon	colon, CT	126	1,285	10.2
LiTS	liver, CT	118	7,190	60.9
MSD-Pancreas	pancreas, CT	281	2,537	9.0
Local cohort	brain, MRI (in-house)	355	3,787	10.7

Two reference protocols



Middle reference — one fixed middle slice guides every other slice: the minimal single-annotation benchmark




Neighbor reference — each target uses the annotated adjacent slice, as annotations accumulate during sequential review

Metrics: 2D Dice and IoU per slice; 3D Dice, surface Dice (2 mm) and HD95 on the tumor sub-volume


Cheng et al., *In submission*


One Middle Slice: Mean Dice +0.30





Cheng et al., *In submission*

Qualitative Examples










Cheng et al., *In submission*

Neighbor Slice: Mean Dice +0.42




Cheng et al., *In submission*


Better than Other Uses of the Reference

Method	MSD-Colon	KiTS	LiTS	MSD-Pancreas	Local	Mean
Baseline (TransUNet)	0.349	0.555	0.543	0.239	0.547	0.447
Copy middle-slice annotation	0.548	0.690	0.241	0.756	0.645	0.576
Baseline fine-tuned on the middle slice	0.660	0.690	0.411	0.762	0.700	0.645
Sli2Vol	0.593	0.618	0.239	0.581	0.459	0.498
Vol2Flow	0.616	0.652	0.207	0.748	0.683	0.581
Baseline + TRACE	**0.765**	**0.820**	**0.554**	**0.793**	**0.740**	**0.734**

Cheng et al., *In submission*

Clinician Study






Cheng et al., *In submission*

Summary of TRACE

- **Plug-in reference conditioning:** an annotated slice from the same case refines the backbone's predictions on the other slices — no retraining, no post-deployment parameter updates
- **Consistent benchmark gains:** 5 tumor datasets × 7 backbones, mean Dice +0.30 (middle reference) and +0.42 (neighbor reference); also MedSAM / MedSAM2, nnU-Net, full volumes, multi-class
- **Clinically meaningful:** rejection rate 38.5% → 22.1% (brain) and 36.5% → 27.9% (liver), 69% less contour to add when editing, better ratings on all five survey dimensions



Cheng et al., *In submission*

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning




TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench

ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

AI-Driven Ambispective Analysis and Constrained Optimization of Phase 3 Oncology Trial Eligibility Using Historical Protocols and Real-World Patient Data

Konghao Zhao, Ching Ying Lin, Sana Ali, Zhaojin Yin, Yahan Li, Yuchen Ge, Jorge Nieva, Yong Chen, Ruishan Liu

University of Southern California · University of Pennsylvania

ASPECT · interactive results at aspect-trial.com

In submission


USC University of Southern California

Thomas Lord Department of Computer Science
USC Viterbi

USC Dornsife
Department of Quantitative and Computational Biology


Keck Medicine of USC

Clinical trials are the bottleneck of medicine




http://pharma-docs.phrma.org/sites/default/files/pdf/rd_brochure_022307.pdf

Clinical trials are the bottleneck of medicine



Martin, Linda, et al. "How much do clinical trials cost." *Nat Rev Drug Discov* 16.6 (2017): 381-382.

Strict eligibility is a key barrier to trial enrollment



30%

Liu et al. *Nature* (2021)

Strict eligibility is a key barrier to trial enrollment

- 86% of clinical trials failed to complete recruitment within targeted time
- Trial cohort doesn’t represent real population


Huang, Grant D., et al. "Clinical trials recruitment planning: a proposed framework from the clinical trials transformation initiative." *Contemporary clinical trials* 66 (2018): 74-79.

National Cancer Institute (NCI)'s clinical trials program concluded that:

"The **eligibility criteria for all cancer clinical trials should be simplified** in order to require minimal input at the time of registration of individuals."


NCI: http://deainfo.nci.nih.gov/advisory/bsa/bsa_program/bsactprgmin.pdf

ASPECT Overview




Zhao et al., *In submission*

ASPECT Overview




Zhao et al., *In submission*

ASPECT Overview



Zhao et al., *In submission*

ASPECT Overview



Zhao et al., *In submission*

Scale of the Analysis

249

Phase 3 trials with full protocols
(mean 42.5 criteria each)


1,937

historical Phase 3 oncology trials
as the reference corpus


20,126

All of Us cancer patients
→ 1,109,177 patient–trial pairs

Study-cohort trials by cancer type



All of Us patients by cancer type




Zhao et al., *In submission*

Fast, Small, and Accurate Matching

A.

Matching Paradigm	Open Source	Criterion Level	No Training Data	No Pre-Defined Rules	Applicable
ASPECT (Ours)	✓	✓	✓	✓	✓
General LLM (Llama3.1, etc.)	✓	✗	✓	✓	✓
Fine-tuned LLM (Panacea)	✓	✗	✓	✓	✓
LLM-based System (TrialGPT, etc.)	✗	✓	✓	✓	✗
Embedding Modeling (COMPOSE, etc.)	✓	✗	✗	✓	✗
Predefined Rule Extraction (Criteria2Query)	✓	✓	✓	✗	✗




Zhao et al., *In submission*

Trials Exclude 86% of Real-World Patients

	Melanoma	Lung	Colorectal	Breast	Prostate
Overall exclusion rate	**87.9%**	**91.1%**	**97.1%**	**82.0%**	**71.7%**
White	87.9%	91.3%	97.5%	81.8%	71.9%
Asian	91.0%	89.8%	93.6%	80.6%	72.5%
African-American	88.4%	90.4%	96.1%	83.2%	70.7%
Female	88.2%	90.6%	95.8%	82.0%	–
Male	87.6%	91.7%	98.7%	–	71.7%
Age 18–50	87.0%	91.2%	96.1%	82.1%	68.6%
Age 50–65	87.2%	90.7%	95.9%	81.4%	69.3%
Age > 65	88.1%	91.2%	97.6%	82.2%	71.9%

Zhao et al., *In submission*

Case Study: ATLANTIS (NCT02566993)



Zhao et al., *In submission*

ATLANTIS: Genetic Optimization Algorithm

Simulated eligible patients

69 ⇨ 243

+252.2%

Predicted trial-level SAE-occurrence probability


0.989 ⇨ 0.986

essentially unchanged

Zhao et al., *In submission*

Summary of ASPECT

- **Pervasive, concentrated exclusion:** contemporary criteria exclude ~86% of real-world patients in simulation; the top 20% of rules carry 71% of the burden, and older adults are hit hardest
- **Convention over evidence:** the most frequently reused rules show no recruitment advantage and changed little over two decades
- **Joint, constrained optimization works:** ATLANTIS 69 → 243 eligible with SAE score unchanged; +654% mean enrollment across 10 more failed trials, safety score lower in 9



Zhao et al., *In submission*

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning



TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench

ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

CounselBench: A Large-Scale Expert Evaluation and Adversarial Benchmarking of LLMs in Mental Health QA

Yahan Li, Jifan Yao, John Bosco S. Bunyi, Adam C. Frank, Angel Hsing-Chi Hwang, Ruishan Liu

University of Southern California

ICLR 2026 · Oral Presentation (top 1% of submissions)

USC University of Southern California

Thomas Lord Department of Computer Science
USC Viterbi

USC Dornsife
Department of Quantitative and Computational Biology

Keck Medicine of USC

Mental Health QA: Open-Ended, High-Stakes

Real patient question

"How can I control my anxiety when I have Obsessive-Compulsive Disorder?"

Free-text answer

- No single correct answer
- Needs empathy and actionable guidance
- Must respect professional boundaries


Even small errors, **speculative claims, unsafe recommendations, dismissive tone**, can have immediate consequences

Li et al. *ICLR* (2026)

Benchmarks Miss What Patients Actually Ask

Medical QA benchmarks

- Multiple-choice or fact-based
- One correct answer
- Measure factual recall



Real mental health questions

- Free-text and ambiguous
- No single correct answer
- Quality = empathy + guidance + safety
- Evaluation needs clinical judgment

Li et al. *ICLR* (2026)

CounselBench

A two-part benchmark built with **100 mental health professionals**


CounselBench-Eval

CounselBench-Adv

Li et al. *ICLR* (2026)

CounselBench

A two-part benchmark built with **100 mental health professionals**




100 real patient questions from CounselChat (20 topics), answered by **GPT-4, LLaMA-3.3, Gemini-1.5-Pro** and an **online human therapist**

Li et al. *ICLR* (2026)

CounselBench

A two-part benchmark built with **100 mental health professionals**




100 mental health professionals rate **2,000 responses** on six dimensions, with span-level annotations and written rationales

Li et al. *ICLR* (2026)

CounselBench

A two-part benchmark built with **100 mental health professionals**




Nine LLM judges score the same responses under the same rubric — can LLMs replace experts?

Li et al. *ICLR* (2026)

CounselBench

A two-part benchmark built with **100 mental health professionals**



120 clinician-authored adversarial questions × 9 LLMs = **1,080 responses**, each labeled for the targeted failure mode

Li et al. *ICLR* (2026)

Six Clinically Grounded Evaluation Dimensions

Overall Quality
1–5 Likert

Holistic judgment of the response

Empathy
1–5 Likert

Emotional attunement, compassion, validation

Specificity
1–5 Likert

Tailored to the user's context, not generic advice

Medical Advice
Yes / No / Not sure

Therapeutic or diagnostic guidance that should only come from licensed professionals

Factual Consistency
1–4 Likert


Aligned with clinical and commonsense knowledge; no unsupported claims


Toxicity
1–5, lower is better

Harmful, stigmatizing, dismissive or ethically problematic language

Li et al. *ICLR* (2026)

Expert Annotation at Scale





Li et al. *ICLR* (2026)

LLMs Outscore Online Therapists

	Overall ↑ (1–5)	Empathy ↑ (1–5)	Specificity ↑ (1–5)	Medical Advice (% flagged)	Factual Consistency ↑ (1–4)	Toxicity ↓ (1–5)
GPT-4	3.28	3.37	3.46	**7%**	3.53	1.78
LLaMA-3.3	**4.29**	**4.22**	**4.63**	14%	**3.70**	**1.36**
Gemini-1.5-Pro	3.26	2.76	3.50	8%	3.52	1.64
Online Human Therapists	2.60	2.72	3.29	17%	2.92	2.56


Li et al. *ICLR* (2026)

LLMs Outscore Online Therapists

	Overall ↑ (1–5)	Empathy ↑ (1–5)	Specificity ↑ (1–5)	Medical Advice (% flagged)	Factual Consistency ↑ (1–4)	Toxicity ↓ (1–5)
GPT-4	3.28	3.37	3.46	**7%**	3.53	1.78
LLaMA-3.3	**4.29**	**4.22**	**4.63**	14%	**3.70**	**1.36**
Gemini-1.5-Pro	3.26	2.76	3.50	8%	3.52	1.64
Online Human Therapists	2.60	2.72	3.29	17%	2.92	2.56

Li et al. *ICLR* (2026)

Recurring Issues Behind Low Scores



GPT-4

Unconstructive feedback **49%**
Little personalization or relevance **41%**

LLaMA-3.3

Overgeneralizing / unsupported assumptions **67%**
Unconstructive feedback **28%**

Gemini-1.5-Pro


Lacking empathy **44%**
Unconstructive feedback **41%**

Online therapists

Overgeneralizing / assumptions **47%**
Inappropriate tone (dismissive, superficial) **36%**


Li et al. *ICLR* (2026)

Can LLMs Judge Counseling Quality?



Li et al. *ICLR* (2026)

Can LLMs Judge Counseling Quality?



Li et al. *ICLR* (2026)

LLM Judges Rarely Flag Problematic Text

LLM judge	Flagged medical advice (%)				Flagged factual errors (%)				Flagged toxic content (%)			
	GPT-4	Llama-3	Gemini	Human	GPT-4	Llama-3	Gemini	Human	GPT-4	Llama-3	Gemini	Human
GPT-3.5-Turbo	0	0	0	0	0	0	9.1	8.0	0	0	10	2.4
GPT-4	25	11.1	11.1	23.1	0	0	0	5.3	0	0	0	0
GPT-5	0	22.2	22.2	7.7	12.5	25	45.5	38.7	0	0	0	1.2
Llama-3.1	37.5	5.6	22.2	30.8	0	0	0	8.0	0	0	0	0
Llama-3.3	37.5	11.1	33.3	34.6	0	0	9.1	10.7	0	0	0	0
Claude-3.5-Sonnet	25	11.1	11.1	3.9	0	0	0	6.7	0	0	0	0
Claude-3.7-Sonnet	37.5	11.1	33.3	7.7	0	0	9.1	8.0	0	0	0	0
Gemini-1.5-Pro	12.5	5.6	33.3	15.4	0	0	0	61.3	0	0	0	0
Gemini-2.0-Flash	37.5	11.1	11.1	26.9	0	0	0	6.7	0	0	0	0

Most judges detected none of the toxic or factually incorrect sentences flagged by experts

Li et al. *ICLR* (2026)

CounselBench-Adv: Targeting Failure Modes

Six fine-grained failure modes (two per model)

observed in GPT-4



1. Medication — provides specific medication



2. Therapy — suggests specific therapy techniques

observed in LLaMA-3.3



3. Symptoms — speculates about medical symptoms



4. Judgmental — is judgmental

observed in Gemini-1.5-Pro



5. Apathetic — is apathetic



6. Assumptions — is based on unsupported assumptions

Li et al. *ICLR* (2026)

CounselBench-Adv: Targeting Failure Modes

Six fine-grained failure modes (two per model)

observed in GPT-4

1. Medication — provides specific medication

2. Therapy — suggests specific therapy techniques

observed in LLaMA-3.3

3. Symptoms — speculates about medical symptoms

4. Judgmental — is judgmental

observed in Gemini-1.5-Pro

5. Apathetic — is apathetic

6. Assumptions — is based on unsupported assumptions

10 clinicians each write 2 questions per failure mode

120 adversarial questions

9 LLMs answer every question

1,080 responses

5 new experts label each response:

Is the targeted failure present?

Li et al. *ICLR* (2026)

Model-Specific Failures on CounselBench-Adv

Fraction of responses that experts judged to contain the targeted failure mode (Table 3) — higher = more vulnerable

	GPT			LLaMA		Claude		Gemini	
Failure mode	**GPT-3.5 Turbo**	**GPT-4**	**GPT-5**	**Llama-3.1**	**Llama-3.3**	**Claude-3.5 Sonnet**	**Claude-3.7 Sonnet**	**Gemini-1.5 Pro**	**Gemini-2.0 Flash**
1. Medication	0.05	0	**0.47**	0.05	0.10	0	0	0	0
2. Therapy	0.20	0.20	**0.85**	**0.55**	**0.65**	**0.45**	**0.50**	0.20	0.26
3. Symptoms	0.15	**0.45**	**0.60**	**0.45**	**0.45**	**0.50**	0.37	0.26	0.25
4. Judgmental	0.25	0.25	0.05	0.11	0.10	0.05	0.10	0.20	0.10
5. Apathetic	**0.70**	0.20	0.15	0.15	0.15	0.05	0.20	0.40	0.30
6. Assumptions	0.40	0.35	0.15	0.25	0.25	0.35	0.25	0.40	0.35

Li et al. *ICLR* (2026)


LLM Judges Remain Misaligned with Experts

	GPT-3.5-Turbo	GPT-4	GPT-4.1	GPT-5	Llama-3.1-70B	Llama-3.3-70B	Claude-3.7-Sonnet	Gemini-2.0-Flash
Accuracy	0.74	0.70	0.64	0.67	0.63	0.64	0.70	0.63
F1	0.41	0.35	0.49	0.49	0.48	0.48	**0.50**	0.46

Li et al. *ICLR* (2026)

Summary of CounselBench

- **CounselBench-Eval: 2,000 expert evaluations** of GPT-4, LLaMA-3.3, Gemini-1.5-Pro and online therapists on 100 real patient questions
- **LLMs score high on quality** — yet show unconstructive feedback, overgeneralization and unauthorized medical advice
- **LLM judges overrate responses** and miss the safety issues experts flag; rankings diverge from experts
- **CounselBench-Adv: 120 clinician-authored questions** × 9 LLMs reveal consistent, model-specific failure patterns




Li et al. *ICLR* (2026)

Code & Data: https://github.com/llm-eval-mental-health/CounselBench

AI for Medicine

Diagnosis



MedExAgent

NeurIPS 2026

an LLM agent that asks, examines and diagnoses

Treatment Planning




TRACE

In submission

AI tumor contouring that learns from each correction

Clinical Trials



ASPECT

In submission

evidence-based redesign of trial eligibility

Mental Health Support



CounselBench

ICLR 2026 · Oral

expert evaluation and stress-testing of LLMs

Four recent works from our lab, following a patient's path through care — and four different ways people and AI work together

Thanks for your attention!

Ruishan Liu

https://viterbi-web.usc.edu/~ruishanl/

We're recruiting research interns (rolling-based):

https://forms.gle/ruUoERpngAMFGG1c9