

# Simulation of Biological Neuronal Networks

## Cortical networks: Background and simple models

Jyotika Bahuguna, Abigail Morrison

[morrison@fz-juelich.de](mailto:morrison@fz-juelich.de)

Inst. of Neuroscience and Medicine (INM-6)  
Computational and Systems Neuroscience  
Research Center Jülich, Germany

# Outline

Biological background: Cortex structure

Some graph-theoretical concepts

Random networks

the balanced-random-network model (brunel, 2000)

Literature

# **Biological background: Cortex structure**

- ▶ **Macroscopic structure:**

- ▶ visible with your eyes
- ▶ thousand/millions of cells forming a structure
- ▶ e.g brain areas, long range connections

- ▶ **Microscopic structure:**

- ▶ magnifying instruments are needed to recognize structure
- ▶ depending on resolution: group of cells, single cell, molecules
- ▶ e.g small neuronal network, short range connections

# Microscopic structure: Cell types

- ▶ cortical tissue is mainly composed of two cell types:
  - ▶ **neuroglia (glia):**
    - ▶ important role in development of the brain
    - ▶ metabolic supportive role
    - ▶ control ionic composition of extracellular space
    - ▶ form myelin sheath around axons of neurons
    - ▶ do not take part in interaction between neurons on a millisecond scale  
(however, may play a role in slow modulations)
  - ▶ **nerve cells (neurons):**
    - ▶ carry out information processing and storage in the brain


# Microscopic structure: Neuron types

## ► pyramidal cell:

- ▶ dendrites covered with spines
- ▶ axon leaves cortex into white matter but has numerous branches close to cell body
- ▶ axons makes **excitatory** synapses
- ▶ cell receives input from inhibitory cells mainly at the soma and from excitatory cells at the basal and apical dendrite

## Microscopic structure: Neuron types

## Pyramidal cell




# Microscopic structure: Neuron types

- ▶ pyramidal cell:
  - ▶ dendrites covered with spines
  - ▶ axon leaves cortex into white matter but has numerous branches close to cell body
  - ▶ axons makes **excitatory** synapses
  - ▶ cell receives input from inhibitory cells mainly at the soma and from excitatory cells at the basal and apical dendrite
- ▶ spiny stellate cell:
  - ▶ axon branches out within the cortex (rarely leaves the cortex)
  - ▶ axons makes **excitatory** synapses
  - ▶ soma receives exclusively inhibitory and dendrites mostly excitatory inputs

# Microscopic structure: Neuron types

## Spiny stellate cell




# Microscopic structure: Neuron types


- ▶ pyramidal cell:
  - ▶ dendrites covered with spines
  - ▶ axon leaves cortex into white matter but has numerous branches close to cell body
  - ▶ axons makes **excitatory** synapses
  - ▶ cell receives input from inhibitory cells mainly at the soma and from excitatory cells at the basal and apical dendrite
- ▶ spiny stellate cell:
  - ▶ axon branches out within the cortex (rarely leaves the cortex)
  - ▶ axons makes **excitatory** synapses
  - ▶ soma receives exclusively inhibitory and dendrites mostly excitatory inputs
- ▶ smooth stellate cell (e.g. basket cell):
  - ▶ axon branches out only in the cortex
  - ▶ axons makes **inhibitory** synapses (GABA)
  - ▶ both the soma and the dendrites receive a mixture of excitatory and inhibitory inputs

# Microscopic structure: Neuron types

## Smooth stellate cell





# Microscopic structure: Neuron types



(Binzegger et al., 2004)


# Microscopic structure: Cortex layers

- subdivision of neocortex into 6 (or 5) layers
  - **layer I**: very few neurons, only dendrites and mesh of horizontal axons
  - **layer II/III**: small pyramidal cells
  - **layer IV**: small and medium-size pyramidal cells and stellate cells in layer IVA, almost exclusively stellate cells in IVB
  - **layer V**: all cell types, very large pyramidal cells dominate
  - **layer VI**: small and medium-size pyramidal cells



# Cortex connectivity: Summary - some numbers

Table 1.5.1. *Densities of neurons in the cortex (thousands per cubic millimeter)*




Animal	Density	A		B		C	
		Region	Density	Layer	Density	Layer	Density
Mouse	142.5	Visual	106	I	20		
Rat	105.0	Somatosensory	60	II	82		
Guinea pig	52.5	Auditory	43	III	62		
Rabbit	43.8	Motor	30	IV	67		
Cat	30.8			V	61		
Dog	24.5			VI	77		
Monkey	21.5						
Human	10.5						
Elephant	6.9						
Whale	6.8						

Table 1.5.4. *Typical compositions of cortical tissues*

Variable	Value
Neuronal density	40,000/mm <sup>3</sup>
Neuronal composition:	
Pyramidal	75%
Smooth stellate	15%
Spiny stellate	10%
Synaptic density	$8 \cdot 10^8/\text{mm}^3$
Axonal length density	$3,200 \text{ m/mm}^3$
Dendritic length density	$400 \text{ m/mm}^3$
Synapses per neuron	20,000
Inhibitory synapses per neuron	2,000
Excitatory synapses from remote sources per neuron	9,000
Excitatory synapses from local sources per neuron	9,000
Dendritic length per neuron	10 mm

# Cortex connectivity: Summary - some numbers

Table 1.5.1. Densities of neurons in the cortex (thousands per cubic millimeter)



A		B		C	
Animal	Density	Region	Density	Layer	Density
Mouse	142.5	Visual	106	I	20
Rat	105.0	Somatosensory	60	II	82
Guinea pig	52.5	Auditory	43	III	62
Rabbit	43.8	Motor	30	IV	67
Cat	30.8			V	61
Dog	24.5			VI	77
Monkey	21.5				
Human	10.5				
Elephant	6.9				
Whale	6.8				

Table 1.5.4. Typical compositions of cortical tissues

Variable	Value
Neuronal density	40,000/mm <sup>3</sup>
Neuronal composition:	
Pyramidal	75%
Smooth stellate	15%
Spiny stellate	10%
Synaptic density	$8 \cdot 10^{14}/mm^3$
Axonal length density	3,200 m/mm <sup>3</sup>
Dendritic length density	400 m/mm <sup>3</sup>
Synapses per neuron	20,000
Inhibitory synapses per neuron	2,000
Excitatory synapses from remote sources per neuron	9,000
Excitatory synapses from local sources per neuron	9,000
Dendritic length per neuron	10 mm

# Some graph-theoretical concepts


- ▶ Graph:  $G := (V, E)$

$V$ : set of 'vertices' (neurons)

$E$ : set of (ordered) pairs of vertices  $\curvearrowright$  'edges' (synapses)

example:

$$V = \{a, b, c\} \quad , \quad E = \{\{a, b\}, \{b, a\}, \{b, c\}, \{a, c\}\}$$



# Some graph-theoretical concepts

- ▶ Graph:  $G := (V, E)$

$V$ : set of 'vertices' (neurons)

$E$ : set of (ordered) pairs of vertices  $\curvearrowright$  'edges' (synapses)

example:


$$V = \{a, b, c\} \quad , \quad E = \{\{a, b\}, \{b, a\}, \{b, c\}, \{a, c\}\}$$

- ▶ more convenient:

presynaptic  
adjacency matrix  $A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$  postsynaptic

(columns=axons, rows=dendrites)

with  $A_{ij} = \begin{cases} 1 & \text{edge (synapse) } j \rightarrow i \text{ present} \\ 0 & \text{else} \end{cases}$



# Some graph-theoretical concepts

- Graph:  $G := (V, E)$

$V$ : set of 'vertices' (neurons)

$E$ : set of (ordered) pairs of vertices  $\curvearrowright$  'edges' (synapses)

example:

$$V = \{a, b, c\} \quad , \quad E = \{\{a, b\}, \{b, a\}, \{b, c\}, \{a, c\}\}$$

- more convenient:



presynaptic  
adjacency matrix  $A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$  postsynaptic

(columns=axons, rows=dendrites)

with  $A_{ij} = \begin{cases} 1 & \text{edge (synapse) } j \rightarrow i \text{ present} \\ 0 & \text{else} \end{cases}$

weight matrix  $W = \begin{pmatrix} 0 & J & 0 \\ J & 0 & 0 \\ J & -J & 0 \end{pmatrix}$

(for weighted graphs)



# Some graph-theoretical concepts

- **in-degree:**  $K_i^{\text{in}} = \sum_j A_{ij}$  (number of inputs)
- **out-degree:**  $K_j^{\text{out}} = \sum_i A_{ij}$  (number of targets)

# Some graph-theoretical concepts

- ▶ **in-degree:**  $K_i^{\text{in}} = \sum_j A_{ij}$  (number of inputs)
- ▶ **out-degree:**  $K_j^{\text{out}} = \sum_i A_{ij}$  (number of targets)
- ▶ note:
  - ▶  $A$  and  $W$  generally NOT symmetric

# Some graph-theoretical concepts

- ▶ **in-degree:**  $K_i^{\text{in}} = \sum_j A_{ij}$  (number of inputs)
- ▶ **out-degree:**  $K_j^{\text{out}} = \sum_i A_{ij}$  (number of targets)
- ▶ note:
  - ▶  $A$  and  $W$  generally NOT symmetric
  - ▶ 'space' not *explicitly* represented (depends on neuron labeling)

# Some graph-theoretical concepts

- ▶ in-degree:  $K_i^{\text{in}} = \sum_j A_{ij}$  (number of inputs)
- ▶ out-degree:  $K_j^{\text{out}} = \sum_i A_{ij}$  (number of targets)
- ▶ note:
  - ▶  $A$  and  $W$  generally NOT symmetric
  - ▶ 'space' not *explicitly* represented (depends on neuron labeling)
- ▶ structure of  $W$  determines features of network dynamics

# Some graph-theoretical concepts

- ▶ **in-degree:**  $K_i^{\text{in}} = \sum_j A_{ij}$  (number of inputs)
- ▶ **out-degree:**  $K_j^{\text{out}} = \sum_i A_{ij}$  (number of targets)
- ▶ note:
  - ▶  $A$  and  $W$  generally NOT symmetric
  - ▶ 'space' not *explicitly* represented (depends on neuron labeling)
- ▶ structure of  $W$  determines features of network dynamics
  - example: linear firing-rate network model (Wilson-Cowan model)

$$\tau \frac{dy}{dt} = -y + Wy$$

with firing rates  $y(t) = [y_1(t), \dots, y_N(t)]^T$  of neurons  $1, \dots, N$

↪ stability of fixed point determined by eigenvalue spectrum of  $W - 1$

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, . . . (reflects lack of knowledge)

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, . . . (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

Be precise when describing network structure!

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

Be precise when describing network structure!

- ▶ popular random networks in Computational Neuroscience:
  - ▶ Erdős-Renyi graphs:
    - ▶ Each possible connection is randomly and independently drawn with probability  $\epsilon$ .

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

Be precise when describing network structure!

- ▶ popular random networks in Computational Neuroscience:
  - ▶ Erdős-Renyi graphs:
    - ▶ Each possible connection is randomly and independently drawn with probability  $\epsilon$ .
  - ▶ random networks with fixed in- (or out-) degree:
    - ▶ random distribution of a fixed number of 1's in each row (column) of the adjacency matrix

# Random networks

- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

Be precise when describing network structure!

- ▶ popular random networks in Computational Neuroscience:
  - ▶ Erdős-Renyi graphs:
    - ▶ Each possible connection is randomly and independently drawn with probability  $\epsilon$ .
  - ▶ random networks with fixed in- (or out-) degree:
    - ▶ random distribution of a fixed number of 1's in each row (column) of the adjacency matrix
  - ▶ small-world networks:
    - ▶ (deterministic) nearest-neighbour connections
    - ▶ random redistribution of individual links with probability  $\alpha$

# Random networks


- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!

Be precise when describing network structure!

- ▶ popular random networks in Computational Neuroscience:
  - ▶ Erdős-Renyi graphs:
    - ▶ Each possible connection is randomly and independently drawn with probability  $\epsilon$ .
  - ▶ random networks with fixed in- (or out-) degree:
    - ▶ random distribution of a fixed number of 1's in each row (column) of the adjacency matrix
  - ▶ small-world networks:
    - ▶ (deterministic) nearest-neighbour connections
    - ▶ random redistribution of individual links with probability  $\alpha$
  - ▶ networks with specified high-order connectivity statistics (motifs)

# Random networks



Song et al. (2005), Highly Nonrandom Features  
of Synaptic Connectivity in Local Cortical Circuits

# Random networks


- ▶ connection matrix  $W$  often treated as *random* matrix with defined statistical properties, e.g. connection probability, in- or out-degree distribution, weight distribution, ... (reflects lack of knowledge)

There is no such thing as **THE** random-network model!


Be precise when describing network structure!

- ▶ popular random networks in Computational Neuroscience:
  - ▶ Erdős-Renyi graphs:
    - ▶ Each possible connection is randomly and independently drawn with probability  $\epsilon$ .
  - ▶ random networks with fixed in- (or out-) degree:
    - ▶ random distribution of a fixed number of 1's in each row (column) of the adjacency matrix
  - ▶ small-world networks:
    - ▶ (deterministic) nearest-neighbour connections
    - ▶ random redistribution of individual links with probability  $\alpha$
  - ▶ networks with specified high-order connectivity statistics (motifs)
  - ▶ ... and infinitely many more

# the balanced-random-network model (brunel, 2000)




Connection matrix  $J = \{J_{ij}\}$




- ▶  $N_E$  excitatory,  $N_I$  inhibitory neurons,  $N = N_E + N_I \sim 10^4 \dots 5$ ,  $N_E:N_I=4:1$
- ▶ random Dale-conform connectivity with fixed in-degrees  $K_{E/I} = N_{E/I}/10$
- ▶ free parameters:  $g$  (ratio of inhibitory to excitatory synaptic weight),  $\frac{\nu_{\text{ext}}}{\nu_{\text{thr}}}$  (ratio of external rate to threshold rate)

# the balanced-random-network model (brunel, 2000)



Connection matrix  $J = \{J_{ij}\}$



- ▶  $N_E$  excitatory,  $N_I$  inhibitory neurons,  $N = N_E + N_I \sim 10^4 \dots 5$ ,  $N_E:N_I=4:1$
- ▶ random Dale-conform connectivity with fixed in-degrees  $K_{E/I} = N_{E/I}/10$
- ▶ LIF dynamics with current synapses
- ▶ free parameters:  $g$  (ratio of inhibitory to excitatory synaptic weight),  $\frac{\nu_{\text{ext}}}{\nu_{\text{thr}}}$  (ratio of external rate to threshold rate)

# Activity regimes



- ▶ analysis of network dynamics in diffusion approximation (small synaptic weights, high firing rates, small correlations) (see Brunel, 2000)

# Activity regimes

- ▶ analysis of network dynamics in diffusion approximation (small synaptic weights, high firing rates, small correlations) (see Brunel, 2000)

## ↵ stationary and oscillatory states

SR: synchronous regular  
AI: asynchronous irregular  
SI: synchronous irregular (slow and fast)



# Activity regimes



- ▶ analysis of network dynamics in diffusion approximation (small synaptic weights, high firing rates, small correlations) (see Brunel, 2000)

- ↪ stationary and oscillatory states

SR: synchronous regular

AI: asynchronous irregular

SI: synchronous irregular (slow and fast)



# the balanced-random-network model


(brunel, 2000)

# the balanced-random-network model

(brunel, 2000)

Activity states

- ▶ Cortical neuron receives much higher proportion of excitatory synapses as compared to inhibitory synapses




# the balanced-random-network model

(brunel, 2000)

Activity states

- ▶ Cortical neuron receives much higher proportion of excitatory synapses as compared to inhibitory synapses
- ▶ In-vivo like activity
  - large membrane potential fluctuations
  - low firing rates
  - irregular spiking




# the balanced-random-network model

(brunel, 2000)

Activity states

- ▶ Cortical neuron receives much higher proportion of excitatory synapses as compared to inhibitory synapses
- ▶ In-vivo like activity
  - large membrane potential fluctuations
  - low firing rates
  - irregular spiking
- ▶ Possible explanation: cortex operates in inhibition dominated regime, i.e stronger synaptic strengths for inhibitory neurons




# the balanced-random-network model

(brunel, 2000)

## Chaos in random networks

- ▶ Cortical neuron receives much higher proportion of excitatory synapses as compared to inhibitory synapses
- ▶ In-vivo like activity
  - large membrane potential fluctuations
  - low firing rates
  - irregular spiking
- ▶ Possible explanation: cortex operates in inhibition dominated regime, i.e stronger synaptic strengths for inhibitory neurons
- ▶ chaotic spiking (high sensitivity)



Simulation of two identical networks ( $N \sim 10000$ ), slight perturbation of initial membrane potential of **one** neuron ( $\Delta V = 0.1$  mV)

# Literature: Cortex anatomy

- ▶ Abeles,  
*Corticonics: Neural Circuits of the Cerebral Cortex*,  
Cambridge University Press, 1991
- ▶ Braitenberg & Schüz,  
*Cortex: Statistics and Geometry of Neuronal Connectivity*,  
Springer, 2nd edition, 1998
- ▶ Thomson & Bannister (2003), Interlaminar Connections in the neocortex, *Cerebral Cortex* 13:5–14
- ▶ Binzegger et al. (2004), A Quantitative Map of the Circuit of Cat Primary Visual Cortex, *J Neurosci* 39(24):8441–8453
- ▶ Hellwig (2000), A quantitative analysis of the local connectivity between pyramidal neurons in layers 2/3 of the rat visual cortex, *Biological Cybernetics* 82:111–121
- ▶ Lund et al. (2003), Anatomical Substrates for Functional Columns in Macaque Monkey Primary Visual Cortex, *Cerebral Cortex* 13:15–24
- ▶ Voges et al. (2010), A modeler's view on the spatial structure of intrinsic horizontal connectivity in the neocortex, *Progress in Neurobiology* 92(3):277–292
- ▶ Song et al. (2005), Highly nonrandom features of synaptic connectivity in local cortical circuits, *PLoS Biology* 3(3):0507–0519

# Literature: Spiking random-network models

## Random matrices:

1. ML Mehta, *Random matrices*, 3rd edition, Elsevier Ltd., 2004

## Recurrent-network dynamics:

2. LF Abbott, C van Vreeswijk, Asynchronous states in a network of pulse-coupled oscillators, *Phys Rev E* 48:1483–1490, 1993
3. N Brunel, Dynamics of sparsely connected networks of excitatory and inhibitory spiking neurons, *J Comput Neurosci* 8:183–208, 2000
4. A Renart, N Brunel, XJ Wang, Mean-field theory of irregularly spiking neuronal populations and working memory in recurrent cortical networks, in *Computational neuroscience – A comprehensive approach*, J Feng (ed.), Chapman & Hall/CRC, London, 2003