

# Simulation of Biological Neuronal Networks

## Topological Networks

Robin Pauli, Abigail Morrison

{r.pauli|morrison}@fz-juelich.de



Inst. of Neuroscience and Medicine (INM-6)  
Computational and Systems Neuroscience  
Research Center Jülich, Germany

some time July

# Macroscopic structure


## ► forebrain (anterior part of the brain):

- ▶ diencephalon, e.g. thalamus and hypothalamus
- ▶ telencephalon (cerebrum)
  - ▶ basal ganglia (motor control, cognition, emotions, learning)
  - ▶ limbic system, e.g. amygdala (emotions)
  - ▶ olfactory bulb (smell)
  - ▶ cerebral cortex (archeocortex + neocortex)




# Macroscopic structure: Cortex folds

- ▶ neocortex = surface of the mammalian brain (gray matter)
- ▶ strongly folded in 'higher' mammals
  - ▶ large surface (relatively to skull volume)
  - ▶ efficient wiring between cortical areas through white matter




# Macroscopic structure: Cortex folds

- ▶ neocortex = surface of the mammalian brain (gray matter)
- ▶ strongly folded in 'higher' mammals
  - ▶ large surface (relatively to skull volume)
  - ▶ efficient wiring between cortical areas through white matter
- ▶ folds = 'sulci'
- ▶ region between adjacent folds = 'gyrus'
- ▶ location of (deep) sulci is consistent across individuals of the same species



# Macroscopic structure: Cerebral hemispheres

- ▶ two mirror-symmetric cerebral hemispheres
- ▶ interconnected via the Corpus callosum (white matter)




# Macroscopic structure: Cerebral hemispheres

- ▶ two mirror-symmetric cerebral hemispheres
- ▶ interconnected via the Corpus callosum (white matter)
- ▶ brain function is sometimes lateralised  
(i.e. preference for one hemisphere)
- ▶ in general, however, involvement of both hemispheres

e.g. visual processing:


each hemisphere receives information from both eyes

however, preference of input from left part of the visual field for right hemisphere  
(and vice versa)



# Cortical areas


- ▶ subdivision of cortex into areas according to
  - ▶ functional properties




© 2007 Encyclopædia Britannica, Inc.

# Cortical areas

- ▶ subdivision of cortex into areas according to
  - ▶ functional properties
  - ▶ histo-anatomical features (cytoarchitecture, e.g. Brodmann areas)




© 2007 Encyclopaedia Britannica, Inc.




# Cortical areas

- ▶ subdivision of cortex into areas according to
  - ▶ functional properties
  - ▶ histo-anatomical features (cytoarchitecture, e.g. Brodmann areas)
- ▶ frequent coincidence of both definitions  
e.g. primary visual cortex (V1)  
= Brodmann area 17





© 2007 Encyclopaedia Britannica, Inc.




# Cortical areas

- ▶ subdivision of cortex into areas according to
  - ▶ functional properties
  - ▶ histo-anatomical features (cytoarchitecture, e.g. Brodmann areas)
- ▶ frequent coincidence of both definitions  
e.g. primary visual cortex (V1)  
= Brodmann area 17
- ▶ further segmentation of cortical areas into functional subdomains



# Cortical areas

- ▶ subdivision of cortex into areas according to
  - ▶ functional properties
  - ▶ histo-anatomical features (cytoarchitecture, e.g. Brodman areas)
- ▶ frequent coincidence of both definitions  
e.g. primary visual cortex (V1)  
= Brodman area 17
- ▶ further segmentation of cortical areas into functional subdomains





# Cortical areas

- ▶ subdivision of cortex into areas according to
  - ▶ functional properties
  - ▶ histo-anatomical features (cytoarchitecture, e.g. Brodmann areas)
- ▶ frequent coincidence of both definitions
  - e.g. primary visual cortex (V1)  
= Brodmann area 17
- ▶ further segmentation of cortical areas into functional subdomains
  - e.g. rat 'barrel' cortex


# Macroscopic structure: Cortex layers

- subdivision of neocortex into 6 (or 5) layers
  - **layer I**: very few neurons, only dendrites and mesh of horizontal axons
  - **layer II/III**: small pyramidal cells
  - **layer IV**: small and medium-size pyramidal cells and stellate cells in layer IVA, almost exclusively stellate cells in IVB
  - **layer V**: all cell types, very large pyramidal cells dominate
  - **layer VI**: small and medium-size pyramidal cells




# Cortex connectivity: Afferent connections

- ▶ external (non-local) input from
  - ▶ thalamus  
(thalamo-cortical afferents)  
mainly to layer IV
  - ▶ other cortical areas  
(cortico-cortical afferents)  
via white matter  
mostly to superficial layers
- ▶ input from cortical neurons  
in the local vicinity



# Cortex connectivity: Vertical connectivity

- layer- and cell-type specific local connections





ChMap/ThalCtxInputs3.cdr

(Thomson et al., 2003)

# Cortex connectivity: Horizontal connectivity


- ▶ local synapses established by local axon collaterals arborizing within ~ 0.5 mm (all neuron types)
- ▶ intrinsic horizontal long-range connections of pyramidal cells over distances up to several millimeter (within gray matter)
- ▶ extrinsic long-range connections of pyramidal cells through white matter



(Voges et al., 2007)


# Cortex connectivity: Local connectivity

- ▶ probability of synaptic connection between adjacent cortical neurons decays to zero within a horizontal distance of  $\sim 0.5$  mm



# Cortex connectivity: Long-range connectivity


- intrinsic long-range connections form 'patchy' projection patterns, i.e. pyramidal cells project to distant clusters of target cells




(Lund et al., 2003)

# Random is not enough (II)


- ▶ Highly non-random distribution of connectivity motifs



[?]









## Dale versus hybrid connectivity



[?]


# Dale versus hybrid connectivity



[?]

- ▶ Same mean firing rate
- ▶ Hybrid less synchronous than Dale



# Distance dependent connectivity




[?]

- ▶ Probability of rewiring  $p_r$
- ▶  $0 < p_r < 1 \rightarrow$  small-world network ([?])

# Distance dependent connectivity




[?]



- ▶ Clustered networks are locally more synchronous.





# Distance dependent connectivity



pre-synaptic neurons  
▲ exc. neuron  
△ inh. neuron

post-synaptic exc. targets  
✖ exc. neuron  
● inh. neuron


post-synaptic inh. targets  
✖ exc. neuron  
● inh. neuron




Voges et al. 2012

# Distance dependent connectivity



## Local transitions




## Overlapping waves



## Melted stripes







## AI with SR



Voges et al. 2012

# Stimulus-induced variability reduction

Connectivity has impact on dynamics,  
implications for function of networks?



[?]

## Summary

- ▶ Random network models can capture many features of cortical activity but not all.
- ▶ The type of structured connectivity has a strong impact on the level of synchrony in the network.
- ▶ Clustered network can generate metastable dynamics that are highly variable.
- ▶ The stimulation of a cluster in a clustered network can reproduce the stimulus-induced drop in firing rate variability.

# Bibliography (I)

-  Pernice, V and Staude, B and Cardanobile, S and Rotter, S  
How structure determines correlations in neuronal networks  
*PLoS Comput Biol*, 7(5), 2011
-  Kriener, B and Tetzlaff, T and Aertsen, A and Diesmann, M and Rotter, S  
Correlations and population dynamics in cortical networks pathways and memory networks  
*Neural Comput*, 20(9):2185-2226, 2008
-  Kriener, B and Helias, M and Aertsen, A and Rotter, S  
Correlations in spiking neuronal networks with distance dependent connections  
*J Comput Neurosci*, 27(2):177-200, 2009

## Bibliography (II)



Watts, D J and Strogatz, S H

Collective dynamics of small-world networks

*Nature*, 393(6684):440-442, 1998



Litwin-Kumar, A and Doiron, B

Slow dynamics and high variability in balanced cortical networks with clustered connections

*Nature Neurosci*, 15(11):1498-1505, 2012



Doiron, B and Litwin-Kumar, A

Balanced neural architecture and the idling brain

*Front Comput Neurosci*, 8. 2014

## Bibliography (III)

-  **Hennequin, G and Vogels, T P and Gerstner, W**  
Optimal control of transient dynamics in balanced networks  
supports generation of complex movements  
*Neuron*, 82(6):1394-1406, 2014
-  **Churchland, M M and Cunningham, J P, ... and Shenoy, K V**  
Neural population dynamics during reaching  
*Nature*, 2012
-  **Churchland, M M, Byron, M Y, ... and Shenoy, K V**  
Stimulus onset quenches neural variability: a widespread cortical phenomenon  
*Nature Neurosci*, 13(3):369-378, 2010

## Bibliography (IV)



Roxin, A

The role of degree distribution in shaping the dynamics in networks of sparsely connected spiking neurons

*Front Comput Neurosci*, 5, 2011



Boucsein, C and Nawrot, M and Schnepel, P and Aertsen, A

Beyond the cortical column: abundance and physiology of horizontal connections imply a strong role for inputs from the surround

*Front Neurosci*, 5(322), 2011



Song, S and Sjöström, P J and Reigl, M and Nelson, S and Chklovskii, D B

Highly nonrandom features of synaptic connectivity in local cortical circuits

*PLoS Biol*, 3(3), e68, 2005