

Fermi–Amaldi-Based Local Mixing Function

Ivan P. Bosko and Viktor N. Staroverov

Department of Chemistry, The University of Western Ontario, London, Ontario N6A 5B7, Canada

(Dated: March 12, 2026)

I. MAIN TEXT

Consider the exact exchange energy density (closed-shell):

$$e_x^{\text{exact}}(\mathbf{r}) = -\frac{1}{4} \int \frac{|\gamma(\mathbf{r}; \mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}', \quad (1)$$

where

$$\gamma(\mathbf{r}, \mathbf{r}') = 2 \sum_{i=1}^{N/2} \varphi_i^*(\mathbf{r}) \varphi_i(\mathbf{r}') \quad (2)$$

is one-electron reduced density matrix [1, 2], and the Fermi–Amaldi [3, 4] energy density expressed via electron density, $\rho(\mathbf{r}) \geq 0$, and electron number, N ,

$$e_x^{\text{FA}}(\mathbf{r}) = -\frac{1}{2N} \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}'. \quad (3)$$

For one-orbital systems (one-electron and two-electron singlet), two quantities are equal for all values of $\mathbf{r}$ [5–7]:

$$e_x^{\text{exact}}(\mathbf{r}) = e_x^{\text{FA}}(\mathbf{r}) \quad N \leq 2 \quad \text{for all } \mathbf{r}. \quad (4)$$

For systems with more than one molecular orbital, the spatial behavior of both quantities needs to be studied closer to establish a relation similar to Eq. (4). Which is convenient to do analysing the corresponding exchange holes. The exchange hole is defined via

$$e_x(\mathbf{r}) = \frac{1}{2} \int \frac{\rho(\mathbf{r})\rho_x(\mathbf{r}, \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' \quad (5)$$

leading to expressions for the exact and the Fermi–Amaldi exchange holes, respectively:

$$\rho_x^{\text{exact}}(\mathbf{r}, \mathbf{r}') = -\frac{1}{2} \frac{|\gamma(\mathbf{r}, \mathbf{r}')|^2}{\rho(\mathbf{r})}, \quad (6)$$

$$\rho_x^{\text{FA}}(\mathbf{r}') = -\frac{1}{N} \rho(\mathbf{r}'). \quad (7)$$

Both quantities are strictly non-positive, hence, we will consider their non-negative absolute values, $|\rho_x^{\text{exact}}(\mathbf{r})|$ and $|\rho_x^{\text{FA}}(\mathbf{r})|$, for convenience.

Effectively, comparing equations (6) and (7), we are looking for zeros of the equation:

$$N|\gamma(\mathbf{r}, \mathbf{r}')|^2 - 2\rho(\mathbf{r})\rho(\mathbf{r}') = 0, \quad N > 2, \quad (8)$$

where, by Cauchy–Schwarz inequality:

$$\rho(\mathbf{r})\rho(\mathbf{r}') \geq |\gamma(\mathbf{r}, \mathbf{r}')|^2, \quad N > 2, \quad \text{for all } \mathbf{r}. \quad (9)$$

II. SCRAP

A local hybrid exchange functional is given by

$$E_x^{\text{LH}} = \int \{g(\mathbf{r})e_x^{\text{exact}}(\mathbf{r}) + [1 - g(\mathbf{r})]e_x^{\text{SL}}(\mathbf{r})\} d\mathbf{r}, \quad (10)$$

where the integrand is a mixture of two exchange density energies weighed by a local mixing function (LMF), $g(\mathbf{r})$, which is position-dependent. The e_x^{SL} and e_x^{exact} quantities are a semi-local and the exact exchange energy densities, respectively.

A particular case of a one-electron limit, i.e., the asymptotic limit ($r \rightarrow \infty$) where the exact exchange must cancel out an excessive self-repulsion of the Coulomb potential, is also satisfied:

$$\lim_{r \rightarrow \infty} \frac{e_x^{\text{FA}}}{e_x^{\text{exact}}} = 1 \quad (11)$$

since both energy densities approach $-1/2r$ as $r \rightarrow \infty$.

-
- [1] V. N. Staroverov, Density-functional approximations for exchange and correlation, in *A Matter of Density* (John Wiley & Sons, Ltd, 2012) Chap. 6, pp. 125–156.
 [2] R. Parr and Y. Weitao, *Density-Functional Theory of Atoms and Molecules*, International Series of Monographs on Chemistry (Oxford University Press, 1989).

- [3] E. Fermi and E. Amaldi, Le orbite ∞ degli elementi, *Mem. Reale Accad. Italia* **6**, 117 (1934).
 [4] P. Bénilan, J. A. Goldstein, and G. R. Rieder, The fermi-amaldi correction in spin polarized thomas-fermi theory, in *Differential Equations and Mathematical Physics*, Mathematics in Science and Engineering, Vol. 186 (Elsevier,

- 1992) pp. 25–37.
- [5] I. P. Bosko and V. N. Staroverov, Derivation and reinterpretation of the fermi–amaldi functional, *J. Chem. Phys.* **159**, 131101 (2023).
 - [6] I. P. Bosko and V. N. Staroverov, Exchange energies and density functionals for systems of fermions of arbitrary spin, *Phys. Rev. A* **108**, 052803 (2023).
 - [7] P. W. Ayers, R. C. Morrison, and R. G. Parr, Fermi-amaldi model for exchange-correlation: atomic excitation energies from orbital energy differences, *Mol. Phys.* **103**, 2061 (2005).
 - [8] J. Jaramillo, G. E. Scuseria, and M. Ernzerhof, Local hybrid functionals, *The Journal of chemical physics* **118**, 1068 (2003).
 - [9] T. M. Maier, A. V. Arbuznikov, and M. Kaupp, Local hybrid functionals: Theory, implementation, and performance of an emerging new tool in quantum chemistry and beyond, *Wiley Interdisciplinary Reviews: Computational Molecular Science* **9**, e1378 (2019).