

Fermi–Amaldi-Based Local Mixing Function

Ivan P. Bosko and Viktor N. Staroverov

Department of Chemistry, The University of Western Ontario, London, Ontario N6A 5B7, Canada

(Dated: March 17, 2026)

I. MAIN TEXT

Consider the exact exchange energy density (closed-shell):

$$e_x^{\text{exact}}(\mathbf{r}) = -\frac{1}{4} \int \frac{|\gamma(\mathbf{r}; \mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}', \quad (1)$$

where

$$\gamma(\mathbf{r}, \mathbf{r}') = 2 \sum_{i=1}^{N/2} \varphi_i^*(\mathbf{r}) \varphi_i(\mathbf{r}') \quad (2)$$

is the one-electron reduced density matrix [1, 2], and the Fermi–Amaldi [3, 4] energy density expressed via electron density, $\rho(\mathbf{r}) \geq 0$, and electron number, N ,

$$e_x^{\text{FA}}(\mathbf{r}) = -\frac{1}{2N} \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}'. \quad (3)$$

For one-orbital systems (one-electron and two-electron singlet), the two quantities are equal for all values of $\mathbf{r}$ [5–7]:

$$e_x^{\text{exact}}(\mathbf{r}) = e_x^{\text{FA}}(\mathbf{r}) \quad N \leq 2 \quad \text{for all } \mathbf{r}. \quad (4)$$

For systems with more than one molecular orbital, the spatial behavior of both quantities needs to be studied closer to establish a relationship between the exact and

the Fermi–Amaldi exchange energy densities. By the Cauchy–Schwarz inequality:

$$\rho(\mathbf{r})\rho(\mathbf{r}') \geq |\gamma(\mathbf{r}, \mathbf{r}')|^2 \quad N > 2 \quad \text{for all } \mathbf{r}, \mathbf{r}'. \quad (5)$$

However, the Fermi–Amaldi global prefactor, $1/(2N)$, is strictly smaller than the exact exchange prefactor, $1/4$, when $N > 2$. Because these opposing conditions compete, neither energy density integrand strictly dominates the other everywhere in space.

As $|\mathbf{r}| \rightarrow \infty$, the denominator can be approximated as $|\mathbf{r} - \mathbf{r}'| \approx |\mathbf{r}|$, revealing that both exchange energy densities approach the same asymptotic limit:

$$e_x^{\text{exact}}(\mathbf{r}) \sim e_x^{\text{FA}}(\mathbf{r}) \sim -\frac{\rho(\mathbf{r})}{2|\mathbf{r}|}. \quad (6)$$

Thus, in the asymptotic limit:

$$\lim_{|\mathbf{r}| \rightarrow \infty} \frac{e_x^{\text{FA}}}{e_x^{\text{exact}}} = 1. \quad (7)$$

Conversely, in the limit of infinitesimally small inter-electronic distances ($\mathbf{r}' \rightarrow \mathbf{r}$), the spatial parts of both integrands approach equality:

$$|\gamma(\mathbf{r}, \mathbf{r}')|^2 \rightarrow \rho(\mathbf{r})\rho(\mathbf{r}') \quad \text{as } \mathbf{r}' \rightarrow \mathbf{r}. \quad (8)$$

However, because the exact exchange prefactor, $1/4$, is larger than the Fermi–Amaldi prefactor, $1/(2N)$, for $N > 2$, the region where $|\mathbf{r} - \mathbf{r}'| \rightarrow 0$ is dominated by the exact exchange energy density.

-
- [1] V. N. Staroverov, Density-functional approximations for exchange and correlation, in *A Matter of Density* (John Wiley & Sons, Ltd, 2012) Chap. 6, pp. 125–156.
 - [2] R. Parr and Y. Weitao, *Density-Functional Theory of Atoms and Molecules*, International Series of Monographs on Chemistry (Oxford University Press, 1989).
 - [3] E. Fermi and E. Amaldi, Le orbite ∞ s degli elementi, *Mem. Reale Accad. Italia* **6**, 117 (1934).
 - [4] P. Bénilan, J. A. Goldstein, and G. R. Rieder, The fermi-amaldi correction in spin polarized thomas-fermi theory, in *Differential Equations and Mathematical Physics*, Mathematics in Science and Engineering, Vol. 186 (Elsevier,

- 1992) pp. 25–37.
- [5] I. P. Bosko and V. N. Staroverov, Derivation and reinterpretation of the fermi–amaldi functional, *J. Chem. Phys.* **159**, 131101 (2023).
- [6] I. P. Bosko and V. N. Staroverov, Exchange energies and density functionals for systems of fermions of arbitrary spin, *Phys. Rev. A* **108**, 052803 (2023).
- [7] P. W. Ayers, R. C. Morrison, and R. G. Parr, Fermi-amaldi model for exchange-correlation: atomic excitation energies from orbital energy differences, *Mol. Phys.* **103**, 2061 (2005).