

# Fermi–Amaldi-Based Local Mixing Function

Ivan P. Bosko and Viktor N. Staroverov

*Department of Chemistry, The University of Western Ontario, London, Ontario N6A 5B7, Canada*

(Dated: March 17, 2026)

## I. MAIN TEXT

Consider the exact exchange energy density (closed-shell):

$$e_x^{\text{exact}}(\mathbf{r}) = -\frac{1}{4} \int \frac{|\gamma(\mathbf{r}; \mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}', \quad (1)$$

where

$$\gamma(\mathbf{r}, \mathbf{r}') = 2 \sum_{i=1}^{N/2} \varphi_i^*(\mathbf{r}) \varphi_i(\mathbf{r}') \quad (2)$$

is one-electron reduced density matrix [1, 2], and the Fermi–Amaldi [3, 4] energy density expressed via electron density,  $\rho(\mathbf{r}) \geq 0$ , and electron number,  $N$ ,

$$e_x^{\text{FA}}(\mathbf{r}) = -\frac{1}{2N} \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}'. \quad (3)$$

For one-orbital systems (one-electron and two-electron singlet), two quantities are equal for all values of  $\mathbf{r}$  [5–7]:

$$e_x^{\text{exact}}(\mathbf{r}) = e_x^{\text{FA}}(\mathbf{r}) \quad N \leq 2 \quad \text{for all } \mathbf{r}. \quad (4)$$

For systems with more than one molecular orbital, the spatial behavior of both quantities needs to be studied closer to establish a relation similar to Eq. (4). This is conveniently done by analyzing the corresponding exchange holes.

The exchange hole,  $\rho_x(\mathbf{r}, \mathbf{r}')$ , is defined via

$$e_x(\mathbf{r}) = \frac{1}{2} \int \frac{\rho(\mathbf{r})\rho_x(\mathbf{r}, \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' \quad (5)$$

leading to expressions for the exact and the Fermi–Amaldi exchange holes, respectively:

$$\rho_x^{\text{exact}}(\mathbf{r}, \mathbf{r}') = -\frac{1}{2} \frac{|\gamma(\mathbf{r}, \mathbf{r}')|^2}{\rho(\mathbf{r})}, \quad (6)$$

$$\rho_x^{\text{FA}}(\mathbf{r}') = -\frac{1}{N} \rho(\mathbf{r}'). \quad (7)$$

Both exchange holes are strictly non-positive, and both are integrating to  $-1$  over all space.

When  $N > 2$ , we cannot establish an unambiguous inequality between these exchange holes for all  $\mathbf{r}$ . Although the Cauchy–Schwarz inequality states that

$$\rho(\mathbf{r})\rho(\mathbf{r}') \geq |\gamma(\mathbf{r}, \mathbf{r}')|^2 \quad N > 2 \quad \text{for all } \mathbf{r}, \quad (8)$$

the Fermi–Amaldi prefactor,  $1/N$ , decays faster than the constant prefactor,  $1/2$ , of the exact exchange hole.

Therefore, there are some ranges of  $|\mathbf{r} - \mathbf{r}'|$  dominated by one hole and others dominated by another.

As such, in the limit of infinitesimally small interelectronic distances ( $|\mathbf{r}| \rightarrow |\mathbf{r}'|$ ) we have:

$$|\gamma(\mathbf{r}, \mathbf{r})|^2 = \rho(\mathbf{r})^2, \quad (9)$$

consequently, the depths of the exchange holes are

$$\rho_x^{\text{exact}}(\mathbf{r}, \mathbf{r}) = -\frac{\rho(\mathbf{r})}{2} \quad (10)$$

and

$$\rho_x^{\text{FA}}(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{N} \quad (11)$$

thus, for  $N > 2$ , the well of the exchange hole is dominated by the exact exchange hole.

Regarding the behavior far from the reference electron, the asymptotic decay of the Fermi–Amaldi hole follows that of the total electron density at large distances:

$$\rho_x^{\text{FA}}(\mathbf{r}') \sim -\frac{1}{N} \exp\left(-2\sqrt{2I_{\min}}|\mathbf{r}'|\right), \quad (12)$$

where  $I_{\min}$  is the exact first ionization energy [2, 8]. For the exact exchange hole, the asymptotic behavior is given by:

$$\rho_x^{\text{exact}}(\mathbf{r}, \mathbf{r}') \sim -\frac{2|\varphi_{\text{HOMO}}(\mathbf{r})|^2}{\rho(\mathbf{r})} \exp\left(-2\sqrt{2I_{\min}}|\mathbf{r}'|\right), \quad (13)$$

given that of the molecular orbital [2, 8].

## II. SCRAP

A local hybrid exchange functional is given by

$$E_x^{\text{LH}} = \int \{g(\mathbf{r})e_x^{\text{exact}}(\mathbf{r}) + [1 - g(\mathbf{r})]e_x^{\text{SL}}(\mathbf{r})\} d\mathbf{r}, \quad (14)$$

where the integrand is a mixture of two exchange density energies weighed by a local mixing function (LMF),  $g(\mathbf{r})$ , which is position-dependent. The  $e_x^{\text{SL}}$  and  $e_x^{\text{exact}}$  quantities are a semi-local and the exact exchange energy densities, respectively.

A particular case of a one-electron limit, i.e., the asymptotic limit ( $r \rightarrow \infty$ ) where the exact exchange must cancel out an excessive self-repulsion of the Coulomb potential, is also satisfied:

$$\lim_{r \rightarrow \infty} \frac{e_x^{\text{FA}}}{e_x^{\text{exact}}} = 1 \quad (15)$$

since both energy densities approach  $-1/2r$  as  $r \rightarrow \infty$ .

- 
- [1] V. N. Staroverov, Density-functional approximations for exchange and correlation, in *A Matter of Density* (John Wiley & Sons, Ltd, 2012) Chap. 6, pp. 125–156.
  - [2] R. Parr and Y. Weitao, *Density-Functional Theory of Atoms and Molecules*, International Series of Monographs on Chemistry (Oxford University Press, 1989).
  - [3] E. Fermi and E. Amaldi, Le orbite  $\infty$ s degli elementi, *Mem. Reale Accad. Italia* **6**, 117 (1934).
  - [4] P. Bénilan, J. A. Goldstein, and G. R. Rieder, The fermi-amaldi correction in spin polarized thomas-fermi theory, in *Differential Equations and Mathematical Physics*, Mathematics in Science and Engineering, Vol. 186 (Elsevier, 1992) pp. 25–37.
  - [5] I. P. Bosko and V. N. Staroverov, Derivation and reinterpretation of the fermi–amaldi functional, *J. Chem. Phys.* **159**, 131101 (2023).
  - [6] I. P. Bosko and V. N. Staroverov, Exchange energies and density functionals for systems of fermions of arbitrary spin, *Phys. Rev. A* **108**, 052803 (2023).
  - [7] P. W. Ayers, R. C. Morrison, and R. G. Parr, Fermi-amaldi model for exchange-correlation: atomic excitation energies from orbital energy differences, *Mol. Phys.* **103**, 2061 (2005).
  - [8] M. M. Morrell, R. G. Parr, and M. Levy, Calculation of ionization potentials from density matrices and natural functions, and the long-range behavior of natural orbitals and electron density, *The Journal of Chemical Physics* **62**, 549 (1975).