

METIS documentation for output data structure

Introduction:

This document describes the content of output data structure (named "post") of METIS code. The first part describes the list of substructures in data post structure. The second part describes the output data model of METIS code, made of the list of time dependent scalar data with their description, followed by the list of time dependent profile data with their description.

Access to output data is not available in the standalone-compiled version of METIS.

The syntax to access to structure field in Matlab is
"post.<substructure_name>.<field_name>".

Example: "post.zerod.ip".

List of substructures of METIS output structure:

.

1- profil0d:

Time dependent profile data.

2- simout:

Data use by Simulink during run of Simulink workflow including METIS block named "simmetis". This substructure is empty when METIS is used without Simulink.

3- z0dinput:

Copy of METIS input data used during the computation of the simulation.

4- zerod:

Time dependent scalar data.

1. Substructure: zerod

Time dependent scalar data.

1.1. RR

plasma resistor (ohm).

1.2. aitb

effect of itb on temperature peaking factor.

1.3. ane

exponent of electron density profile.

1.4. ane_actual

actual exponent of electron density profile obtained taking into account the reference and constraints from transport.

1.5. ape

exponent of electron pressure profile.

1.6. asser

if 1, feedback on vloop is on.

1.7. ate

exponent of electron temperature profile.

1.8. betan

normalized total beta of the plasma.

1.9. betap

poloidal normalized pressure of the plasma (thermal, ITER definition : betap_th(1)).

1.10. betaptot

poloidal normalized pressure of the plasma (total, ITER definition : betap(1)).

1.11. d0

Shafranov shift (m).

1.12. diboot

relative error on lboot, at the end of convergence.

1.13. difcurconv

current diffusion solver state.

1.14. dini

relative error on lni, at the end of convergence.

1.15. disrupt

if = 1, radiative disruption flag.

1.16. dlh

width of the power deposition of LH (su).

1.17. dpfus

relative error on pfus, at the end of convergence.

1.18. drmdt

$d \sqrt{\Phi/\pi/B_0} / dt$ of LCMS(m/s) .

1.19. dsol

characteristic SOL width (λ_q , m).

1.20. dw

relative error on w, at the end of convergence.

1.21. dwbpdt

time derivative of plasma poloidal field energy (W).

1.22. dwdt

time derivative of total plasma energy (W).

1.23. dwmagtordt

time derivative of toroidal magnetic plasma energy (W).

1.24. dwthdt

time derivative of thermal plasma energy (W).

1.25. ecrit_he

alpha critical energy (eV): for He3 if option.gaz == 5, otherwise for He4.

1.26. ecrit_he3_DDn

critical energy (eV) of helium-3 fusion particules from D-D reactions.

1.27. ecrit_he4_DHe3

critical energy (eV) of helium-4 fusion particules from D-He3 reactions.

1.28. ecrit_he4_DT

critical energy (eV) of helium-4 fusion particules from D-T reactions.

1.29. ecrit_icrh

critical energy of ICRH accelerated fast ions (eV).

1.30. ecrit_nbi

critical energy of NBI beam (eV).

1.31. ecrit_p_DDp

critical energy (eV) of proton fusion particules from D-D reactions.

1.32. ecrit_p_DHe3

critical energy (eV) of proton fusion particules from D-He3 reactions.

1.33. ecrit_t_DDp

critical energy (eV) of tritium fusion particules from D-D reactions.

1.34. eddy_current

Eddy current in passive structure for breakdown description (A).

1.35. edgeflux

poloidal flux at the edge of plasma (V s).

1.36. efficiency

Fisch like LH current drive efficiency for LHCD(A/W/m²).

1.37. einj_icrh

mean energy of fast ion produce by ICRH in minority scheme (eV).

1.38. einj_lh

mean LH electron energy, only define if rip = 1 (eV).

1.39. einj_nbi_icrh

beam energy including ICRH effects (eV; reserved for internal used only).

1.40. esup_fus

D-T fusion fast alpha suprathermal energy (J).

1.41. esup_icrh

ICRH fast ions suprathermal energy (J).

1.42. esup_lh

LH fast ions suprathermal energy (J).

1.43. esup_nbi

NBI fast ions suprathermal energy (J).

1.44. etalh

LH current drive efficiency (A W⁻¹ m⁻²).

1.45. etalh0

LH current drive efficiency @ vloop = 0 (A W⁻¹ m⁻²).

1.46. etalh1

LH current drive efficiency correction due to hot conductivity (A W⁻¹ m⁻²).

1.47. firstorb_nbi

fraction of NBI power lost due to fast ion first orbit loss.

1.48. flux_edge_cor

Poloidal flux modification of reference cons.flux due to eddy current ($\text{Wb} / (2 \cdot \pi)$).

1.49. frac_pellet

fraction of fuelling due to pellet.

1.50. fracmino

minority ions fraction accelerated by ICRH.

1.51. frloss_icrh

part of icrh power lost due to fast ions losses.

1.52. frnbi

fraction of NBI power absorded in the plasma.

1.53. fwcorr

internal data to compute tem.

1.54. harm

ICRH minority scheme harmonic.

1.55. hitb

H factor of ITB.

1.56. hmhd

H factor limitation due to MHD (BetaN limit).

1.57. ialign

current drive alignment quality parameter (1 = good , 0 = bad).

1.58. iboot

bootstrap current (A).

1.59. *icd*

total current drive (A).

1.60. *ieccd*

ECCD current drive(A).

1.61. *ifus*

fast alpha (fusion) "bootstrap" current (A) .

1.62. *ifwcd*

FWCD current drive(A).

1.63. *ilh*

LH current drive(A).

1.64. *inbicd*

NBI current drive (A).

1.65. *indice_inv*

sawtooth invert radius indice.

1.66. *ini*

total non inductive current (A).

1.67. *iohm*

ohmic current (A).

1.68. *ip*

plasma current (A).

1.69. *ipar*

plasma current // B (A).

1.70. *irun*

runaway current (A).

1.71. j0fus

Jalpha at center (usage interne,su).

1.72. jxfus

Jalpha at xfus (su).

1.73. kidds_evol

facteur applique au coefficient de transport de Te pour la simulation de dents de scie en mode evolution.

1.74. li

internal self inductance (formule 3 of ITER FDR).

1.75. meff

effective mass (number of atomic mass, hydrogenoid ions).

1.76. modeh

confinement mode versu time: 0 = L et 1 = H.

1.77. mu0_nbi

initial value of pitch angle for NBI (V_{\parallel} / V').

1.78. n0a

number of cold neutral that input in plasma at the edge every second coming from recycling and gaz puff (s^{-1}).

1.79. n1m

volume averaged density of H + D + T ions (m^{-3}) or H+D if option.gaz = 11.

1.80. nDm

volume averaged density of deuterium ions (m^{-3}).

1.81. nTm

volume averaged density of tritium ions (m^{-3}); if option.gaz = 11, encode for volume averaged boron density; in this case the tritium density is null.

1.82. nb

number of convergence loops made.

1.83. nbar

line averaged electron density (m^{-3}).

1.84. nbar_nat

natural density (m^{-3}).

1.85. ndd

total DD neutrons per second (s^{-1}).

1.86. ndd_nbi_nbi

beam/beam DD neutrons per second (s^{-1}).

1.87. ndd_nbi_th

beam/plasma DD neutrons per second (s^{-1}).

1.88. ndd_th

plasma/plasma DD neutrons per second (s^{-1}).

1.89. ne0

estimation central electron density (m^{-3}).

1.90. nebord

estimation of plasma edge electron density (m^{-3}).

1.91. negr

Greenwald limit for electron density (m^{-3}).

1.92. nelim

plasma electron density temperature estimation near the divertor plate (m^{-3}).

1.93. nem

volume averaged electron density (m^{-3}).

1.94. neped

top electron pedestal density (m^{-3}).

1.95. nhe4m

volume averaged density of helium-4 (m^{-3}) in case of option.gaz == 5, adding accumulation of He4 on option.frhe0 * zs.nem for initial volume averaged He4 density.

1.96. nhem

volume averaged density of helium (m^{-3}); if option.gaz == 5, encode for volume averaged He3 density and option.frhe0 * zs.nem encode for volume averaged He4 density.

1.97. ni0

estimation of central ionic density (m^{-3}).

1.98. nibord

edge ion density (m^{-3}).

1.99. nim

sum of volume averaged ions density (m^{-3}).

1.100. nimpbm

volume averaged density of impurity (other than helium) (m^{-3}).

1.101. niped

top ion pedestal density (m^{-3}).

1.102. nmino

volume averaged ICRH minority density.

1.103. nsat

saturation electron density, use for the calculation of density (m^{-3}).

1.104. nwm

volume averaged tungsten density (m^{-3}); if option.Sn_fraction > 0, nwm become the sum of the density of W and Sn: $\langle n_W \rangle = (1 - \text{option.Sn_fraction}) * nwm$ & $\langle n_{Sn} \rangle = \text{option.Sn_fraction} * nwm$.

1.105. pbrem

Bremsstrahlung radiation losses (W).

1.106. pyclo

cyclotron radiation losses (W).

1.107. pddfus

fusion power from DD reactions (W).

1.108. peakdiv

divertor peak power surface density estimation (W/m^2).

1.109. pecrh

ECRH power(W).

1.110. pei

equipartition power (W).

1.111. pel

total thermal power deposition on electron, use in the calculation of T_i/T_e (W).

1.112. pel_fus

Alpha power going on electron (W).

1.113. pel_icrh

ICRH power going on electron (W).

1.114. pel_nbi

NBI power going on electron (W).

1.115. pelec

reactor electric power provide to the network (W).

1.116. peri

length of the LCMS (m).

1.117. pfus

total fusion power of alpha for D-T plasma and p-B11 plasma heating the plasma or from charged products for D + He3 (W).

1.118. pfus_loss

fusion power loss due to first orbit losses of fast alpha (estimation, W).

1.119. pfus_nbi

NBI induce fusion power heating the plasma (W).

1.120. pfus_th

fusion thermal power deposition of alpha particles (W).

1.121. phiplasma

toroidal magnetic flux due to the plasam (Wb).

1.122. picrh

ICRH power, decrease of ripple losses in TS (W).

1.123. picrh_th

ICRH thermal power deposition (W).

1.124. pin

total heat power (W).

1.125. pion

total thermal power deposition on ions, use in the calculation of T_i/T_e (W).

1.126. pion_fus

Alpha power going on ions (W).

1.127. pion_icrh

ICRH power going on ions (W).

1.128. pion_nbi

NBI power going on ions (W).

1.129. pioniz

power losses due to cold neutral ionization and charge exchange between ion and cold neutrals(W).

1.130. p_{ioniz_i}

power losses due to charge exchange between ion and cold neutrals (W).

1.131. p_{iqj}

peaking factor of current profile.

1.132. p_{iqnbi}

peaking factor of the NBI power deposition profile (su).

1.133. p_{lh}

LH power, decrease of ripple losses in TS (W).

1.134. p_{lh_th}

LH thermal power deposition (W).

1.135. p_{lhrip}

LH power loss in ripple for Tore Supra (W).

1.136. p_{lhthr}

additional power crossing the LCMS; must be compared to L->H threshold power (Ryter PPCF 2002).

1.137. p_{lim}

total power estimation on divertor plate (W).

1.138. p_{loss}

plasma losses power, as defined in ITER basis (W).

1.139. p_{lossl2h}

threshold power for transition from L mode to H mode for the selected scaling law (W).

1.140. p_{nbi}

NBI power (W).

1.141. p_{nbi_icrh}

equivalent beam power increase needs for neutron rate enhancement due to ICRH synergy effects (W).

1.142. pnbi_th

NBI thermal power deposition (W).

1.143. pohm

ohmic power (W).

1.144. poynting

flux of Poynting vector through the LCMS (W).

1.145. pped

total pressure @ pedestal top (Pa).

1.146. ppedmax

maximal allowed total pressure @ pedestal top (Pa).

1.147. prad

impurity radiation losses in core plasma, without Bremsstrahlung (W).

1.148. pradsol

radiation losses in the SOL (W).

1.149. priptherm

TS ripple losses, thermal part (W).

1.150. pth

thermal power input, define as $\tau_E \cdot P_{th} = W_{th}$ (W).

1.151. pttfus

fusion power from TT reactions (W).

1.152. pw

effective power define as $\tau_{eff} \cdot pw = W$ (W).

1.153. q0

estimation of central value of safety factor.

1.154. q95

safety factor @ 95% of poloidal flux.

1.155. qa

edge safety factor.

1.156. qeff

effective safety factor at the edge (computed with ITER formula).

1.157. qmin

estimation of minimal value of safety factor.

1.158. rm

$\sqrt{\Phi I / \pi B_0}$ of LCMS (m) .

1.159. rres

ICRF resonance layer radial position (m).

1.160. salpha

total number of alpha fusion particules from D-T reactions, from p-B11 reactions per second or from D-He3 reactions (s^{-1}).

1.161. salpha_he3

total number of helium-3 fusion particules per second from D-D reactions (s^{-1}).

1.162. salpha_he4

total number of helium-4 fusion particules per second from D-T and D-He3 reactions (s^{-1}).

1.163. salpha_n

total number of neutrons per second from D-D and D-T reactions (s^{-1}).

1.164. salpha_p

total number of proton fusion particules per second from D-He3 and D-D reactions (s^{-1}).

1.165. salpha_t

total number of tritium fusion particules per second from D-D reactions (s^{-1}).

1.166. sext

external plasma surface (m^2).

1.167. slh

rotation torque due to LHCD and ECCD (N m).

1.168. sn0fr

friction source on edge neutral that damps the toroidal rotation (N m).

1.169. snbi

rotation torque due to NBI (N m).

1.170. sp

poloidal plasma surface (m^2).

1.171. stf

number of wrong data in zs data structure containing NaN or Imag.

1.172. taue

energy confinement time (s).

1.173. taue_alt

energy confinement time, Helander & Sigmar definition (s).

1.174. tau ee

scale of electron heat confinement time (s).

1.175. tau ei

exchange electron/ion heat time (s).

1.176. tau h

confinement time of energy in H mode for the selected scaling law (s).

1.177. tau he

confinement time of helium impurity/ashes (s).

1.178. tauhe_h

confinement time of helium impurity/ashes in H mode (s).

1.179. tauhe_l

confinement time of helium impurity/ashes in L mode (s).

1.180. tauii

scale of ion heat confinement time (s).

1.181. tauip

carateristic time of R-L equivalent plasma circuit (s).

1.182. tauj

diffusion time of current (s).

1.183. taup

confinement time of matter (s).

1.184. taus_he

characteristic slowing down time for alpha particules (s).

1.185. taus_icrh

characteristic slowing down time for ICRH fast ions (s).

1.186. taus_nbi

characteristic slowing down time for NBI fast ions (s).

1.187. tauthl

confinement time of energy in L mode for the selected scaling law (s).

1.188. te0

estimation of central electron temperature (eV).

1.189. tebord

estimation of plasma edge electron temperature (eV).

1.190. telim

plasma electron temperature estimation near the divertor plate (eV).

1.191. tem

volume averaged electron temperature (eV).

1.192. temps

time slices vector.

1.193. teped

top electron pedestal temperature (eV).

1.194. tibord

edge ion temperature (eV).

1.195. tiped

top ion pedestal temperature (eV).

1.196. tite

volume averaged ratio T_i / T_e .

1.197. vloop

loop voltage, as $vloop \cdot iohm = pohm$ (V).

1.198. vmes

loop voltage as measured on a fixed magnetic loop (V).

1.199. vp

plasma volume (m^3).

1.200. w

total plasma energy (J).

1.201. wbp

poloidal field energy of the plasma (J).

1.202. wdia

$3/2$ perpendicular plasma energy (J).

1.203. wmagtor

toroidal magnetic plasma energy (J).

1.204. wrad

estimation of bulk volume averaged toroidal rotation velocity (rad/s).

1.205. wrlw

plasma energy contents for electron computed with RLW sacling law (J).

1.206. wrot

toroidal plasma rotation stored energy (J).

1.207. wth

thermal plasma energy (J).

1.208. xeccd

position of the maximum power deposition of ECCD(su).

1.209. xfus

Jalpha maximum position (su).

1.210. xitb

radius of itb due to a revserse shear (estimation,su).

1.211. xlh

position of the maximum power deposition of LH deposition maximum position (su).

1.212. xnbi

position of the maximum power deposition of NBI(su).

1.213. xped

pedestal normalized position.

1.214. xpoint

flag for diverted plasma; for plasma in limiter mode = 0; for plasma in divertor mode = 1.

1.215. xres

ICRF resonance layer normalized radius position.

1.216. zeff

plasma effective charge with alpha particles.

1.217. zeffsc

zeff scaling (private data, do not use).

1.218. zmszl

ratio between volume averaged zeff and line averaged zeff .

2. Substructure: profil0d

Time dependent profile data.

2.1. C2

C2 geometrical coefficient (see CRONOS technical document).

2.2. C3

C3 geometrical coefficient (see CRONOS technical document).

2.3. Raxe

of major radius of the centre of each flux surface (m).

2.4. Rsepa

radial coordinate of the LCMS points (m).

2.5. Zsepa

vertical coordinate of the LCMS points (m).

2.6. bpol

average poloidal magnetic field (T).

2.7. df2dpsi

$d(F^2)/d\Psi$ for FREEBIE ($4\pi T$).

2.8. dn

density diffusivity estimation (m^2/s).

2.9. dphidx

toroidal flux space derivative (Wb).

2.10. dpsidt

time derivative of poloidal flux (V).

2.11. dptotdpsi

$dP_{total}/d\Psi$ for FREEBIE ($2\pi A/m^3$).

2.12. drot

toroidal rotation diffusion coefficient (m^2/s).

2.13. dx

flux surface geometrical triangularity .

2.14. ej

ohmic power density (W/m^3).

2.15. epar

parallel electric field (V/m) .

2.16. epsi

of aspect ratio (m).

2.17. er

neoclassical radial electric field (V/m) = E_r / gradient(ρ).

2.18. eta

neoclassical resistivity ($\text{ohm} * \text{m}$).

2.19. fdia

diamagnetic function ($\text{T} * \text{m}$).

2.20. fprad

line radiative power shape; reserved to an internal used only.

2.21. frot

toroidal rotation moment flux (N m^{-1}).

2.22. ftrap

effective trap fraction profile .

2.23. fxdurplh

Hard X ray inversed profile.

2.24. ge

electrons flux (m^{-2}/s).

2.25. grho

$\langle |\text{gradient}(\rho)| \rangle$ (see CRONOS technical document).

2.26. grho2

$\langle |\text{gradient}(\rho)|^2 \rangle$ (see CRONOS technical document).

2.27. grho2r2

$\langle |\text{gradient}(\rho)|^2 / R^2 \rangle$ (see CRONOS technical document).

2.28. jboot

bootstrap current density (A/m^2).

2.29. jeccd

parallel current density source due to ECCD ($j = \langle J.B \rangle / B_0$, A/m^2).

2.30. jeff

$\langle J.B \rangle / B_0$ (A/m^2).

2.31. jfus

fast alpha bootstrap current density (A/m^2).

2.32. jfusshape

alpha particles bootstrap current density shape; reserved to an internal used only .

2.33. jfwcd

parallel current density source due to FWCD ($j = \langle J.B \rangle / B_0$, A/m^2).

2.34. jgs

averaged current density, raw computation for FREEBIE (A/m^2).

2.35. jlh

parallel current density source due to LHCD ($j = \langle J.B \rangle / B_0$, A/m^2).

2.36. jli

averaged current density, equivalent at j_{moy} in CRONOS (A/m^2).

2.37. jnbicd

parallel current density source due to NBICD ($j = \langle J.B \rangle / B_0$, A/m^2).

2.38. jnbishape

NBICD current density shape; reserved to an internal used only.

2.39. jni

total current density source (A/m^2).

2.40. jrun

average runaway current density (A/m^2).

2.41. kx

flux surface elongation .

2.42. n0

neutral density coming from edge, hot neutral (m^{-3}).

2.43. n0m

neutral density coming from edge, cold neutral (m^{-3}).

2.44. n1p

density of HDT ions (m^{-3}) or H+D if `option.gza = 1`.

2.45. nbinesource

electron source due to fast neutral ionisation ($\text{Number s}^{-1} \text{ m}^{-3}$).

2.46. nbishape_el

NBI power deposition on electrons shape; reserved to an internal used only.

2.47. nbishape_ion

NBI power deposition on ions shape; reserved to an internal used only.

2.48. nep

electron density (m^{-3}).

2.49. nhep

density of helium (m^{-3}) or helium 3 density if option.gza = 5.

2.50. nip

ions density (m^{-3}).

2.51. nwp

density of W when tungsten effects are taking into account (m^{-3}); if option.Sn_fraction > 0, this profile become the sum of the density of W and Sn: $n_W = (1 - \text{option.Sn_fraction}) * nwp$ & $n_{Sn} = \text{option.Sn_fraction} * nwp$.

2.52. nzp

density of main impurity (m^{-3}).

2.53. omega

plasma solid rotation frequency in toroidal direction (rad/s) $\frac{\sum(nk * mk * \langle V_{k,\phi} * R \rangle)}{\sum(nk * mk)}$.

2.54. palf

alpha power source (W/m^{-3}).

2.55. pbrem

bremsstrahlung power sink (W/m^3).

2.56. pcyclo

cyclotron radiation power source (W/m^3).

2.57. pecrh

power density due to ECRH (W/m^3).

2.58. pfus

power density heating the plasma due to fusion reactions (W/m^3).

2.59. pfus_ion

power density heating the plasma due to fusion reactions, coupled to ions (W/m^3).

2.60. pfweh

power density due to FWEH (W/m^3).

2.61. phi

toroidal flux (Wb).

2.62. picrh

power density due to ICRH minority scheme (W/m^3).

2.63. picrh_ion

power density due to ICRH minority scheme, coupled to ions (W/m^3).

2.64. pioniz

loss power due to cold neutral ionization and charge exchange between ions and cold neutrals (W m^{-3}).

2.65. pioniz_i

loss power due charge exchange between ions and cold neutrals (W m^{-3}).

2.66. pitch

NBI pitch angle profile = $\cos(\text{beam}, B)$.

2.67. plh

power density due to LH (W/m^3).

2.68. pnbi

power density due to NBI (W/m^3).

2.69. pnbi_ion

power density due to NBI, coupled to ions (W/m^3).

2.70. pohm

ohmic power deposition (W/m^3).

2.71. prad

radiated power sink (W/m^3).

2.72. ψ_i

poloidal flux (Wb).

2.73. p_{tot}

total pressure profile (Pa).

2.74. q_e

electron heat flux (W).

2.75. q_{ei}

electron to ion heat flux (W).

2.76. q_i

ion heat flux (W).

2.77. q_{jli}

safety factor.

2.78. r_{2i}

$\langle 1/R^2 \rangle$ (see CRONOS technical document).

2.79. r_i

$\langle 1/R \rangle$ (see CRONOS technical document).

2.80. r_{mx}

of average radius of each flux surface: $r_{mx} = \sqrt{\psi_i / \pi / b_0}$ (m).

2.81. rot_{lh}

toroidal torque source from LHCD and ECCD ($N\ m^{-2}$).

2.82. rot_{n0}

toroidal torque sink from edge neutral friction ($N\ m^{-2}$).

2.83. rot_{nbi}

toroidal torque source from NBI ($N\ m^{-2}$).

2.84. rtor

toroidal rotation moment density ($\text{kg m}^{-1} \text{s}^{-1}$).

2.85. s0

ionisation sources coming from edge, hot neutral (m^{-3}/s).

2.86. s0m

ionisation sources coming from edge, cold neutral (m^{-3}/s).

2.87. salf

alpha particles source ($\text{m}^{-3} \text{s}^{-1}$).

2.88. source_el

total heat power density coupled to electrons (W/m^3).

2.89. source_ion

total heat power density coupled to ions (W/m^3).

2.90. spelllet

equivalent continue source due to pellet injection (m^{-3}/s).

2.91. spr

surface element (m^2 , $\text{int}(\text{spr}, x= 0..1) = \text{plasma poloidal section surface}$) .

2.92. temps

vectors of time associate to the s (only time at which the s are computed are stored).

2.93. tep

electron temperature (eV).

2.94. tip

ion temperature (eV).

2.95. utheta

neoclassical poloidal rotation speed ,for main impurity ($\langle V_k . \theta \rangle / \langle B . \theta \rangle$, m/s/T).

2.96. vn

anormal density convection velocity estimation (m^2/s).

2.97. vpr

volume element (m^3 , $\text{int}(\text{vpr}, x=0..1) = \text{plasma volume}$).

2.98. vpr_tor

volume element, reserved for internal used (m^2 , $\text{rhomax} * \text{int}(\text{vpr}, x=0..1) = \text{plasma volume}$) .

2.99. vrot

toroidal rotation pinch coefficient (m/s).

2.100. vtheta

fluid velocity, theta component, at $R = R_{\text{max}}$ of each flux surface, for main impurity (m/s).

2.101. vtor

toroidal rotation speed (m/s), at $R = R_{\text{max}}$ of each flux surface, for main impurity .

2.102. ware

Ware pinch estimation (m/s).

2.103. web

rotation shear (NClass definition).

2.104. xie

electron diffusivity estimation (m^2/s).

2.105. xieshape

heat transport coefficient (Ke) shape without ITB effect .

2.106. xieshape_itb

heat transport coefficient (Ke) shape with ITB effect.

2.107. xii

ion diffusivity estimation (m^2/s).

2.108. xli

radial normalized coordinate (Lao coordinate $\sim r/a$, normalized to 1 at the edge)..

2.109. zeff

effective charge.