


The development of coupled NWP forecasting at the Met Office

Dan Copsey, Tim Graham, Michael Vellinga and Chris Harris.



Structure of presentation



Comparing coupled NWP vs atmosphere-only NWP



Operational implementation



Future changes after operational implementation

Introduction

- The Met Office plans to move to coupled NWP for its operational weather forecasts in the next few years.
- Results presented here is for PS41 coupled NWP which consists of components shown on right

UM atmosphere using
GA6.1 physics



JULES land surface
using GL8.0 physics



Internal
communication



OASIS3-MCT
coupler (every hour)




Internal
communication



NEMO ocean

using NEMO3.6 and GO6
physics



CICE sea ice using
CICE5.1.2 and GSI8
physics

Models to compare

	Operational atmosphere only NWP model (UNCPLD)	Coupled NWP model (CPLDNWP)
Physics options	PS41	PS41
Atmospheric horizontal resolution	N1280 = 10km grid spacing in mid latitudes	N1280 = 10km grid spacing in mid latitudes
Atmospheric vertical levels	70 levels	70 levels
Sea surface temperatures and sea ice	Provided by OSTIA SST and sea ice analysis and kept constant	Provided by NEMO/CICE
Atmospheric model's land sea mask	Derived from IGBP including large lakes as sea points	Derived from NEMO mask
Ocean/sea-ice horizontal resolution		ORCA025 = 25km grid spacing
Ocean vertical levels		75
Coupling interval		Every hour

Verification statistics

Based on RMSE statistics against UM analysis.


Green up arrow = CPLDNWP improved over UNCPLD

Blue down arrow = CPLDNWP degraded over UNCPLD

Almost all fields improved apart from 2m temperatures (T_{2m}). CPLDNWP penalised as not using OSTIA SSTs and OSTIA lakes (which both UM analysis and UNCPLD use).


1st Dec 2018 – 28th Feb 2019. 0Z forecasts.

% Difference (CPLDNWP vs.
Control) - overall -0.05%
RMSE against ownanal for 20181201 to 20190228




Met Office SST forecast performance vs persistence


a) MAE CPLD - persist. day 2




b) MAE CPLD - persist. day 7



c) VAR CPLD - persist. day 2



d) VAR CPLD - persist. day 7




Verification against FOAM analysis
MAE = mean absolute error
VAR = variance of difference
between forecast and observations

SST forecast improved in North
Pacific, North Atlantic and
southern sub-tropics

SST forecast degraded in
Southern Ocean


26th Sept 2018 – 25th Sept 2019. 0Z forecasts.

[Vellinga et al. 2020](#)




500 hPa geopotential height (Z500) and 250 hPa zonal wind (U250)


a) $\Delta RMSE$ Z500




b) ΔVar Z500



c) $\Delta RMSE$ U250



d) ΔVar U250




Plots are for T+168 (day 7)


Verification against UM analysis
VAR = variance of difference
between forecast and
observations

Largest improvements to the
atmosphere are near the
equator

26th Sept 2018 – 25th Sept 2019. 0Z fcst.



Met Office Tropical cyclone tracks



Track error superior counts averaged over all tropical cyclones:

CS = CPLDNWP is Superior

US = UNCPLD is Superior

SIM = Both systems Similar

Coupled model gives lower track errors most often


11th July 2017 – 1st Nov 2019. 0Z forecasts.

[Vellinga et al. 2020](#)

Operational implementation



Uncoupled & weakly coupled DA




Operational Implementation


- Plan to make coupled NWP with weakly coupled DA operational from Autumn 2021
- Resolution will be N1280L70 (~10km) ORCA025L75 for the deterministic model and N640 (~20km) ORCA025L75 for the ensemble
- Model version is expected to be GC4 which was frozen in March.
- Ensemble will use ocean analysis from the deterministic model + SST perturbations but development of an ocean ensemble DA system is underway for future implementation.

Future changes after operational implementation


Next Generation Modelling System (NGMS) at the Met Office



UM




LFRic




Atmospheric component no longer scalable on huge number of processors. Met Office is moving atmosphere component to a cube-sphere grid with new dynamical core (GungHo). This new model will be called the LFRic model.

Current coupled NWP



Future coupled NWP



Summary

- Using a coupled NWP model gives better verification scores than an atmosphere only model (except 2m temperatures).
- Remaining 2m temperature issues can be fixed by verifying vs a weakly coupled DA analysis and by using OSTIA lake SSTs.
- SST forecasts are improved over persistence, (except in Southern Ocean).
- 500 hPa heights, 250 hPa winds and tropical cyclones are all (on average) improved.
- We are aiming for the operational coupled NWP model to be GC4 in autumn 2021.
- The Met Office coupled NWP is moving to a new sea ice model (SI³ – GC5), a higher resolution ocean (ORCA12) and then a cubesphere grid atmosphere (LFRic – GC6), with optional wave model.