

Warm-up Exercises

Zheng Tian

[2017-04-01 Sat]

Question 1

In the simple linear regression model $Y_i = \beta_0 + \beta_1 X_1 + u_i$,

- A the intercept is typically small and unimportant.
- B $\beta_0 + \beta_1 X_1$ represents the population regression function.
- C the absolute value of the slope is typically between 0 and 1.
- D $\beta_0 + \beta_1 X_1$ represents the sample regression function.

Question 1

In the simple linear regression model $Y_i = \beta_0 + \beta_1 X_1 + u_i$,

- A the intercept is typically small and unimportant.
- B $\beta_0 + \beta_1 X_1$ represents the population regression function.
- C the absolute value of the slope is typically between 0 and 1.
- D $\beta_0 + \beta_1 X_1$ represents the sample regression function.

Answer: B

Question 1

In the simple linear regression model $Y_i = \beta_0 + \beta_1 X_1 + u_i$,

- A the intercept is typically small and unimportant.
- B $\beta_0 + \beta_1 X_1$ represents the population regression function.
- C the absolute value of the slope is typically between 0 and 1.
- D $\beta_0 + \beta_1 X_1$ represents the sample regression function.

Answer: B

Question 2

The OLS estimator is derived by

- A connecting the Y_i corresponding to the lowest X_i observation with the Y_i corresponding to the highest X_i observation.
- B making sure that the standard error of the regression equals the standard error of the slope estimator.
- C minimizing the sum of absolute residuals.
- D minimizing the sum of squared residuals.

Question 2

The OLS estimator is derived by

- A connecting the Y_i corresponding to the lowest X_i observation with the Y_i corresponding to the highest X_i observation.
- B making sure that the standard error of the regression equals the standard error of the slope estimator.
- C minimizing the sum of absolute residuals.
- D minimizing the sum of squared residuals.

Answer: D

Question 2

The OLS estimator is derived by

- A connecting the Y_i corresponding to the lowest X_i observation with the Y_i corresponding to the highest X_i observation.
- B making sure that the standard error of the regression equals the standard error of the slope estimator.
- C minimizing the sum of absolute residuals.
- D minimizing the sum of squared residuals.

Answer: D

Question 3

The following are all least squares assumptions with the exception of:

- A The conditional distribution of u_i given X_i has a mean of zero.
- B The explanatory variable in regression model is normally distributed.
- C $(X_i, Y_i), i = 1, \dots, n$ are independently and identically distributed.
- D Large outliers are unlikely.

Question 3

The following are all least squares assumptions with the exception of:

- A The conditional distribution of u_i given X_i has a mean of zero.
- B The explanatory variable in regression model is normally distributed.
- C $(X_i, Y_i), i = 1, \dots, n$ are independently and identically distributed.
- D Large outliers are unlikely.

Answer: B