

Definition 0.0.1 (Δ_h^k)

We define $\forall h > 0 : \Delta_h$ (h -central difference) of $f : \mathbb{R} \rightarrow \mathbb{R}$ as follows:

$$\Delta_h f(t) = f(t+h) - f(t-h), \quad (1)$$

and $\forall k \in \mathbb{N} : \Delta_h^k$ is defined as applying Δ_h applying k times and $k = 0$ as the identity. Note that Δ_h^k is a linear operator. We will always take differences in respect to the t variable.

Lemma 0.0.2 ($\Delta_h^k e^{at}$)

$$\forall k \in \mathbb{N} : \Delta_h^k e^{at} = e^{at} (e^{ah} - e^{-ah})^k. \quad (2)$$

Proof. This follows simply by induction. $k = 0$ is trivial. Base case:

$$\Delta_h e^{at} = (e^{a(t+h)} - e^{a(t-h)}) \quad (3)$$

$$= e^{at} (e^{ah} - e^{-ah}). \quad (4)$$

Induction hypothesis:

$$\Delta_h^{k-1} e^{at} = e^{at} (e^{ah} - e^{-ah})^{k-1}. \quad (5)$$

Induction step:

$$\Delta_h^k e^{at} = \Delta_h \Delta_h^{k-1} (e^{at}) \quad (6)$$

$$= \Delta_h \left(e^{at} (e^{ah} - e^{-ah})^{k-1} \right) \quad (7)$$

$$= \Delta_h (e^{at}) (e^{ah} - e^{-ah})^{k-1} \quad (8)$$

$$= e^{at} (e^{ah} - e^{-ah}) (e^{ah} - e^{-ah})^{k-1} \quad (9)$$

$$= e^{at} (e^{ah} - e^{-ah})^k. \quad (10)$$

□

Lemma 0.0.3 (Δc)

$$\forall c \in \mathbb{R} : \Delta_h c = c - c = 0. \quad (11)$$

Lemma 0.0.4 ($\Delta_h t^n$)

$\forall n > 0 \in \mathbb{N} : \Delta_h t^n$ is a polynomial of degree $n - 1$ with leading coefficient $2hn$.

Proof.

$$\Delta_h t^n = (t+h)^n - (t-h)^n. \quad (12)$$

So $\Delta_h t^n$ is a polynomial, verify the degree and leading coefficient by looking at its derivatives (or use the binomial theorem):

$$\frac{d^n}{dt^n} ((t+h)^n - (t-h)^n) = n! - n! = 0. \quad (13)$$

$$\frac{d^{n-1}}{dt^{n-1}} ((t+h)^n - (t-h)^n) = n!(t+h) - n!(t-h). \quad (14)$$

$$= 2hn! \neq 0. \quad (15)$$

□

Lemma 0.0.5 ($\Delta_h^k t^n$)

$\forall n, k \in \mathbb{N}, n < k : \Delta_h^k t^n = 0.$

Proof. Follows from the previous 2 lemmas and linearity of Δ_h^k . □

Lemma 0.0.6 ($\Delta_h^k t^k$)

$\forall k \in \mathbb{N} : \Delta_h^k t^k = (2h)^k k!.$

Proof. Proof by induction. Base case $k = 0$ is trivial. Induction hypothesis:

$$\forall l < k : \Delta_h^l t^l = (2h)^l l!. \quad (16)$$

Induction step, use previous lemmas:

$$\Delta_h^k t^k = \Delta_h^{k-1} \Delta_h t^k \quad (17)$$

$$= \Delta_h^{k-1} ((t+h)^k - (t-h)^k) \quad (18)$$

$$= \Delta_h^{k-1} (2hkt^{k-1} + \dots) \quad (19)$$

$$= \Delta_h^{k-1} (2hkt^{k-1}) \quad (20)$$

$$= 2h\Delta_h^{k-1} (t^{k-1}) \quad (21)$$

$$= 2hk(2h)^{k-1}(k-1)! \quad (22)$$

$$= (2h)^k k! \quad (23)$$

□

Theorem 0.0.7 (central finite difference formula)

$\forall k \in \mathbb{N} \forall k$ -continuously differentiable f 's at 0:

$$f^{(k)}(0) = \lim_{h \rightarrow 0} \frac{\Delta_h^k|_{t=0} f(t)}{(2h)^k}. \quad (24)$$

Proof. Follows from Taylor's theorem and previous lemma.

$$\lim_{h \rightarrow 0} \frac{\Delta_h^k|_{t=0} f(t)}{(2h)^k} \quad (25)$$

$$= \lim_{h \rightarrow 0} \frac{\Delta_h^k|_{t=0} \left(\sum_{n=0}^{k-1} \frac{f^{(n)}(0)}{n!} t^n + \left(\frac{f^{(k)}(0)}{k!} + h_k(t) \right) t^k \right)}{(2h)^k} \quad (26)$$

$$= \lim_{h \rightarrow 0} \frac{\Delta_h^k|_{t=0} \left(\left(\frac{f^{(k)}(0)}{k!} + h_k(t) \right) t^k \right)}{(2h)^k} \quad (27)$$

$$= \lim_{h \rightarrow 0} \frac{\left(\frac{f^{(k)}(0)}{k!} + h_k(t) \right) \Delta_h^k|_{t=0} (t^k)}{(2h)^k} \quad (28)$$

$$= \lim_{h \rightarrow 0} \frac{\left(\frac{f^{(k)}(0)}{k!} + h_k(t) \right) (2h)^k k!}{(2h)^k} \quad (29)$$

$$= \lim_{h \rightarrow 0} f^{(k)}(0) + h_k(t) k! \quad (30)$$

$$= f^{(k)}(0) \quad (31)$$

□

Lemma 0.0.8

$\forall k \in \mathbb{N} : \phi_X(t)$ is $2k$ -time continuous differentiable $\Rightarrow E[X^{2k}] < \infty$

Proof. Use the central finite difference formula and previous lemma's:

$$|\phi_X^{(2k)}(0)| = \left| \lim_{h \rightarrow 0} \frac{\Delta_h^{2k}|_{t=0} \phi_X(t)}{(2h)^{2k}} \right| \quad (32)$$

$$= \left| \lim_{h \rightarrow 0} \frac{\Delta_h^{2k}|_{t=0} E[e^{itX}]}{(2h)^{2k}} \right| \quad (33)$$

$$= \left| \lim_{h \rightarrow 0} \frac{E[\Delta_h^{2k}|_{t=0} e^{itX}]}{(2h)^{2k}} \right| \quad (34)$$

$$= \left| \lim_{h \rightarrow 0} \frac{E[e^{itX}|_{t=0} (e^{ihX} - e^{-ihX})^{2k}]}{(2h)^{2k}} \right| \quad (35)$$

$$= \left| \lim_{h \rightarrow 0} \frac{E[(e^{ihX} - e^{-ihX})^{2k}]}{(2h)^{2k}} \right| \quad (36)$$

$$= \left| \lim_{h \rightarrow 0} \frac{1}{i^{2k}} E \left[\frac{(e^{ihX} - e^{-ihX})^{2k}}{(2h)^{2k}} \right] \right| \quad (37)$$

$$= \lim_{h \rightarrow 0} E \left[\frac{\sin(Xh)^{2k}}{h^{2k}} \right] \quad (38)$$

$$= \lim_{h \rightarrow 0} E \left[X^{2k} \frac{\sin(Xh)^{2k}}{X^{2k} h^{2k}} \right] \quad (39)$$

$$= \lim_{h \rightarrow 0} E[X^{2k} \text{sinc}(Xh)^{2k}] \quad (40)$$

$$\geq E[X^{2k}] \quad (41)$$

Where last line follows by Fatou's lemma. □