



MASTERPROEF SCRIPTIE

# *Unbiased Monte Carlo for Recursive Integrals*

Auteur: *Isidoor Pinillo Esquivel*

Promotor: *Wim Vanroose*

ACADEMIEJAAR 2022-2023

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Introductory Example . . . . .	2
1.2	Contributions . . . . .	3
1.3	Related Work . . . . .	3
<b>2</b>	<b>Background</b>	<b>3</b>
2.1	Modifying Monte Carlo . . . . .	3
2.2	Monte Carlo Trapezoidal Rule . . . . .	5
2.3	Unbiased Non-Linearity . . . . .	5
2.4	Recursion . . . . .	7
2.5	Green Functions . . . . .	8
<b>3</b>	<b>1-Dimensional Recursive Integrals</b>	<b>8</b>
3.1	Linear Recursive Integrals . . . . .	8
3.2	IVPs ODEs . . . . .	8
3.3	BVPs ODEs . . . . .	8
<b>4</b>	<b>Higher Dimensional Recursive Integrals</b>	<b>8</b>
4.1	Complicated Geometry . . . . .	8
4.2	Recursive Brownian Motion . . . . .	8
4.3	Heat Equation . . . . .	9
4.4	Wave Equation . . . . .	9
<b>5</b>	<b>Appendix</b>	<b>11</b>

## Abstract

We will write this at the end.

# 1 Introduction

## 1.1 Introductory Example

To get familiar with Monte Carlo for estimating recursive integrals we demonstrate it on following problem:

$$y' = y, y(0) = 1. \quad (1)$$

By integrating both sides of (1) following integral equation can be derived:

$$y(t) = 1 + \int_0^t y(s)ds. \quad (2)$$

Equation (2) is a recursive integral equation or to be more specific a linear Volterra integral equation of the second type. By naively using Monte Carlo on the recursive integral of equation (2) one derives following estimator:

$$Y(t) = 1 + tY(Ut).$$

where  $U = \text{Uniform}(0, 1)$ . If  $y$  is well behaved then  $E[Y(t)] = y(t)$  but we can't calculate  $Y(t)$  without accesses to  $y(s), s < t$ . Notice that we can replace  $y$  by a unbiased estimator of it without changing  $E[Y(t)] = y(t)$  by the law of total expectance ( $E[X] = E[E[X|Z]]$ ). By replacing  $y$  by  $Y$  itself we obtain a recursive expression for  $Y$ :

$$Y(t) = 1 + tY(Ut). \quad (3)$$

Equation (3) is a recursive random variable equation (RRVE). If you would implement equation (3) with recursion it will run indefinitely. A biased way of around this is by approximating  $Y(t) \approx 1$  near  $t = 0$ . Later we discuss Russian roulette (2.1.1) which can be used as an unbiased stopping mechanism.

### Python Code 1.1.1 (implementation of (3))

```
1 from random import random as U
2 def Y(t, eps): return 1 + t*Y(U()*t, eps) if t > eps else 1
3 def y(t, eps, nsim):
4     return sum(Y(t, eps) for _ in range(nsim))/nsim
5 print(f"y(1) approx {y(1,0.01,10**3)}")
6 # y(1) approx 2.710602603240193
```

An issue with (1.1.1) is that the variance increases rapidly when  $t$  increases. Which we later solve in the section on ODEs. Note that (1.1.1) keeps desirable properties from unbiased Monte Carlo methods such as: being embarrassingly parallel, robustness and having simple error estimates.

## 1.2 Contributions

We write this at the end. Probably a lot of conjectures.

## 1.3 Related Work

work on

- alternative methods for recursive integrals
- MC work on ODEs
- MC work on PDEs
- WoS

This is just to give a general overview we probably reference specific ideas when we first introduce them.

# 2 Background

## 2.1 Modifying Monte Carlo

Once we have a RRVE it is possible to modify it to have more desirable properties. Our favorite work that discusses these techniques is [Vea].

**Definition 2.1.1** (Russian roulette)

Define Russian roulette on  $X$  with free parameters  $Y_1, Y_2 : E[Y_1] = E[Y_2]$ ,  $p \in [0, 1]$  and  $U$  independent of  $Y_1, Y_2, X$  the following way:

$$X \rightarrow \begin{cases} \frac{1}{p}(X - (1-p)Y_1) & \text{if } U < p \\ Y_2 & \text{else} \end{cases}.$$

The idea behind Russian roulette is replacing  $X$  with a cheaper approximation  $Y_2$  sometimes.

**Example 2.1.2**

Say that we are interested in estimating  $E[Z]$  with  $Z$  defined in the following way:

$$Z = U + \frac{f(U)}{1000}.$$

where  $f : \mathbb{R} \rightarrow [0, 1]$  expensive to compute. Estimating  $Z$  directly would require calling  $f$  each simulation. We can modify  $Z$  to

$$\tilde{Z} = U + B\left(\frac{1}{100}\right) \frac{f(U)}{10}.$$

where  $B(\frac{1}{100}) \sim \text{Bernouli}(\frac{1}{100})$ . Now  $\tilde{Z}$  just requires calling  $f$  on average once every 100 simulations with the variance only increasing slightly compared to  $Z$ .

Maybe this wasn't the best example because you could also estimate the expectance of the 2 terms of  $Z$  separately.

**Example 2.1.3** (Russian roulette on (3))

Russian roulette can fix the indefinite recursion issue of equation (3) by approximating  $Y$  near  $t = 0$  with 1. Concretely we replace the  $t$  in front of the recursive term with  $B(t) \sim \text{Bernouli}(t)$  when  $t < 1$ .

$$Y(t) = \begin{cases} 1 + B(t)Y(Ut) & \text{if } t < 1 \\ 1 + tY(Ut) & \text{else} \end{cases}.$$

**Python Code 2.1.4** (Russian roulette on (3))

```
1 from random import random as U
2 def Y(t):
3     if t>1: return 1 + t*Y(U()*t)
4     return 1 + Y(U()*t) if U() < t else 1
5 def y(t, nsim): return sum(Y(t) for _ in range(nsim))/nsim
6 print(f"y(1) approx {y(1,10**3)}")
7 # y(1) approx 2.698
```

**Definition 2.1.5** (splitting)

Splitting  $X$  means using multiple  $X_j \sim X$  not independent per se to lower variance by averaging them:

$$\bar{X} = \frac{1}{N} \sum_{j=1}^N X_j.$$

Splitting the recursive term in a RRVE can lead to (additive) branching recursion. Extra care should be taken that all branches get terminated with probability 1. This can be achieved by termination strategies already discussed and later we discuss coupled recursion for alleviating additive branching recursion in RRVEs.

**Example 2.1.6** (splitting on (3))

We can "split" the recursive term of (3) in 2:

$$Y(t) = 1 + \frac{t}{2}(Y_1(Ut) + Y_2(Ut)).$$

with  $Y_1(t), Y_2(t)$  i.i.d.  $Y(t)$ .

**Python Code 2.1.7** (splitting on (3))

```
1 from random import random as U
2 def Y(t):
3     u = U()
4     if t > 1:
5         return 1 + t*(Y(u*t)+Y(u*t))/2
6     return 1 + (Y(u*t)+Y(u*t))/2 if U() < t else 1
7 def y(t, nsim): return sum(Y(t) for _ in range(nsim))/nsim
8 print(f"y(1) approx {y(1,10**3)}")
9 # y(1) approx 2.73747265625
```

**Definition 2.1.8** (2-level MC)

2-level MC on  $X$  with parameters  $\tilde{X}, Y : E[\tilde{X}] = E[Y]$ :

$$X \rightarrow X - \tilde{X} + Y.$$

This gets discussed in [Gil13].

**Definition 2.1.9** (control variates)

Control variate on  $f(X)$  is

$$f(X) \rightarrow f(X) - \tilde{f}(X) + E[\tilde{f}(X)].$$

Control variates are a special case of 2-level MC. Usually  $\tilde{f}$  is an approximation of  $f$  to reduce variance.

**Example 2.1.10** (control variate on (3))

To make a control variate for (3) that reduces variance we use following approximation of  $y(t) \approx 1 + t$ :

$$Y(t) = 1 + t + \frac{t^2}{2} + t(Y(Ut) - 1 - Ut).$$

Notice that we can cancel the constant term of the control variate but that would affect the Russian roulette negatively.

**Python Code 2.1.11** (control variate on (3))

```

1 from random import random as U
2 def Y(t):
3     u = U()
4     if t > 1: return 1+t**2/2 + t*(Y(u*t)-u*t)
5     return 1 + t + t**2/2 + (Y(u*t)-1-u*t if U() < t else 0)
6 def y(t, nsim): return sum(Y(t) for _ in range(nsim))/nsim
7 print(f"y(1) approx {y(1,10**3)}")
8 # y(1) approx 2.734827303480301

```

## 2.2 Monte Carlo Trapezoidal Rule

The Monte Carlo trapezoidal rule has some interesting convergence behavior [Wu20]. We aren't yet comfortable with our understanding of the randomized trapezoidal rule.

comparing normal vs Monte Carlo trapezoidal rule and highlighting the "half variance phenomenon". + maybe integrating polynomials for intuition

## 2.3 Unbiased Non-Linearity

At first sight it looks only possible to deal with linear problems in an unbiased way but by using independent samples it possible to deal with polynomial non-linearity's which practically extend to any continuous functions by the Weierstrass approximation theorem. A similar approach to non-linearity can be found in [ET19]. It is not always easy to transform non-linearity into polynomials but it is not difficult to come up with biased alternative approaches based on linearization or approximate polynomial non-linearity.

**Example 2.3.1** ( $y' = y^2$ )

Let's do following example:

$$y' = y^2. \quad (4)$$

with  $y(1) = -1$ . This has solution  $-\frac{1}{t}$ . Integrate both sides of equation (4) to arrive at following integral equation:

$$y(t) = -1 + \int_1^t y(s)y(s)ds. \quad (5)$$

To estimate the recursive integral in equation (2) we use 2 independent estimators  $Y_1, Y_2$  of  $y$ :

$$Y(t) = -1 + (t-1)Y_1(S)Y_2(S).$$

With  $S \sim \text{Uniform}(1, t)$ . This is a branching RRVE this is typical when dealing with non-linearity.

**Python Code 2.3.2** ( $y' = y^2$ )

```

1 from random import random as U
2 def Y(t):
3     if t>2: raise Exception("doesn't support t>2")
4     S = U()*(t-1)+1
5     # Y(u)**2 != Y(u)*Y(u) !!!
6     return -1 + Y(S)*Y(S) if U()<t-1 else -1
7 def y(t, nsim): return sum(Y(t) for _ in range(nsim))/nsim
8 print(f"y(2) approx {y(2,10**3)}")
9 # y(2) approx -0.488

```

**Example 2.3.3** ( $e^{E[X]}$ )

$e^{\int x(s)ds}$  is common expression encountered when studying ODEs. In this example we demonstrate how you can generate unbiased estimates of  $e^{E[X]}$  with simulations of  $X$ . The Taylor series of  $e^x$  is:

$$e^{E[X]} = \sum_{n=0}^{\infty} \frac{E^n[X]}{n!} \quad (6)$$

$$= 1 + \frac{1}{1}E[X] \left( 1 + \frac{1}{2}E[X] \left( 1 + \frac{1}{3}E[X] (1 + \dots) \right) \right). \quad (7)$$

Change the fractions of equation (7) to Bernoulli processes and replace all  $X$  with independent  $X_j$  with  $E[X] = E[X_i]$ .

$$\begin{aligned}
e^{E[X]} &= E \left[ 1 + B \left( \frac{1}{1} \right) E[X_1] \left( 1 + B \left( \frac{1}{2} \right) E[X_2] \left( 1 + B \left( \frac{1}{3} \right) E[X_3] (1 + \dots) \right) \right) \right] \\
&= E \left[ 1 + B \left( \frac{1}{1} \right) X_1 \left( 1 + B \left( \frac{1}{2} \right) X_2 \left( 1 + B \left( \frac{1}{3} \right) X_3 (1 + \dots) \right) \right) \right]
\end{aligned}$$

What is inside the expectation is something that we can simulate with simulations of  $X_j$ .

### Python Code 2.3.4 ( $e^{E[X]}$ )

The following python code estimates  $e^{\int_0^t s^2 ds}$ :

```
1 from random import random as U
2 from math import exp
3 def X(t): return -t**3*U()**2
4 def num_B(i): # = depth of Bernoulli's = 1
5     return num_B(i+1) if U()*i < 1 else i-1
6 def res(n, t): return 1 + X(t)*res(n-1, t) if n != 0 else 1
7 def expE(t): return res(num_B(0), t)
8
9 t, nsim = 1, 10**3
10 sol = sum(expE(t) for _ in range(nsim))/nsim
11 exact = exp(-t**3/3)
12 print(f"sol = {sol} %error={(sol- exact)/exact}")
13 #sol = 0.7075010309320893 %error=-0.01260277046
```

## 2.4 Recursion

In this section we discuss recursion related techniques.

### Technique 2.4.1 (coupled recursion)

The idea behind coupled recursion is sharing recursion calls of multiple RVVEs for simulation. This does make them dependent. This is like assuming 2 induction hypotheses at the same time and proving both inductions steps at the same time vs doing separate induction proofs. Which should be easier because you have accesses to more assumptions at the same time.

### Example 2.4.2 (coupled recursion)

Lets say you are interested in calculating the sensitivity of the solution of an ODE to a parameter  $a$ :

$$y' = ay, y(0) = 1 \Rightarrow \quad (8)$$

$$\partial_a y' = y + a \partial_a y' \quad (9)$$

Turn (8) and (9) into RRVEs. To emphasize that they are coupled, that they should recurse together we write them in a matrix equation:

$$\begin{bmatrix} Y(t) \\ \partial_a Y(t) \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} a & 0 \\ 1 & a \end{bmatrix} \begin{bmatrix} Y(Ut) \\ \partial_a Y(Ut) \end{bmatrix}. \quad (10)$$

### Python Code 2.4.3 (implementation of (10))

```
1 from random import random as U
2 import numpy as np
3 def X(t, a): # only supports t<1
4     q, A = np.array([1, 0]), np.array([[a, 0], [1, a]])
5     return q + A @ X(U()*t, a) if U() < t else q
6 def sol(t, a, nsim): return sum(X(t, a) for _ in
7     range(nsim))/nsim
8 print(f"x(1,1) = {sol(1,1,10**3)}")
9 # x(1,1) = [2.7179 2.7104]
```



**Example 2.4.4** (recursion in recursion)  
maybe induction in induction proof example

**Example 2.4.5** (tail recursion)  
discuss problems with implementing recursion and solutions.  
inverse problem example

## 2.5 Green Functions

green function stuff that we will be needing, we aren't sure in how much detail we're going to go.

**Example 2.5.1** (numerical green functions)  
There will be probably some green functions that we need that don't have an analytic expression yet.

## 3 1-Dimensional Recursive Integrals

### 3.1 Linear Recursive Integrals

We have algo in mind for this case based on coupled recursion on disjunct sets.

### 3.2 IVPs ODEs

An IVP example probably using DRRMC maybe compare it to parareal. Maybe also non-linear algo

### 3.3 BVPs ODEs

A BVP example using yet another algo that hopefully has the half variance phenomenon.

## 4 Higher Dimensional Recursive Integrals

### 4.1 Complicated Geometry

**Example 4.1.1** (nasty 2D integral)  
2D integral that is difficult because of its geometry

### 4.2 Recursive Brownian Motion

WoS like way to simulate Brownian motion which is related to the green function of the heat equation

**Example 4.2.1** (recursive Brownian motion)  
see period5

### **4.3 Heat Equation**

a geometric robust way to solve the heat equation and maybe a higher order method to solve the heat equation

### **4.4 Wave Equation**

probably won't get to it

## References

- [ET19] S. M. Ermakov and T. M. Tovstik. “Monte Carlo Method for Solving ODE Systems”. en. In: *Vestnik St. Petersburg University, Mathematics* 52.3 (July 2019), pp. 272–280. issn: 1063-4541, 1934-7855. DOI: 10.1134/S1063454119030087. URL: <https://link.springer.com/10.1134/S1063454119030087> (visited on 01/20/2023).
- [Gil13] Michael B. Giles. *Multilevel Monte Carlo methods*. en. arXiv:1304.5472 [math]. Apr. 2013. URL: <http://arxiv.org/abs/1304.5472> (visited on 12/04/2022).
- [Vea] Eric Veach. “ROBUST MONTE CARLO METHODS FOR LIGHT TRANSPORT SIMULATION”. en. In: ().
- [Wu20] Yue Wu. *A randomised trapezoidal quadrature*. en. arXiv:2011.15086 [cs, math]. Dec. 2020. URL: <http://arxiv.org/abs/2011.15086> (visited on 03/06/2023).

## 5 Appendix

Derivation of the green functions and some expressions.