


Nanomagnetism I:

Theoretical and Experimental Techniques in the Study of Nanomagnetic Systems

Sean Langridge

Large Scale Structures Group
Rutherford Appleton Laboratory, Chilton, Oxfordshire


s.langridge@rl.ac.uk



1

The big picture

- In addition the surfaces and interfaces are particularly important
- Why?
- A break in symmetry




The length scale from

- $10^{-10} - 10^{-6}$ m
- covers a huge range of condensed matter science and technological applications



2


Applications



OSNS10
September 2007

3

Relevance of scattering techniques to nanomagnetism



Adapted from I.K. Schuller *et al.* J. Magn. Magn. Mater. 200 (1999) 571.



4

References

- Polarized Neutrons, W.G. Williams, Oxford (1988)
- Theory of Magnetic neutron and photon scattering, E. Balcar & S.W. Lovesey, Oxford (1988)
- Introduction to Thermal Neutron Scattering, G.L. Squires, Cambridge (1978)
- Elements of Modern X-Ray Physics, Als-Nielsen and McMorrow, Wiley & Sons (2001)
- Magnetism: from fundamentals to nanoscale dynamics, Stohr and Siegmann, Springer (2006)
- www.ill.fr
- www.isis.rl.ac.uk
- www.esrf.fr
- www.diamond.ac.uk




OSNS10
September 2007
ISIS
5

Neutron and X-ray compendiums

<http://xdb.lbl.gov/>



<http://www.ill.fr/>



OSNS10
September 2007
ISIS
6

Complementarity

- Bulk measurements
 - Magnetometry
 - Transport...



- X-rays
 - Element specific
 - Good Q-resolution
 - Weak interaction
 - High incident flux
 - Poor energy resolution (?)
 - Polarisation analysis
- Imaging
 - XPEEM
 - Lorentz



OSNS10
September 2007
ISIS
7


Science & Technology Facilities Council

ISIS

7

Motivation

- Spin-Engineering
 - Transport of spin polarised currents
 - Spin Torque etc.
- Buried Interfaces
 - c.f. μ -scopy
 - Need for large area patterning if PN to contribute
- Interfacial Phenomena
 - Influence on GMR
 - Exchange bias
 - Enhanced/reduced magnetic moments
 - Pinned/free moments
 - Magnetic Roughness



OSNS10
September 2007
ISIS
8

<http://www.research.lbn.com>

Basic Theoretical Ideas

Polarised Neutron Cross-Section



Basic Ideas-I

- Magnetic dipole moment:
- Differential cross-section

$$\frac{d^2\sigma}{d\Omega dE} = \frac{k}{k'} \left(\frac{m}{2\pi\hbar^2} \right)^2 \left| \langle k' \sigma' \lambda' | V_m | k \sigma \lambda \rangle \right|^2 \delta(E_\lambda - E_{\lambda'} + \hbar\omega)$$

(See e.g. Squires eqn 7.15)

$$\begin{aligned} \hat{V}_m &= -\hat{\mu}_n \cdot \mathbf{B} \\ \langle k' | \hat{V}_m | k \rangle &= -r_0 \hat{\mathbf{e}} \cdot \hat{\mathbf{Q}}_\perp \\ \hat{\mathbf{Q}}_\perp &= \sum_i \exp(i\mathbf{k} \cdot \mathbf{r}_i) \left(\mathbf{k} \times (\hat{\mathbf{s}}_i \times \mathbf{k}) - \frac{i}{\hbar\kappa} (\mathbf{k}_i \times \hat{\mathbf{p}}_i) \right) \\ \mathbf{Q} &= -\frac{1}{2\mu_B} \mathbf{M}(\kappa) \end{aligned}$$

$$Q = Q_s + Q_l = -\frac{1}{2\mu_B} \mathbf{M}(\kappa)$$



Historical Overview

- Pre-reactor
- Bloch 1936
 - theoretical description of the interaction between magnetic atoms and "magnetic" neutrons
- Schrödinger 1937
 - Detailed QM treatment
 - Phys Rev 51 544
- Halpern *et al.* (1937-1941)
 - Nuclear scattering
 - Paramagnetic scattering
- Alvarez and Block (1940)
 - Determination of the neutron moment
 - Phys Rev 57,111
- Blume (1963)
 - Polarisation effects in the magnetic elastic scattering of slow neutrons Phys. Rev. 130 (1963) 16702
- Moon *et al.* (1969)
 - Polarisation Analysis of Thermal Neutron scattering Phys. Rev. 181 (1969) 920
- Nobel Prize (1994)
 - "for pioneering contributions to the development of neutron scattering techniques for studies of condensed matter"



- Halpern *et al.* (1941)
 - Polarisation by reflection from magnetised mirrors
- Mid-late 1940's Intense reactor sources
- Shull *et al.* (1951)
 - Neutron Scattering and polarisation by ferromagnetic materials Phys. Rev. 84 (1951) 912
- Blume (1963)
 - Polarisation effects in the magnetic elastic scattering of slow neutrons Phys. Rev. 130 (1963) 16702
- Moon *et al.* (1969)
 - Polarisation Analysis of Thermal Neutron scattering Phys. Rev. 181 (1969) 920
- Nobel Prize (1994)
 - "for pioneering contributions to the development of neutron scattering techniques for studies of condensed matter"



Polarisation Analysis

$$\left(\frac{d^2\sigma}{d\Omega dE'} \right)_{\sigma\lambda \rightarrow \sigma'\lambda'} = \frac{k'}{k} \left| \sum_i \langle \sigma' \lambda' | \sigma \cdot \mathbf{Q}_\perp | \sigma \lambda \rangle \right|^2 \delta(E_\lambda - E_{\lambda'} + \hbar\omega)$$


- Components of the magnetisation, \mathbf{m} give rise to
- $\mathbf{m} \parallel \mathbf{H}$: Non Spin Flip Scattering (NSF)
- $\mathbf{m} \perp \mathbf{H}$: Spin Flip Scattering (SF)

$$\langle \uparrow | \sigma \cdot \mathbf{Q}_\perp | \uparrow \rangle = Q_{\perp z} \quad (1)$$


$$\langle \downarrow | \sigma \cdot \mathbf{Q}_\perp | \downarrow \rangle = -Q_{\perp z} \quad (2)$$

$$\langle \uparrow | \sigma \cdot \mathbf{Q}_\perp | \downarrow \rangle = Q_{\perp x} + iQ_{\perp y} \quad (3)$$

$$\langle \downarrow | \sigma \cdot \mathbf{Q}_\perp | \uparrow \rangle = Q_{\perp x} - iQ_{\perp y} \quad (4)$$




Spin Flip and non-spin flip



OSNS10
September 2007
 ISIS

13

Resonance Enhancement



J. P. Hamon, J. P., et al. 1988, Phys. Rev. Lett. **61**, 1245
J. P. Hill and D. F. McMorrow, Acta Crystallogr., Sect. A: Found. Crystallogr. **52**, 236 (1996)

OSNS10
September 2007
 ISIS

14


Specular Scattering

Neutron and x-ray reflectivity

Science & Technology Facilities Council
 ISIS

Spin polarised Neutron Reflectivity

- Measure the reflected neutrons as a function of their perpendicular momentum and spin eigenstate
- $k^\pm = \sqrt{k - 4\pi N(b \pm cB)}$
- Can observe the magnitude and orientation of atomic magnetic moments in thin films and multilayer media.
- Probe length scale (<1nm to >1000nm): covers many aspects of thin film structure and magnetism
- Complementary to:
 - VSM/SQUID
 - MOKE average magnetisation over the sample thickness
 - SEMPA, Lorentz surface domain magnetisation
 - XMCD/XRMS-element specific
- PNR gives the microscopic *in-plane vector* magnetisation *depth profile*.





OSNS10
September 2007
 ISIS

16

Index of Refraction: Neutrons

$$n = \frac{\sin \gamma_i}{\sin \gamma_t} = \frac{|\vec{k}_t|}{|\vec{k}_i|}$$

$$\begin{aligned} n^2 &= \frac{|\vec{k}_t|^2}{|\vec{k}_i|^2} \\ &= \frac{E_t}{E_i} \quad V_n = \frac{2\pi\hbar^2}{m_n} Nb \\ &= \frac{E_i - V_n}{E_i} \\ &= 1 - \frac{4\pi}{k_i^2} Nb \end{aligned}$$



OSNS10
September 2007
ISIS


17

Critical Reflection

- $\frac{\cos \alpha_i}{\cos \alpha_f} = \frac{n_2}{n_1}$
- At the critical angle
 $\frac{\cos \alpha_i}{\cos \theta} = n$

$$\begin{aligned} Q_c &= \frac{4\pi}{\lambda} 2k \sin \alpha_c \\ &= 2k \sqrt{1 - \cos^2 \alpha_c} \\ &= \sqrt{4k^2(1 - n^2)} \\ &\cong \sqrt{4k^2 \cdot 2\delta} \\ &= \sqrt{16\pi Nb} \end{aligned}$$

- Q_c only depends on the material!



OSNS10
September 2007
ISIS

18

Basic Ideas


■ Neutron Potential

$$\begin{aligned} V &= V_{Nuclear} + V_{Magnetic} \\ &= \frac{\hbar^2}{2m_n} Nb + \mathbf{B} \cdot \hat{\mathbf{s}} \end{aligned}$$

$$F_j = \frac{k_{j+1} - k_j}{k_{j+1} + k_j}$$

$$r_j = \exp(-ik_j d_j) \times \frac{r_{j+1} + F_j}{r_{j+1} F_j + 1}$$

$$R = |r|^2$$



■ Kinematical Approximation

$$R(Q) = 16 \frac{\pi^2}{Q^4} \left| \int \rho(z) \exp(iq/s) z dz \right|^2$$

OSNS10
September 2007
ISIS

19

■ QM solution

$$-\frac{\hbar^2}{2m_n} \nabla^2 \Psi(\vec{r}) + V(\vec{r}) \Psi(\vec{r}) = E \Psi(\vec{r}) \quad (1)$$

$$k_m = \sqrt{k_z^2 + \Gamma_0 - \Gamma_z} \quad (2)$$

$$E = \frac{\hbar^2}{2m_n} |\vec{k}|^2 + V_0 = \frac{\hbar^2}{2m_n} (k_x^2 + k_y^2 + k_z^2) + V_0 \quad (2)$$

$$k_z^2 + \Gamma_0 - \Gamma(z) = 0 \quad (3)$$

$$V(\vec{r}) = \frac{2\pi\hbar^2}{m_n} N(\vec{r}) b(\vec{r}) \quad (3)$$

$$\text{for } -\infty < z \leq 0$$

$$\Gamma(\vec{r}) = 4\pi N(\vec{r}) b(\vec{r}) = \frac{2m_n V(\vec{r})}{\hbar^2} \quad (4)$$

$$\psi(z) = \exp(ik_m z) + r \exp(-ik_m z) \quad (5)$$

$$\nabla^2 \Psi(\vec{r}) + (k^2 + \Gamma_0 - \Gamma(\vec{r})) \Psi(\vec{r}) = 0 \quad (5)$$

$$R = rr^* \quad (6)$$

$$\frac{\partial^2 \psi(z)}{\partial z^2} + (k^2 + \Gamma_0 - \Gamma(z)) \psi(z) = 0 \quad (6)$$

$$R(q_i) = \frac{|q_i - q_{i+1}|^2}{|q_i + q_{i+1}|^2} \quad (7)$$




OSNS10
September 2007
ISIS

20

Using the neutron's spin

- The neutron is a spin $\frac{1}{2}$ particle
- The neutron possesses an intrinsic magnetic moment: spin
- Caution...




Caveat to Classical Description


- CD predicts a continuous variation of critical edge

$$\frac{4\pi \sin(\alpha_c^\pm)}{\lambda} = Q_c^\pm = \sqrt{\frac{2m_n}{\hbar^2}} (V_n \pm |\mu| |B_s| \cos(\theta))$$


- Stern-Gerlach effect? Only 2 eigenstates



PNR from a single layer




PNR from a multiple layers



Spin dependent cross-section

- In-plane orientation of magnetisation obtainable from 4 spin dependent cross-sections



- Components of the magnetisation, m give rise to

$m \parallel H$: Non Spin Flip Scattering (NSF)

$m \perp H$: Spin Flip Scattering (SF)

- Dynamical analysis gives absolute depth dependence profile

$$b = b + p \sin \phi$$

$$p_m \cos \phi = p_x$$

$$\left[\frac{-\hbar^2}{2m_n} \nabla^2 + V(r) \right] \psi^{\uparrow,\downarrow} = E \psi^{\uparrow,\downarrow}$$





OSNS10
September 2007

25

Reflectometers

- The CRISP reflectometer at ISIS is a white beam time of flight (tof) polarised neutron reflectometer viewing a 20K hydrogen moderator.




<http://www.ill.fr/YellowBook/D17/>



<http://www.ill.fr/YellowBook/ADAM/>

- Continuous or tof mode of operation
- 5Å Monochromator (6Å Polarised)
- White beam flux 9.6×10^9 n/s/cm²



OSNS10
September 2007

26

Simulation Packages



Simulation Packages: neutron

Neutron

Polly

http://www.isis.rl.ac.uk/LargeScale/CRISP/da_ta_analysis_software.htm



OSNS10
September 2007

28

Simulation Packages: photon

X-ray

- <http://sergey.gmca.aps.anl.gov/>
- <http://solst.phys.msu.su/andreeva>



29


Depth dependent vector magnetometry

Interlayer Exchange Coupling




Spintronics

- Spintronics offer:
 - Nonvolatile, high speed, low power, scaleable
 - Reconfigurable
 - Integrated MO devices
 - Quantum information
- MgO tunnel barriers
- Spin Accumulation
 - δ -doping
 - Forthcoming experiments at SRS, NSLS
- Spin-torque
 - Large area arrays
 - e-beam
- Precessional pumping



31

The GMR Effect



Baibich *et al.* Phys. Rev. Lett. **61** 2472 (1988)


- Applications
 - Computer read/write heads
 - ABS sensors
 - Opto-isolators
 - MRAM




32

Quantum Well State

- Band matching important e.g.




Co/Cu GMR



OSNS10
September 2007
ISIS

33

Interlayer Coupling




<http://www.dspc.vt.edu>

Thin Fe Overlayer

Au Spacer Layer

Cu Whisker Substrate



J. Unguris, et al. J. Appl. Phys. 75, 6437 (1994).

OSNS10
September 2007
ISIS

34


$$\frac{\mathbf{E}}{\mathbf{A}} = -J\hat{\mathbf{m}}_1 \cdot \hat{\mathbf{m}}_2 \quad (1)$$

$$\frac{\mathbf{E}}{\mathbf{A}} = -J_2(\hat{\mathbf{m}}_1 \cdot \hat{\mathbf{m}}_2)^2 \quad (2)$$

M Stiles (2002)

Spin valve structures

Si/Cu(494Å)/FeNi(71Å)/Cu(31Å)/Co(46Å)/Cu(31Å)/FeNi(74Å)/Cu(46Å)



FeNi



Co

FeNi

600G

- Saturation field all moments align along applied external field

J.A.C Bland et al. Phys. Rev B 60 (1999) 7304




OSNS10
September 2007
ISIS



35

20G applied field

- In 20G the Co is able to rotate by 60°
- PNR gives absolute moments with layer sensitivity




20G



OSNS10
September 2007
ISIS



36

GMR and MOKE




$$\frac{\Delta\rho}{\rho} \Rightarrow -\left(\frac{M_0}{M_s}\right)^2$$

- Room temperature measurements
- Reduction in gmr and zero field remanence
- Biquadratic coupling?




$$\approx \frac{\sqrt{2}M}{2}$$



OSNS10
September 2007
ISIS

37

Clean AF coupled sample




- Small spin dependence
- Large gmr
- Ferromagnetic contribution at bottom of stack

OSNS10
September 2007
ISIS

38

Biquadratic sample



- Damaged sample
- Spin dependence
- Reduced gmr
- Direct observation of 90° coupling
- Single domain





■ Marrows *et al.* Physical Review B 62 11340 (2000)

OSNS10
September 2007
ISIS

39

Front-back reflectivity

- Ni₈₀Fe₂₀(50)/Fe₅₅Pt₄₅(20nm)
- Exchange Spring magnet



■ K.V. O'Donovan *et al.* Phys. Rev. Lett. 88 (2002) 067201

OSNS10
September 2007
ISIS

40

Proximity Magnetism

Interfacial magnetism




Roughness

Structural and magnetic interfacial phenomena




Surpressed Magnetisation through PNR/EELS

- $\text{La}_{0.7}\text{Ca}_{0.3}\text{MnO}_3/\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ superlattices
- Proximity effects
 - Variation in superconducting transition temperature
 - Critical current
- Half-metallicity
- Charge transfer key




Hoffmann *et al.* Phys. Rev. B 72 140407 (R) (2005)





OSNS10
September 2007

42

Structural Roughness



$$R(Q) = R_F \exp(-Q^2 \sigma^2)$$





44

Off-specular scattering

Probing in-plane lengthscales



Diffuse scattering




Holy and Baumbach, prb, 49, 10668, (1994)



47

Experimental Geometry



PHYSICAL REVIEW B

VOLUME 38, NUMBER 4


1 AUGUST 1988

X-ray and neutron scattering from rough surfaces

S. K. Sinha, E. B. Sirota, and S. Garoff*
Corporate Research Science Laboratory, Exxon Research and Engineering Company, Clinton Township, Route 22 East, Annandale, New Jersey 08801

H. B. Stanley†
University of Maryland, College Park, Maryland 20742
(Received 30 November 1987)

$$C(R) = \langle [z(x, y) - z(x', y')]^2 \rangle = \sigma^2 \exp(-r/\xi)^{2h}$$




- σ = roughness
- ξ = cut-off length:
 - for $R > \xi$, interface appears smooth,
 - for $R < \xi$, interface appears rough, fractal behaviour
- h = 3-D Hurst parameter for jaggedness ($0 < h < 1$)
 - smooth: $D=2$, $h=1$
 - very rough $D=3$, $h=0$

OSNS10 September 2007



48


Effects of magnetisation on neutron reflectivity



Magnetic Roughness



Magnetic Surfaces



Modelling of Data

- Consider spatially inhomogeneous magnetisation profile: $m(r)$

$$m(\vec{r}) = m_0(\cos(\phi(\vec{r})), \sin(\phi(\vec{r})), 0)$$

- Introduce a correlation function:

$$M(\vec{r}) = \sigma_m^2 \exp(-r/\xi_m)$$

- Within the Born approximation:

$$S(\vec{Q}) \propto \sum_{\alpha\beta} \int d^3r \exp(i\vec{Q} \cdot \vec{r}) (\delta_{\alpha\beta} - Q_\alpha Q_\beta) \langle m_\alpha(\vec{r}) m_\beta(0) \rangle$$


- Averaging for a Gaussian distribution of orientations:

$$\begin{aligned} S(\vec{Q}) &= S_{spec} + S_{diff} \\ &= 4\pi^2 D \delta(Q_{||}) + D \int d^2r \exp(i\vec{Q}_{||} \cdot \vec{r}) [\mathbf{s} + \mathbf{m} + \mathbf{sm}] \end{aligned}$$




Diffuse scattering

- Coherent AF order
- No evidence for uncorrelated roughness



μ -magnetics


- Solve LLG equation



Probe coherence: *Consideration in scattering experiments*

Probe Coherence (Longitudinal + transverse)

$$\begin{aligned} 2L_{Coh} &= N\lambda \\ &= (N+1)(\lambda - \delta\lambda) \\ N &\approx \frac{\lambda}{\Delta\lambda} \\ L_{Coh} &= \frac{\lambda^2}{2\Delta\lambda} \end{aligned}$$




- Similarly for transverse (d is the slit width, l the distance)

$$L_{Trans} = \frac{\lambda}{2} \left(\frac{l}{d} \right)$$

$L_{\text{Coh}} < \text{Domain size}$

- $I \propto \sum_i (A_i^2)$
- Only specular scattering



OSNS10
September 2007




Science & Technology Facilities Council
ISIS

57

$L_{\text{Coh}} > \text{Domain size}$

- $I \propto (\sum_i A_i)^2$
- Only diffuse scattering



OSNS10
September 2007



Science & Technology Facilities Council
ISIS

58


Domains and the coherence volume

Many thanks to H. Zabel for slides




Science & Technology Facilities Council
ISIS

Simulation of a Co film on a substrate



70 nm Co
Si



Magnetic parameters

$$\langle \cos \gamma \rangle = 0.7$$
$$\langle \sin^2 \gamma \rangle = 0.25$$
$$\langle \cos \Delta\gamma^2 \rangle, \langle \sin^2 \Delta\gamma^2 \rangle$$
$$L_c \approx 100 \mu\text{m}$$


OSNS10
September 2007




Science & Technology Facilities Council
ISIS

60


Domain size: $100 \mu\text{m}$ ($> L_c$)




Domain size: $100 \mu\text{m}$




Domain size: $50 \mu\text{m}$




Domain size: $10 \mu\text{m}$




Domain size: 5 μm



Domain size: 1 μm




Single ferromagnetic film in the domain state




67

Banana shape off-specular scattering from domain state



ISIS
OSNS10
September 2007
68

Simulation of domain state



Complementarity of techniques

- Photon**
 - Element specific
 - Band selective
 - Large resonant enhancements
 - Strong absorption (resonant)
 - Polarisation dependence
 - Small coherence length (<10μm)
- Neutron**
 - Large, absolute cross-section
 - Low absorption
 - Polarisation dependence
 - Long coherence length (~100μm)

OSNS10 September 2007
ISIS 70

Nanomagnetism II:

Applications of the nanomagnetism tools


Sean Langridge
Large Scale Structures Group
Rutherford Appleton Laboratory, Chilton, Oxfordshire

s.langridge@rl.ac.uk

OSNS10 September 2007
ISIS 71

Exchange bias

Shifted hysteresis loop away from zero field due to exchange coupling with the last af layer




Meiklejohn and Bean Phys. Rev. 105 904 (1957)

- Bias field orders of magnitude too large
- $H_E = 2JS^2/(a^2 Mt)$
- No coercivity enhancement
- Temperature dependence
- Compensated interfaces


OSNS10 September 2007
ISIS 72

Domain Wall Structure

- Parallel Domain wall
 - Mauri *et al.* *J. Appl. Phys.* **62**, 3047 (1987)
 - Exchange energy spread into the AF layer




- Perpendicular Domain wall
 - A.P. Malozemoff *Phys. Rev. B* **35**, 3679 (1988)
 - Random field from defects breaks AF into domains



OSNS10
September 2007
ISIS

73

As grown sample




- 150 Å bilayer spacing
- Broadening of Bragg peak
- loss of vertical coherence

OSNS10
September 2007
ISIS

74


$T >$ Blocking Temperature



OSNS10
September 2007
ISIS

75

Hysteresis Measurements




- Field Cooled
- Unpolarised
- Uncorrelated disorder


OSNS10
September 2007
ISIS

76

Field Cooled



- Asymmetric
- Small domain size
- Domain size reduces from $1.5 \mu\text{m}$ to $1 \mu\text{m}$ upon reversal
- $\varepsilon_{fc} < \varepsilon_{t>T_b}$




OSNS10
September 2007
ISIS

77

Artificial Exchange bias: *double superlattices*

Science & Technology Facilities Council
ISIS

Magnetisation





- Good AF (0.25 mJ/m^2) and F coupling in respective components

OSNS10
September 2007
ISIS

79

Spin-flop phase - I


$\blacksquare H=600\text{Oe}$



OSNS10
September 2007
ISIS

80

Exchange Spring phase - II





OSNS10
September 2007
ISIS

81

Energy Minimisation

- Minimise the free energy per unit area
- $$\epsilon = -\sum \mu_0 m H \cos(\theta_i) - J_i \cos(\theta_i - \theta_{i+1})$$

- Monte Carlo algorithm



OSNS10
September 2007
ISIS


82

Patterning of magnetic systems

A route to higher recording densities...

Science & Technology Facilities Council
ISIS

Moore's Law for Media




OSNS10
September 2007
ISIS


84

xrmSAXS

- CoCrPtB
- Amorphous grain boundaries



$$fE1 = [(\vec{e} \cdot \vec{e}) F^0 - i(\vec{e} \times \vec{e}) \cdot \hat{z}_n F^1 + (\vec{e} \cdot \hat{z}_n)(\vec{e} \times \hat{z}_n) F^2]$$



- Expect to observe struc-struc and struc-mag correlations


E.E. Fullerton *et al.*, Nucl. Instr. and Meth. in Phys. Res. B 200 (2003) 202–209

OSNS10
September 2007
ISIS


85

Stamping magnetic structures

- Micro-stamp production
- 200/200nm stripes
- 20nm high



IBM Zurich



A. Kumar and G. M. Whitesides, Appl. Phys. Lett. 63, 2002 (1993).

OSNS10
September 2007
ISIS

86

Co L_{III} resonance






FIG. 1. A scanning electron micrograph of the sample. Inset is a Kerr loop of the sample with the field applied in the plane, both along (+) and across (○) the direction of the lines.





OSNS10
September 2007
ISIS

87

Complementarity: real and reciprocal space

- Combination of real and reciprocal techniques
 - Kerr Microscopy
 - PNR-pa
- CoFe 3 μm periodicity
- Ripple domain structure
- dwba analysis
 - S. K. Sinha, E. B. Sirota, S. Garofalo, and H. B. Stanley, Phys. Rev. B **38**, 2297 (1988)
 - B. P. Toperberg, Physica B **297**, 160 (2001); Appl. Phys. A: Mater. Sci. Process. **74**, S1560 (2002)



Theis-Brohl *et al.* Phys. Rev. B. 020403(R) (2005)

OSNS10
September 2007
ISIS

88

SPM


- Large sample area
- MFM
- Capacitance
- Liquid Surface adapter
- Torsion Mode



OSNS10
September 2007
ISIS

Science & Technology Facilities Council

89



13.5 micron




AFM
MFM



Controlled Surface Morphology

- Self Assembly of $0.2\mu\text{m}$ latex spheres
- $[\text{Co}(2\text{nm})\text{Ru}(1.8\text{nm})]_{50}$

V. Ng, Y. V. Lee, B. T. Chen, and A. O. Adeyeye, *Nanotechnology* **13**, 554 (2002).




OSNS10
September 2007
ISIS

90

Exchange Coupling Strength

- Separation of bilinear/biquadratic
 - Fullerton & Bader Phys. Rev. B. 53, 5112 (1996)
 - $\epsilon \approx \mu_0 H_{\text{ext}} (\cos \theta_1 + \cos \theta_2) - J_1 \cos \theta - J_2 \cos^2 \theta$
 - $\frac{\mu_0 H_{\text{ext}}}{4} = J_1 - 2J_2 \cos \theta + 4J_2 \cos^3 \theta$






Clean	Patterned
$J_1 = -0.62 \text{ mJm}^{-1}$	-0.58 mJm^{-1}
$J_2 = -0.05 \text{ mJm}^{-1}$	-0.07 mJm^{-1}

OSNS10
September 2007
ISIS

91

XTEM

- **320Kx**




OSNS10
September 2007
ISIS

92

XTEM

910Kx

- The brighter lines are the Ru and the darker the Co



- Layer thicknesses in good agreement with nominal structure



OSNS10
September 2007
ISIS

93


Patterned System

$$\theta = \frac{Q_x}{2Q_z} - \frac{Q_x}{2Q_z}$$

Coercive Field



AFM FM




OSNS10
September 2007
ISIS

94

Saturation Field

- Near-Saturation field



- Magnetic Roughness (cross-correlation of structure and magnetism)
 - Diffuse scattering transferred from AFM to FM Bragg peak
 - $\xi_M < 2000 \text{ \AA}$



OSNS10
September 2007
ISIS

95

μ -magnetics

- Simulation reveals rich phase diagram

- Quantitative agreement
- AFM domain size
- Degree of disorder
- Qualitative agreement with correlation length



OSNS10
September 2007
ISIS

96

Superconductivity:


flux penetration and proximity effects

OSNS10
September 2007



97

Flux penetration in superconductors



Flux penetration in superconductors





FIG. 2. The spin-dependent reflectivities R^+ and R^- measured in an applied magnetic field of 6.0×10^4 A/m (750 Oe). The continuous lines are the reflectivities calculated for the two polarization states, with the same instrumental and surface parameters as in Fig. 1, and an exponential decay of magnetic induction with a penetration depth of 39 nm.



- 1 μm Pb film
- PbO surface layer
- $B(z) = \mu_0 H \exp(-z/\lambda)$
- $\lambda = 39 \pm 1$ nm
- No field dependence

PHYSICAL REVIEW B

VOLUME 49, NUMBER 22

1 JUNE 1994 II

Magnetic-inductance profile in a type-I superconductor by polarized-neutron reflectometry

M. P. Nistley*

Institut Laue-Langevin, 3600 Grenoble CEDEX, France

A. T. Boothroyd and C. R. Stodols

Department of Physics, Clarendon Laboratory, University of Oxford, Parks Road, Oxford, OX1 3PU, United Kingdom

D. M. Peacock

Department of Physics, University of York, University, CV1 5FL, United Kingdom

J. Penfold


ISIS Science Division, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

(Received 17 February 1994)

Science & Technology Facilities Council
ISIS

99

Flux distributions in s/c samples



Summary

■ Summary

- Scattering techniques are ideally suited to a wide range of topical problems in Nanomagnetism research across the magnetic periodic table



- Ability to characterise μg to pico-grammes of magnetic material



■ Acknowledgements

- Hours on the beamline with...




OSNS10
September 2007
ISIS
Science & Technology Facilities Council

101

Acknowledgments

- L.A. Michez, *C.H. Marrows*, B.J. Hickey
- P. Steadman
- C. Kinane*
- Rob Dalglish, *Tim Charlton*
- H. Zabel
- D.G. Bucknall, H-L. Zhang
- J. Dvorak, D. Arena
- R. Springell, G.H. Lander, W.G. Stirling
- EPSRC funded research:



OSNS10
September 2007
ISIS
Science & Technology Facilities Council

102

