Daubenmire Datasheet Site: Date: Transect: Observer: Round: Recorder:

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf	Litter
0 m	29	32	9				Tamer #	Opecies	Ground	Litter	Depth (cm)
10 m	100	0	89	0					0	98	
20 m	0	0	2	3					0	9	
30 m	(0	2	0					9	8	
40 m	300	4	89	0				8	0	30	<u> </u>
50 m	8	0	88	2				5	\bigcirc	B	20
60 m	S	0	M	2				0	0	50	\rightarrow
70 m	2	53	M	0				2	0	30	H
30 m	90	Š		0	200			0	0	80	7
90 m	199	88	~					0		B	7

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	d	6	\sim	6	M	4	2	d	00 111	→ m
E	9	d	~	d	M	-	Cl	d		3
s	CP	d	5	4	M	d	~	4		4
W	d	C	d	a	N		7	1		1

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100% CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm

prd2

tprd2a

3

100m