Daubenmire Datasheet

Date:

Observer: Recorder:

Team lead:

Site: prd1

tprd1a

Round: Transect: 100m ω

7.	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody	Bare	Leaf Litter	Litter Depth (cm)
0 m	0	63	88	0				0	6	$\sim$	
10 m	<b>○</b>	63	3/6	0				0	16	5	- S
20 m	0	0/8	80	0				5	_	200	0
30 m	0	NE	88	0				>	6	(2)	0,25
40 m		80	10	Ø				) (	10	Q (3)	0,25
50 m	0	278	O	0	1			Ot	0)	360	0.25
60 m	7	20	Q	-	_			0	W	0 128	0,25
70 m	0	278		0				0		3	0,25
80 m	10	1 (0	fle	0	1			0		8	, s
90 m	N Q	38	02	0				Ö	0	2/0	20
Dobal Pola	Dollo.									9	

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	70 m 80 m
z	5	7	A	Q	//0	16	7	0	8
Е	$\infty$	9	03	٥	13	160	160	7-1	7 4
S	6	ij	8	12	16	6	7	6	16: 10
₹	S	7	C.	5	14	0)/	7	E	E

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Litter Depth - to nearest .25 cm Leaf Litter (horizontal dead material)

Appelgate updated 5/30/17