Daubenmire Datasheet

Observer: Date:

Recorder:

Team lead:

Site: Transect: tgro1a gro1 2

Round: 60m

-	2		7	38	22	W,	0)2	53	(N	CS6
	<			0	0	0	0	0	0	WS6
		7	5	80		8(0	38	S	2	Forbs
	<		1	0	0	-6	0	0	0	Milkweed
		7	A	9	0-	_	0	0	0	Common ramet #
			7	0	0	0	0	C	0	Swamp ramet #
				0	0	0	0	Q	Ò	Butterfly ramet #
			1	0	0	0	0	0	0	Woody Species
			\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	38	6	160	250	W	S S	Bare Ground
	5		2	63	86	20	80	80	W R	Leaf Litter
				0 25	0.25	0.25	0.25	0, 25	0.25	Litter Depth (cm)

Robel Pole:

₩ 06

60 m

50 m 40 m 30 m 20 m 10 m o ₹

	z	Е	S	≷
0 a	7	Q	W	7
10 m	S	(0)	(6)	X
20 m	6	(0	(0)	0
30 m	1 10	-	9	
40 m	.33 X	A	(N.	W
50 m	4	N	0	6
60 m		7		
70 m	A Section of the Sect		X	
80 m	The second secon		V	
90 n	V		The same of the sa	1

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm