Daub...mire Datasheet

Date:

Recorder:

Team lead:

Observer:

Site: Transect: tnor1b nor1

50m

Round:

	Swamp Butterfly Woody amet # ramet # Species	Butterfly ramet #	CSG WSG Forbs Milkweed ramet # r	0 m 10 0 98 0 0	10 m 10 0 98 0 0	20 m /0 0 98 0 0	30 m 10 0 80 0 0		40 m /2 0 98 0 0	76	16 0	76 0	0
Manager 1		Woody Species	Milkweed   Common   Swamp   ramet #	0	98 0 0	78 0 0	C 0 0 0%	))	0	0	0 0		0 0
	Leaf Litter		Litter Depth (cm)	0.20	0,25		5,0	1.0	5.0	5.0	0,5	0,5	0,5

	т О ж	10 m	20 m	30 m	40 m	50 m	786	60 m	60 m 70 m
z	JW	100	0)]	12	1/6				
ш	7	14	10	Ñ	) [i				
S	7	(6)	16	元	71				
*	5	0) [	. 16	2	100				

CSG - cool season grasses (including sedges) Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm