Daubenmire Datasheet

Date:

Observer:

Recorder:

Team lead:

Site:

pio2

Transect:

tpio2a

Round:

63 - 50%-75%

2

80m

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	98	Ø	Ø	Ø	Ø,	8_	Ø	Ø	Ø,	86	0.50
10 m	98	8	Ø	Ø	8	10	8	Ø	10	80	2.00
20 m	78	Ø	3	Ø,	Ø	Ø	Ø	Ø,	Ø	80	200
30 m	63	8	38	10	0	8	Ø	8	100	86	5.00
40 m	80	Ø	10	Ø	8	8	8	Ø	8	80	0.75
50 m	86	Ø	1(2	8	Ø	Ø	8	Ø	· Ø	86	0.75
60 m	63	8	38	8	Ø	0	2	0	8	86	0.75
70 m	63	Ø	38	3	\	Ø	Ø	8	8	80	3.00
80 m	1 1 1	$\wedge \wedge \wedge$	001	\triangle	1 1 1	ΛΛ	AAA	AAA	AAA	AMA	MA
90 m	VVV		VVV	\cup \vee \vee	VVV	VVV	1000		V V V	VVV	VVV

Robel Pole:

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m 90 m
N	5	5	\	2	10	3	Q	8	
Ε	4	5	7	10	7	3	8	10	
s	3	0	7	12	10	3	8	10	
W	3	5	F	10	12	5	(0	10	Manual Spirite Manual

senescing destroy season

senescing destroy season

86 - 75%-95% | 98 - 95%-100% person

Solid understory,

Thick ro red clover

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm