50m

Daubenmire Data Sheet

Sampling Date and Year:

Site: Caylord Jones, Bioreactor Webster City

ののちち

2238

Data Collected by: KELDEWS Haw + ROYCH on this date: 4/27/16

Wild grope, smell

	952	WSG	Forbs	Milkweed		Woody	Bare	Leaf	Litter	
			-			Species	Ground	Litter	Depth	
1	38	0	3	0		2	91	92	1.5cm	
2	91	0	38	0		0	35	38	2.5 cm	
8	38	0	9%	0		0	9	98	1 cm	
4	16	0	38	0		9	2	63	SUM	
2	9	0	38	0		3	9/	26	1cm	
9	10	0	63	0		1 6	16	860	SPA	
1						)			,	
<b>∞</b>										
6										
10										

0
0
4
_
0
9
0

N 4 5 6 7 8 9 10  E 3 4 5 6 7 8 9 10  W 5 9 10 6 4 9	-		The second second second second second	The second secon			•				
4 6 8 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7		1	2	3	4	5	-0	7	œ	6	10
2	Z	7	0	2	3	h	rt				
7 9 01 0 01 1	П	8	>	+	~	7	+				
9 01 6	S	+	9	8	10	77	+				
	>	N	6	01	9	h	5				

50m transfect at Bioreoceters

<u>38</u> - 25%-50% <u>63</u> - 50%-75% <u>86</u> - 75%-95% <u>98</u> - 95%-100% 16 - 5%-25% 3 - trace-5% CSG - cool season grasses (including sedges) Categories:  $\underline{0}$  - absent  $\underline{1}$  - trace

WSG - warm season grasses

Forbs - forbs not including milkweed

Milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material), Litter Depth - to nearest .25 cm

jon1 tjon1a 1