Daubenmire Datasheet

Date: 7/2/9

Observer: Recorder:

Team lead:

Site: byd1

Transect: tbyd1a

2

Round: 100m

•	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	58	8	Re	9	0	2	0	DY	38		
10 m	1 Ce	8	86	Ø	X	X	(X)		1/	63	0.50
20 m	38	(X)	103	8	2	X	(2)		10	86	0.50
30 m	160	Ø	Ca	X	AV	~			10	80	0.50
40 m	20	20	0		20		0.		38	(03	0.50
	26	19	103	X	\varnothing	8	8	X	100	No	0.50
50 m	43	Ø	38	8	0	M	00	8	35/	CaZ	A =
60 m	100	8	\$ 8G	8	0				08	US	0.00
70 m	28	Ø	102		3			20	63	38	0.50
80 m	162	OX -	107	XJ N				9	3	878	0.50
	40	X)		· 1	\mathcal{A}		0	0	10	810	0.50
90 m	112	\varnothing	910			2)	9	X	30	(03	0.50

0 m 10 m 20 m 30 m 40 m 50 m 60 m 70 m 80 m 90 m N Ε 2 13 S 2 W

Categories: <u>O</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100% CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm