Daub ... mire Datasheet

Date: 7-17-18

Observer: Mail

Team lead:_

Site: app1
Transect: tapp1a
2
Round: 5000

50m

	CS6	0 m (13	10 m U 3	20 m 80	30 m 38	40 m (23	50 m	50 m	50 m 70 m	50 m 70 m
	Forbs	28	38	160	38	W				
Forbs	Milkweed	0	0	0	0/	O				
	Common ramet #	0	0	0)	0	O ₄				N 2000 200 1000
Milkweed O	Swamp ramet #	\Diamond	0	0	0	0				
Milkweed Common ramet #	Butterfly ramet #	0	0	0	0	0				
Milkweed Common Swamp ramet # ramet #	Woody Species	0	0	0	0	0				
Milkweed Common Swamp Butterfly ramet # ramet # ramet #	Bare	_	10	W	16	16				
Milkweed Common Swamp Butterfly Woody ramet # ramet # species	Leaf Litter	X	63	978	63	26				
Milkweed Common Swamp Butterfly Woody Bare ramet # ramet # Species Ground	Litter Depth (cm)	0.25	0.25	0.5	2,5	Ó				

Robel Pole:

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
z	12	2	0	12	14					
m	14	14	0	80	8					
S	12	8	12	10						
8	N	>°	J	72	12					

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm