50 m 40 m 30 m 20 m 10 m 70 m 60 m 0 3 Date: Recorder: 5 Observer: 5 Dauberimire Datasheet Team lead: ≤ CSG DAZO WS6 5011 Forbs Milkweed ramet # Common ramet # Swamp ramet # Butterfly Site: Species Woody Round: Transect: Ground Bare tuth2a 100m uth2 ω Litter Leaf 0 Depth (cm) Litter

0 m 10 m 20 m 30 m 40 m 50 m 60 m 70 m 80 m 90 m H H 10 5 9 10 9 11 10 12 H 17 17 18 80 m 90 m		S	E /	Z	0	100001 . 010.
20 m 30 m 40 m 50 m 60 m 70 m 80 m 10	4	8	N	7		
30 m 40 m 50 m 60 m 70 m 80 m	0	٠١. دا ،	4	X	10 m	
40 m 50 m 60 m 70 m 80 m	7	0	12	10	20 m	
50 m 60 m 70 m 80 m		7		5	30 m	
60 m 70 m 80 m	<u></u>	2	χ.	2	40 m	
70 m 80 m	0/0		12	0	50 m	
80 m	Ò	//	F	9	60 m	
		N	6		70 m	
90 m	2	2	41	0	80 m	
	15	St	£	7	90 m	

90 m

CSG - cool season grasses (including sedges) Categories: O - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm