Date: Daubenmire Datasheet Recorder: Observer: Site: Transect: Round: tjon1a 50m jon1 2

Team lead:

	CS6	WS6	Forbs	Milkweed	Common #	Swamp ramet #	Buttertly ramet #	Species	Ground	Litter	Depth (cm)
5	3	2	XIO.	3	0	9	0	0	16	6	- 5
10 1	C	0	200	3	\bigcirc	0	0	0	38	(2.5)	0,5
20 5)(0	20) (7)	0	N	1(0	63	0,5
20 =	20	36	2) K) (0	0		0	0.25
M OC	()(- 0) () ()(*	X	1	17,	いしい
40 m	0	CN	28	C	C	(C	(00		0.00
50 m	3	5					1	-			1
60 m		> >	>						5		>
70 m								and the same of th			
80 m									<	5	
90 m		0									

	0 #	10 m	20 m	30 m	40 m	50 m	60 m	A A	70 m	1 / 1
z	7	4	4	工	Q		V		1	
m	4	Jr-	_	N	×	1	V		CONTROL OF THE PARTY OF THE PAR	
S	4	6	(6)	<u>_</u>	Q	\setminus	Λ	1	1	And Part Comments
€	4	T	2)	0	5	M	1		Agenta membership	Control of the c

Categories: $\underline{0}$ - absent | $\underline{1}$ - trace | $\underline{3}$ - trace-5% | $\underline{16}$ - 5%-25% | $\underline{38}$ - 25%-50% | CSG - cool season grasses (including sedges) 00 **6-100%**

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm