Daubenmire Datasheet

Date:

Observer:

Recorder:

Team lead:

Site:

tnie1a

50m

Round: Transect: nie1

90 m	80 m	70 m	60 m	50 m	40 m	30 m	20 m	10 m	0 m	
		a good from part of		>	W	860	_	Ü	U	CS6
C		e de servición de la companya de la	- may a T	3	38	B	63	38	160	WS6
<	and the second	STATE OF THE STATE	and the second	5	NX	38	16	600	22	Forbs
		and the second	and the state of	>	W	Ø	33	Ø	Ø	Milkweed
		ALLOWER TOP	vaniture service	3	_	Ø	W.	Ø	Ø	Common ramet #
		stilet.			Ø	Ø	Q	Ø	B	Swamp ramet #
		and the base		1	Ø	Ø	Q	B	9	Butterfly ramet #
	A Commence of the Commence of	A COLUMN TO SERVICE AND ASSESSMENT OF THE PERSON OF THE PE		3	B	8	Q	Ø	Ø	Woody Species
		en an est est est	and the second	>	W	W	W	W	S	Bare Ground
	Control of the Contro	The state of the s	>	1	20	308	86	0)8	0)8	Leaf Litter
			}	7	0.25	0.50	0.25	0,75	0.50	Litter Depth (cm)

ι	3	•
ġ	ō	
-	U	,
9	<u>2</u>	
	0	

	z	m	S	€
0 m	4	4	W	W
10 m	S	4	W	4
20 m	7	W	W	2
30 m	U	6	6	S
40 m	7	X	S	S
50 m	1	V	~	1
60 m	\	\[		1
70 m	A	1	1	$\mathbb{N}$
80 m		V		V
90 m	1	1	Tonas and the same of the same	

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm