Daubenmire Datasheet

Date:

Observer:

Recorder:

Team lead:

Site:

nkn2

Transect:

tnkn2a

Round:

1

100m

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf	Litter
0 m	38	3	300	7)	6	•	0	Opecies	Ground	Litter	Depth (cm)
10 m	112	3	103	1	6	0	6		(0)	38	0.25
20 m	)(0	1	20	0	0	6	2	0	63	16	0.25
30 m	IÙ	1	(03	7		0	()	Q	86	3	0.25
40 m	16	10	38	8	0	0	O		38	38	0.25
50 m	16	3	43	6		8	- 6		(13)	10	0.25
60 m	10	3	38	0	0	6	0		03	16	0.25
70 m	16		38		0	A	0	5	03	10	0.25
80 m	14	3	1(0			No.	0		10	(0.5	0.25
90 m	16		63	0	0	0	0		10	38	0.25
Robel	Pole:				7	U	(()		38	10	0.25

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	2	2	2	2	2	0	A	1	) III	90 m
Ε	2	2	2	2	3		2	A		1
S	2	2	2	4	)	0	3	2	1	1
W	2	2	2	4	3	5	)		2	2

Categories: 0 - absent 1 - trace 3 - trace-5% 16 - 5%-25% 38 - 25%-50% 63 - 50%-75% 86 - 75%-95% 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm