Daubenmire Datasheet

Date: 8

Observer: Wanty

Team lead: \_

Site: bcr1 tbcr1a

Transect:

3

Round:

50m

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
O m	38	0	38	0	_		/	0	38	16	0,25
10 m	10	Ð	38	Ð				0	10	10	0, 25
20 m	1(0	<del></del>	63	0	-			00	10	38	0,5
30 m	3	0	80	0				-0	110	38	0 0
40 m	3	38	63	0				0	5	10	0,5
50 m											
60 m					4						
70 m											
80 m											
90 m											

Robel Pole:

LODGI I GIG.										
	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	12	9	16	10	10					
Е	10		16	16	8					
s	15	18	16	14	8					
W	13	14	16	160	14					

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm