Daubenmire Datasheet

Date:

Team lead:

Recorder: Observer:

> Site: Transect:

tnkn1a

nkn1

Round:

100m 1

	CS6	WS6	Forbs	Milkweed	Common	Swamp ramet #	Butterfly ramet #	Woody	Bare	Leaf	Litter Depth (cm)
0 m	38	0	8	0				0	0X	2	>
10 m	300	0	(°2	0				a	62	5	0
20 m	200	0	8					0	200	00	2
30 m	000	0	W					0	5	2	· C
40 m	(3)	0	200	<u>ک</u>					5	33	20
50 m	80	0	6	0				0	200	M	2
60 m	38	9	S	0				0	53	5	9
70 m	38	0	38	0				0	S	5	0
80 m	6	0	38	0				0	8 S	6	2
90 m	80	0	6	0				O _r		.;. W	No.
Dobal Bola	Bala:										-

	0	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m
z	P	7	2	6	0	2	2	7	\mathcal{L}
m	V		2	_	78	6	00	7	
S	2	0	2	2	0	O)	0	9	10
€	7	2	57			5	5	0.0	=

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm