Daubenmire Datasheet

Date: 81410

Observer: Kindy

Recorder:

Team lead:

Site:	byd1	
Transect:	tbyd1a	
Round:	3	
Round.	100m	

	CSG WSG Sombo Millowed Common Swamp Butterfly Woody Bo							Bare	Leaf	Litter	
	CSG	WSG	Forbs	Milkweed	ramet #	ramet #	ramet #	Species	Ground	Litter	Depth (cm)
0 m	3	Ø	98	Ø	Ø.	Ø,	Ø	Ø.	63	38	0.75
10 m		63	38	18	Ø	Ø	Ø	Ø	03	38	0.25
20 m	Ø	38	63	Ø	Ø	Ø	Ø	Ø	63	38	0.25
30 m	3	10	86	Ø	Ø	Ø	Ø	0	63	38	0.25
40 m	Ø	16	86	Ø	Ø	Ø	Ó	8	38	(03	0.75
50 m	3	10	86	Ø	Ø	Ø	Ø	0	10	8(0	0.75
60 m	3	10	810	Ø	8	Ø	V D	8	38	03	0.50
70 m	10	10	38	Ø	Ø	10	Ø	8	1(0	860	0.50
80 m	10	10	38	Ø	Ø	9	Ø	Ø	10	810	0.50
90 m	10	10	38	Ø	Ø	Ø	Ø	Ø	1(0	86	0.50

Robel Pole:

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
Ν	15	11	12	10	13		10	10	9	a
Ε	15	11	12	11	13	11	10	10	9	9
s	15	12	12	14	13	11	10		9	0)
W	15	12	12	14	13	11	()	1	9	9

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm