Daubenmire Datasheet

Date:

Observer:

Recorder:

Site: pre6 \_ tpre6a
Transect: 2 \_ \_

Round: 50m

Bare Leaf Butterfly Woody Litter Common Swamp WSG Forbs Milkweed CSG Depth (cm) Litter ramet # ramet # ramet # Species Ground (1) .50 86 0 0 m 86 00. 10 m 0 63 20 m 6 2,00 0 30 m 2.00 0 40 m 6 50 m 60 m 70 m 80 m 90 m

## Robel Pole:

Team lead: \_\_

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	3	7	8	4/	5					
Ε	3	6	4	3	7		100			
s	4	9	9	13	4					
W	5	9	10	7	9					

Categories: 0 - absent 1 - trace 3 - trace-5% 16 - 5%-25% 38 - 25%-50% 63 - 50%-75% 86 - 75%-95% 98 - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm