Daub...mire Datasheet Date:

Observer:

Recorder:

Team lead:

	n lead:	W56	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
					runei n	Tunior ii	- 8	O	16	63	0.5
0 m	3 8	0	86	0				0	36	38	0.25
10 m	38	0	63	0					16	(>	6.25
20 m	86	0	3 %	0			1000	70	38	38	0.25
30 m	63	0	63	0					38	3 %	6.25
0 m	38	Õ	86	0	-				_	38	0.15
50 m	34	0	86	0				C C C C C C C C C C C C C C C C C C C	38	0	~
60 m											
70 m											
80 m											

90 m

obel Po	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	7	4	Ц	3	7	5	\ /			
	1	1-7	//	1	7	5				
Е	6		U)	1-4	5	X			
S		\$	Ч	3	1-/-	5	1/\			
W	7	5	of .	3		1	/ \			

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm

arm1

tarm1b

2

60m

Site:

Round:

Transect: