Daubenmire Datasheet

Date: 7-16-18

Observer: N: -1 D + K 5.15

Recorder: 0.110

Team lead: Non C

> Site: pre3a

Transect: tpre3a

100m 2

Round:

	CS6	WS6	Forbs	Milkweed	Common	Swamp ramet #	Butterfly ramet #	Woody	Bare	Leaf	Litter Depth (cm)
0 m	38	0	R	P	Q	Q	P	0	6	8	0.5
10 m	Ø	0	38	9	0	0	9	8	_	2	2
20 m	16	Ø	98	8	9	Ø	0	2	nagariti-	7	2.0
30 m	16	Ø	98	D	0	0	2	0	3	2	7.5
40 m	0	0	98	0	0	0	B.	2			
50 m	16	\$	63	9	C.	0	Š	2	_	38	2
60 m	38	0	98	0	0	0	0	0		7	~(
70 m	3	0	63	0	0	Ø	D.	8		\(\infty\)	
80 m	a	0	98	0	Ø	0	0	0		0	2
90 m	w	D	38	0	0	0	9	Q			- 1
Robel Pole:	ole:							7		·	

i	5	
i	5	
(	0	
•	_	
•	O	
(	3	
ï	0	
:		

	S l6	m -5	Z	0 m
2 9	_	00	5	n 10 m
0	14	12	w	20 m
2]	16	15	10	30 m
15	16	15	(2)	40 a
0	16	16	51	50 m
3	8	1]	~	60 m
57	5	7	+	70 m
1	16	16	10	80 m
5	-+	ia	5	90 m

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm