Daub ... imire Datasheet

Date: Observer:

Recorder:

Team lead:

Site: arm2

100m

Round: Transect: tarm2a

CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf Litter
30 <u>3</u>	C	702	0	0	0	0	0	0	400
)	22	(0)3	0	0	C	0	0	3	810
. (200	5 () () (¥ ()	ナ	N	Q
0	N	800	0	C	0	C	0	U	O
C	0	00	0	0	0	0	0	O	308
C	10	3	الر		0	0	0	S	2
ادر	<u> </u>	870	0)	d	9	0	S	8(
20	- (000)	3)	7)	7	0	0)1	~
100	1/0	()° ()) (Э	V	K	Q	01	603
SE	- 5	10.2)(0	0)	0	0	0)	8
کور	0)(0	20) (C	C	3	C	\supset	57	20
	20 0 0 0 0 CS6		38 ws6	WS6 Forbs 38 63 38 86 16 78 16 38 16 38 16 38 16 38 16 38	WSG Forbs Milkweed 38 (03 0 38 8(0 0 10 98 0 10 38 8(0 0 10 38 0 10 58 0 10 58 0	WS6 Forbs Milkweed Common ramet # 20 (123 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	WSG Forbs Milkweed Common Swamp O	WSG Forbs Milkweed ramet # ram	WS6 Forbs Milkweed ramet # ramet # Species O

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	
z	6	2)	J	J	V	7		S	
ш	6	4	J	30 .	ىر	7	S	U	
S	()	H	2	00	بر	4	6	S	
€	5	2	0	ϖ)	N	J	U	

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm