Date: Recorder: Observer: Daubenmire Datasheet Team lead: 10-18 Site: Round: Transect: tprd1a 100m prd1 Ь

	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf	Litter Depth (cm)
0 m	77	0)/	0	0				0		\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Ochin (ciii)
10 m	38	50		0				0	Λ-	√) C	03
20 m	38	16	0)]	0				0	N	2	N)
30 m	38	63	0,1	0					_(200	20
40 m	X	39	28	0				0		500	
50 m	0)	56	W	\bigcirc				0		52) .
60 m	W	200								2	N N
70 m	22	W	W.							2	الم
80 m	X	5	_)(5	9
90 m	6	38	W	0						16	<i>S</i>
Robel Pole:	Pole:								(0.0	000

	z	т	S	€
0 m	-	()		S
10 m	0	7	7	6
20 m	0	9	(0	L
30 m	N	2	0	5
40 m	2	H	N	7
50 m	(7)	W		7
60 m	T	5	T	
70 m	二		匚	
80 m	7	7		
90 m				

WSG - warm season grasses CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground – includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm