Daubenmire Datasheet

Date:

Observer:

Recorder:

Site: app1

Transect: tapp1a

Round:

Team lead:

		(had		Common	Swamb	Butterfly	Woody	Bare	Leaf	Litter
	CS6	WS6	Forbs	Milkweed	ramet #	ramet #	ramet #	Species	Ground	Litter	Depth (cm)
0	9	38	N	O/	0	0	0	0	(S)	82	0,25
10 m	W X	N	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	\bigcirc	Q	0	0	0)	/(0	28	0.25
20 m	× 0	6	CC R	0	0	0	α	0)	(6)	\\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\	3,0
30 m	// 0	01/	3	0	0	0	0	0	16	63	9
40 m	63	0)	1(0	E	○	0	O	0	38	CN P	
50 m			1								
60 m											
70 m		=									
80 m											
90 m											
,											

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m
z		7	6	0	12				
m	_	A		7	12				
S	0	00	8	12	12				
₹	0			7	7				

Categories: $\underline{0}$ - absent $|\underline{1}$ - trace $|\underline{3}$ - trace-5% $|\underline{16}$ - 5%-25% $|\underline{38}$ - 25%-50% $|\underline{63}$ - 50%-75% $|\underline{86}$ - 75%-95% $|\underline{98}$ - 95%-100% CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm