Date: Daubenmire Datasheet

Observer:

Team lead:

Recorder:

Site:

Round:

100m

Transect: tbyd2a byd2

	•		1		Common	Swamp	Butterfly	Woody	Bare	Leaf	Litter
	CS6	WS6	rorbs	WIIKWeed	ramet #	ramet #	ramet #	Species	Ground	Litter	Depth (cm)
0	.N.	0	X	0	0	0	0	0	W	8/0	0.50
10 m	∞	0	6 W	0	0	0	0	0	10	03	0.25
20 m	0	0	8(0	0	0	0	C	0	1(2	63	0.25
	N	0	860	0	0	G	C	0	N	876	0.25
	N	0	2 K	0	Q	0	0	0	_	800	1.50
	W	9	200	0	Ó	0	0	0		20	0.50
	ا در	0	86	0	Q	0	0	0	N	8	0.25
70 m	100	0	S.	0	0	0	0	0	W	200	1.25
80	()	0	× 0	0	0	0	0	0	0) 1	67	1.00
	W	Q	2	0	0	0	0	C	38	80	0.25
	-										

Robel Pole:

	Z	æ	S	€
0 m	_	_	_	_
10 m	4	4		6
20 m	W	N	1	≪.
30 m	4	h	A	0
40 m	7	3	2	8
50 m	2	2	7	8
60 m	N	N	M	6
70 m	2	N	W	X
80 m	w	W	N	M
90 m	4	N	4	4

Categories: 0 - absent | 1 - trace | CSG - cool season grasses (including sedges) 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm