Daubenmire Datasheet

Date: 8-13-18

Observer: Service Recorder:

Team lead: Seth

Site: app1

Transect: tapp1a

3 50m

Round:

	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody	Bare	Leaf	Litter Depth (cm)
0 m	3 %	6	91	0	1		(5	36	300	0 25
10 m	38	16	88	Ď	1	(0	~ ;	20 6	R
20 m	500	0	6	0	1)))	- (200	n e
30 m	6,	0	5	0	,)	1	5	N-	0 =	1
40)	3 6)				(0	06	Ü
	100		C	C				0	L	6	° U
50 m	4				1	١	1				
60 m											
70 m											
3											
80 m											
90 m											
Robel Pole:	ole:										

	z	ш	S	€
0	رن	ىيا	a	4
10 m	A	8	0	0
20 m	0)	7	N	ىر
30 m	6	0	Н	6
40 m	6	6	a	32
50 m	1			\
60 m				
70 m				
80 m	\			
90 m				

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100% CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth – to nearest .25 cm